

Dr. Allan Shanin, DDS | 39 Broadway Suite 2115 | New York, NY 10006 | Tel: (212) 422-9229

Post-Operative Care for Implant and Bone Graft

- 1) **DO NOT** rinse for 24 hours. **Avoid** extremely hot food and liquids.
- 2) Medication:
 - 1. Antibiotics to be taken prior to appointment or as instructed.
 - Mild discomfort can be controlled with 2 Ibuprofen/Tylenol (or similar tablets) every 4-6 hours.
 Aspirin should NOT be used. If a PAIN RELIEVER has been prescribed, take only when necessary. If an antibiotic has been prescribed, take ALL medication to help prevent infection.
- AVOID alcoholic beverages during the post-operative period, especially if you take medication.
- 4) **AVOID** smoking during the post-operative period.
- 5) To minimize swelling which may occur, apply an ice pack on your face next to the treated area. Leave it in place for 10 minutes and then remove it for 10 minutes. Repeat this routine. Do not use the ice pack (or application of) after 24 hours following the surgery. If swelling is severe and/or you feel feverish, please call the office.
- 6) **Brush and floss** the teeth well, but not in the treated area. Rinse with Listerine or prescribed rinse 2 times per day.
- 7) AVOID excessive exercise (jogging, swimming, tennis, etc.) for 72 hours.
- 8) Eat on the side OPPOSITE the sutures. Eating suggestion: fish, eggs, soup, cheese, chopped meat, baby food, cooked cereal, cooked vegetables, jell-O, soft fruits, and blended foods.
- 9) If bleeding occurs, DO NOT RINSE! Moisten a gauze pad or fresh tea bag with warm water. Place it over the area and hold firmly for 10-20 minutes without interruption. Repeat if necessary.
- 10) If you have guestions or concerns, please call our office at (212) 422-9229.