

The Singleton Kind Calculus

February 2, 2017

Abstract

In this section we develop the singleton kind calculus. A singleton kind $S(c)$ is the kind of all constructors that are equivalent to c . The addition of these new kinds will be useful to explain module signatures later on.

1 Syntax

The singleton kind calculus is built on top of souped-up F_ω .

$$\begin{aligned} k &::= \mathbf{type} \mid k \rightarrow k \mid k * k \mid S(c) \mid \Pi\alpha : k. k \mid \Sigma\alpha : k. k \\ c &::= \alpha \mid c \rightarrow c \mid \forall\alpha : k. c \mid \lambda\alpha : k. c \mid c \ c \mid \langle c, c \rangle \mid \pi_1 \ c \mid \pi_2 \ c \\ e &::= x \mid \lambda x : c. e \mid e \ e \mid \Lambda\alpha : k. e \mid e[c] \\ \Gamma &::= \cdot \mid \Gamma, x : c \mid \Gamma, \alpha : k \end{aligned}$$

2 Motivation

Consider the following ML signature.

```
sig
  type t
  type 'a u
  type ('a, 'b) v
  type w = int
end
```

The first three types can be assigned kinds in F_ω in a straight forward way.

```
t : type
u : type → type
v : type × type → type
```

But how do we kind w ? Remember, `int` is not a kind, so it doesn't make sense to say $w : \mathbf{int}$. But it's not quite right to say $w : \mathbf{type}$ either, because

\mathbf{w} cannot stand for arbitrary types. We therefore write $\mathbf{w} : S(\mathbf{int})$: $S(\mathbf{int})$ is the kind containing exactly \mathbf{int} and all those types equivalent to \mathbf{int} , such as $(\lambda\alpha : \mathbf{type}. \alpha) \mathbf{int}$. The other new kind constructs, $\Pi\alpha : k. k$ and $\Sigma\alpha : k. k$ (which are called dependent function spaces and dependent sums respectively), exist to solve the analogous problem for kinding assignments to polymorphic types in signatures. Specifically, consider this example:

```
sig
  type 'a t = 'a list
end
```

We can say that

$$\mathbf{t} : \Pi\alpha : \mathbf{type}. \mathcal{S}(\alpha)$$

Σ kinds are used to kind types that are assigned to abstract types. In the signature

```
type t
type s = t
```

we say that

$$\begin{aligned} \mathbf{t} &: \mathbf{type} \\ \mathbf{s} &: \Sigma\alpha : \mathbf{type}. \mathcal{S}(\alpha) \end{aligned}$$

3 Definitions

In this section the following judgements will be defined.

Judgement	Description
$\Gamma \vdash k : \mathbf{kind}$	k is a kind
$\Gamma \vdash k \equiv k' : \mathbf{kind}$	kind equivalence
$\Gamma \vdash k \leq k'$	subkinding
$\Gamma \vdash c : k$	c has kind k
$\Gamma \vdash c \equiv c' : k$	constructor equivalence
$\Gamma \vdash e : \tau$	e has type τ

A complete list would also include the judgement $\Gamma \vdash \tau : \mathbf{type}$ but these rules are exactly the same as in F_ω so we will omit them.

3.1 Well-Formed Kinds

$$\begin{array}{c}
\text{3A} \quad \frac{}{\Gamma \vdash \tau : \text{kind}} \quad \text{3B} \quad \frac{\Gamma \vdash c : \tau}{\Gamma \vdash S(c) : \text{kind}} \quad \text{3C} \quad \frac{\Gamma \vdash k_1 : \text{kind} \quad \Gamma, k_1 : \text{kind} \vdash k_2 : \text{kind}}{\Gamma \vdash \Pi \alpha : k_1. k_2 : \text{kind}} \\
\\
\text{3D} \quad \frac{\Gamma \vdash k_1 : \text{kind} \quad \Gamma, k_1 : \text{kind} \vdash k_2 : \text{kind}}{\Gamma \vdash \Sigma \alpha : k_1. k_2 : \text{kind}}
\end{array}$$

3.2 Definitional Equality of Kinds

$$\begin{array}{c}
\text{3E} \quad \frac{\Gamma \vdash k : \text{kind}}{\Gamma \vdash k \equiv k : \text{kind}} \quad \text{3F} \quad \frac{\Gamma \vdash k_1 \equiv k_2 : \text{kind}}{\Gamma \vdash k_2 \equiv k_1 : \text{kind}} \\
\\
\text{3G} \quad \frac{\Gamma \vdash k_1 \equiv k_2 : \text{kind} \quad \Gamma \vdash k_2 \equiv k_3 : \text{kind}}{\Gamma \vdash k_1 \equiv k_3 : \text{kind}} \quad \text{3H} \quad \frac{\Gamma \vdash c \equiv c' : \text{type}}{\Gamma \vdash S(c) \equiv S(c') : \text{kind}} \\
\\
\text{3I} \quad \frac{\Gamma \vdash k_1 \equiv k'_1 : \text{kind} \quad \Gamma, \alpha : k_1 \vdash k_2 \equiv k'_2 : \text{kind}}{\Gamma \vdash \Pi \alpha : k_1. k_2 \equiv \Pi \alpha : k'_1. k_2 : \text{kind}} \\
\\
\text{3J} \quad \frac{\Gamma \vdash k_1 \equiv k'_1 : \text{kind} \quad \Gamma, \alpha : k_1 \vdash k_2 \equiv k'_2 : \text{kind}}{\Gamma \vdash \Sigma \alpha : k_1. k_2 \equiv \Sigma \alpha : k'_1. k_2 : \text{kind}}
\end{array}$$

3.3 Kind Membership

This judgement defines which constructors belong to a given kind.

$$\frac{3K \quad \alpha : k \in \Gamma}{\Gamma \vdash \alpha : k}$$

$$\frac{3L \quad \Gamma \vdash c_1 : \mathbf{type} \quad \Gamma \vdash c_2 : \mathbf{type}}{\Gamma \vdash c_1 \rightarrow c_2 : \mathbf{type}}$$

$$\frac{3M \quad \Gamma \vdash k : \mathbf{kind} \quad \Gamma, \alpha : k \vdash c : \mathbf{type}}{\Gamma \vdash \forall \alpha : k. c : \mathbf{type}}$$

$$\frac{3N \quad \Gamma \vdash k_1 : \mathbf{kind} \quad \Gamma, \alpha : k_1 \vdash c : k_2}{\Gamma \vdash \lambda \alpha : k_1. c : \Pi \alpha : k_1. k_2}$$

$$\frac{3O \quad \Gamma \vdash c_1 : \Pi \alpha : k. k' \quad \Gamma \vdash c_2 : k}{\Gamma \vdash c_1 c_2 : [c_2/\alpha]k'}$$

$$\frac{3P \quad \Gamma \vdash c_1 : k_1 \quad \Gamma \vdash c_2 : [c_1/\alpha]k_2 \quad \Gamma, \alpha : k_1 \vdash k_2 : \mathbf{kind}}{\Gamma \vdash \langle c_1, c_2 \rangle : \Sigma \alpha : k_1. k_2}$$

$$\frac{3Q \quad \Gamma \vdash c : \Sigma \alpha : k_1. k_2}{\Gamma \vdash \pi_1 c : k_1}$$

$$\frac{3R \quad \Gamma \vdash c : \Sigma \alpha : k_1. k_2}{\Gamma \vdash \pi_2 c : [\pi_1 c/\alpha]k_2}$$

$$\frac{3S \quad \Gamma \vdash c : \mathbf{type}}{\Gamma \vdash c : S(c)}$$

Notice that even though $\Sigma \alpha : k_1. k_2$ is called a dependent sum, it behaves like a product. To avoid the obvious naming confusion here, we will try our best to call $\Sigma \alpha : k_1. k_2$ a "dependent sum" and $\Pi \alpha : k_1. k_2$ a "dependent function space" or sometimes just a "dependent function" because the former is a bit of a mouthful.

3.4 Subkinding

$$\frac{3_T \quad \Gamma \vdash k : \mathbf{kind}}{\Gamma \vdash k \leq k}$$

$$\frac{3_U \quad \Gamma \vdash k_1 \leq k_2 \quad \Gamma \vdash k_2 \leq k_3}{\Gamma \vdash k_1 \leq k_3}$$

$$\frac{3_V \quad \Gamma \vdash c : \mathbf{type}}{\Gamma \vdash S(c) \leq \mathbf{type}}$$

$$\frac{3_W \quad \Gamma \vdash c \equiv c' : \mathbf{type}}{\Gamma \vdash S(c) \leq S(c') : \mathbf{type}}$$

$$\frac{3_X \quad \Gamma \vdash k'_1 \leq k_1 \quad \Gamma \alpha : k_1 \vdash k_2 : \mathbf{kind} \quad \Gamma, \alpha : k'_1 \vdash k_2 \leq k'_2}{\Gamma \vdash \Pi \alpha : k_1. k_2 \leq \Pi \alpha : k'_1. k'_2}$$

$$\frac{3_Y \quad \Gamma \vdash k_1 \leq k'_1 \quad \Gamma, \alpha : k_1 \vdash k_2 \leq k'_2 \quad \Gamma, \alpha : k'_1 \vdash k'_2 : \mathbf{kind}}{\Gamma \vdash \Sigma \alpha : k_1. k_2 \leq \Sigma \alpha : k'_1. k'_2}$$

$$\frac{3_Z \quad \Gamma \vdash c : k \quad \Gamma \vdash k \leq k'}{\Gamma \vdash c : k'}$$

$$\frac{3_{AA} \quad \Gamma \vdash k \equiv k'}{\Gamma \vdash k \leq k'}$$

3.5 Definitional Equality of Constructors

$$\frac{3_{AB} \quad \Gamma \vdash c : k}{\Gamma \vdash c \equiv c : k}$$

$$\frac{3_{AC} \quad \Gamma \vdash c_1 \equiv c_2 : k}{\Gamma \vdash c_2 \equiv c_1 : k}$$

$$\frac{3_{AD} \quad \Gamma \vdash c_2 : k \quad \Gamma, \alpha : k \vdash c_1 : k'}{\Gamma \vdash (\lambda \alpha : k. c_1) c_2 \equiv [c_2/\alpha]c_1 : [c_2/\alpha]k'}$$

$$\frac{3_{AE} \quad \Gamma \vdash c_1 : k_1 \quad \Gamma \vdash c_2 : k_2}{\Gamma \vdash \pi_1 \langle c_1, c_2 \rangle \equiv c_1}$$

$$\frac{3_{AF} \quad \Gamma \vdash c_1 : k_1 \quad \Gamma \vdash c_2 : k_2}{\Gamma \vdash \pi_2 \langle c_1, c_2 \rangle \equiv c_2}$$

$$\frac{3_{AG} \quad \Gamma \vdash c_1 \equiv c_2 : k \quad \Gamma \vdash k \leq k'}{\Gamma \vdash c_1 \equiv c_2 : k'}$$

$$\frac{3_{AH} \quad \Gamma \vdash c \equiv c' : \mathbf{type}}{\Gamma \vdash c \equiv c' : S(c)}$$

$$\frac{3_{AI} \quad \Gamma \vdash c : S(c')}{\Gamma \vdash c \equiv c' : \mathbf{type}}$$

$$\frac{3AJ}{\Gamma \vdash k_1 \equiv k_2 : \mathbf{kind} \quad \Gamma, \alpha : k_1 \vdash c \equiv c' : k_2} \quad \Gamma \vdash \lambda \alpha : k_1. c \equiv \lambda \alpha : k'_1. c' : \Pi \alpha : k_1. k_2$$

$$\frac{3AK}{\Gamma \vdash c \equiv c' : \Pi \alpha : k. k' \quad \Gamma \vdash c_2 \equiv c'_2 : k} \quad \Gamma \vdash c_1 c_2 \equiv c'_1 c'_2 : [c_2/\alpha]k'$$

$$\frac{3AL}{\Gamma \vdash c_1 \equiv c'_1 : k_1 \quad \Gamma \vdash c_2 \equiv c'_2 : [c_1/\alpha]k_2 \quad \Gamma, \alpha : k_1 \vdash k_2 : \mathbf{kind}} \quad \Gamma \vdash \langle c_1, c_2 \rangle \equiv \langle c'_1, c'_2 \rangle : \Sigma \alpha : k_1. k_2$$

$$\frac{3AM}{\Gamma \vdash c \equiv c' : \Sigma \alpha : k_1. k_2} \quad \Gamma \vdash \pi_1 c \equiv \pi_1 c' : k_1$$

$$\frac{3AN}{\Gamma \vdash c \equiv c' : \Sigma \alpha : k_1. k_2} \quad \Gamma \vdash \pi_2 c \equiv \pi_2 c' : [\pi_1 c/\alpha]k_2$$

$$\frac{3AN}{\Gamma \vdash c_1 \equiv c'_1 : \mathbf{type} \quad \Gamma \vdash c_2 \equiv c'_2 : \mathbf{type}} \quad \Gamma \vdash c_1 \rightarrow c_2 \equiv c'_1 \rightarrow c'_2 : \mathbf{type}$$

$$\frac{3AO}{\Gamma \vdash k \equiv k' : \mathbf{kind} \quad \Gamma, \alpha : k \vdash c \equiv c' : \mathbf{type}} \quad \Gamma \vdash \forall \alpha : k. c \equiv \forall \alpha : k'. c' : \mathbf{type}$$

$$\frac{3AP}{\Gamma, \alpha : k_1 \vdash c \alpha \equiv c' \alpha : k_2 \quad \Gamma \vdash c : \Pi \alpha : k_1. k_2 \quad \Gamma \vdash c' : \Pi \alpha : k_1. k_2''} \quad \Gamma \vdash c \equiv c' : \Pi \alpha : k_1. k_2$$

$$\frac{3AQ}{\Gamma \vdash \pi_1 c \equiv \pi_1 c' \quad \Gamma, \alpha : k_1 \vdash k_2 : \mathbf{kind} \quad \Gamma \vdash \pi_2 c \equiv \pi_2 c' : [\pi_1 c/\alpha]k_2} \quad \Gamma \vdash c \equiv c' : \Sigma \alpha : k_1. k_2$$