

The Singleton Kind Calculus

February 2, 2017

Abstract

In this section we develop the singleton kind calculus. A singleton kind $S(c)$ is the kind of all constructors that are equivalent to c . The addition of these new kinds will be useful to explain module signatures later on.

1 Syntax

The singleton kind calculus is built on top of souped-up F_ω .

$$\begin{aligned} k &::= \mathbf{type} \mid k \rightarrow k \mid k * k \mid S(c) \mid \Pi\alpha : k. k \mid \Sigma\alpha : k. k \\ c &::= \alpha \mid c \rightarrow c \mid \forall\alpha : k. c \mid \lambda\alpha : k. c \mid c \ c \mid \langle c, c \rangle \mid \pi_1 \ c \mid \pi_2 \ c \\ e &::= x \mid \lambda x : c. e \mid e \ e \mid \Lambda\alpha : k. e \mid e[c] \\ \Gamma &::= \cdot \mid \Gamma, x : c \mid \Gamma, \alpha : k \end{aligned}$$

2 Motivation

Consider the following ML signature.

```
sig
  type t
  type 'a u
  type ('a, 'b) v
  type w = int
end
```

The first three types can be assigned kinds in F_ω in a straight forward way.

```
t : type
u : type → type
v : type × type → type
```

But how do we kind w ? Remember, `int` is not a kind, so it doesn't make sense to say $w : \mathbf{int}$. But it's not quite right to say $w : \mathbf{type}$ either, because

\mathbf{w} cannot stand for arbitrary types. We therefore write $\mathbf{w} : S(\mathbf{int})$: $S(\mathbf{int})$ is the kind containing exactly \mathbf{int} and all those types equivalent to \mathbf{int} , such as $(\lambda\alpha : \mathbf{type}. \alpha) \mathbf{int}$. The other new kind constructs, $\Pi\alpha : k. k$ and $\Sigma\alpha : k. k$ (which are called dependent function spaces and dependent sums respectively), exist to solve the analogous problem for kinding assignments to polymorphic types in signatures:

```
sig
  type 'a t = 'a list
end
```

This will become more clear once the rules are enumerated.

3 Definitions

In this section the following judgements will be defined.

Judgement	Description
$\Gamma \vdash k : \mathbf{kind}$	k is a kind
$\Gamma \vdash k \equiv k' : \mathbf{kind}$	kind equivalence
$\Gamma \vdash k \leq k'$	subkinding
$\Gamma \vdash c : k$	c has kind k
$\Gamma \vdash c \equiv c' : k$	constructor equivalence
$\Gamma \vdash e : \tau$	e has type τ

A complete list would also include the judgement $\Gamma \vdash \tau : \mathbf{type}$ but these rules are exactly the same as in F_ω so we will omit them. Beginning with the rules for well-formed kinds:

3A $\frac{}{\Gamma \vdash \tau : \mathbf{kind}}$	3B $\frac{\Gamma \vdash c : \tau}{\Gamma \vdash S(c) : \mathbf{kind}}$	3C $\frac{\Gamma \vdash k_1 : \mathbf{kind} \quad \Gamma, k_1 : \mathbf{kind} \vdash k_2 : \mathbf{kind}}{\Gamma \vdash \Pi\alpha : k_1. k_2 : \mathbf{kind}}$
	3D $\frac{\Gamma \vdash k_1 : \mathbf{kind} \quad \Gamma, k_1 : \mathbf{kind} \vdash k_2 : \mathbf{kind}}{\Gamma \vdash \Sigma\alpha : k_1. k_2 : \mathbf{kind}}$	

Definitional equality of kinds:

$$\frac{3E \quad \Gamma \vdash k : \mathbf{kind}}{\Gamma \vdash k \equiv k : \mathbf{kind}}$$

$$\frac{3F \quad \Gamma \vdash k_1 \equiv k_2 : \mathbf{kind}}{\Gamma \vdash k_2 \equiv k_1 : \mathbf{kind}}$$

$$\frac{3G \quad \Gamma \vdash k_1 \equiv k_2 : \mathbf{kind} \quad \Gamma \vdash k_2 \equiv k_3 : \mathbf{kind}}{\Gamma \vdash k_1 \equiv k_3 : \mathbf{kind}}$$

$$\frac{3H \quad \Gamma \vdash c \equiv c' : \mathbf{type}}{\Gamma \vdash S(c) \equiv S(c') : \mathbf{kind}}$$

$$\frac{3I \quad \Gamma \vdash k_1 \equiv k'_1 : \mathbf{kind} \quad \Gamma, \alpha : k_1 \vdash k_2 \equiv k'_2 : \mathbf{kind}}{\Gamma \vdash \Pi \alpha : k_1. k_2 \equiv \Pi \alpha : k'_1. k_2 : \mathbf{kind}}$$

$$\frac{3J \quad \Gamma \vdash k_1 \equiv k'_1 : \mathbf{kind} \quad \Gamma, \alpha : k_1 \vdash k_2 \equiv k'_2 : \mathbf{kind}}{\Gamma \vdash \Sigma \alpha : k_1. k_2 \equiv \Sigma \alpha : k'_1. k_2 : \mathbf{kind}}$$

Kind membership — which constructors belong to a given kind:

$$\frac{3K \quad \alpha : k \in \Gamma}{\Gamma \vdash \alpha : k}$$

$$\frac{3L \quad \Gamma \vdash c_1 : \mathbf{type} \quad \Gamma \vdash c_2 : \mathbf{type}}{\Gamma \vdash c_1 \rightarrow c_2 : \mathbf{type}}$$

$$\frac{3M \quad \Gamma \vdash k : \mathbf{kind} \quad \Gamma, \alpha : k \vdash c : \mathbf{type}}{\Gamma \vdash \forall \alpha : k. c : \mathbf{type}}$$

$$\frac{3N \quad \Gamma \vdash k_1 : \mathbf{kind} \quad \Gamma, \alpha : k_1 \vdash c : k_2}{\Gamma \vdash \lambda \alpha : k_1. c : \Pi \alpha : k_1. k_2}$$

$$\frac{3O \quad \Gamma \vdash c_1 : \Pi \alpha : k. k' \quad \Gamma \vdash c_2 : k}{\Gamma \vdash c_1 c_2 : [c_2/\alpha]k'}$$

$$\frac{3P \quad \Gamma \vdash c_1 : k_1 \quad \Gamma \vdash c_2 : [c_1/\alpha]k_2 \quad \Gamma, \alpha : k_1 \vdash k_2 : \mathbf{kind}}{\Gamma \vdash \langle c_1, c_2 \rangle : \Sigma \alpha : k_1. k_2}$$

$$\frac{3Q \quad \Gamma \vdash c : \Sigma \alpha : k_1. k_2}{\Gamma \vdash \pi_1 c : k_1}$$

$$\frac{3R \quad \Gamma \vdash c : \Sigma \alpha : k_1. k_2}{\Gamma \vdash \pi_2 c : [\pi_1 c/\alpha]k_2}$$

$$\frac{3S \quad \Gamma \vdash c : \mathbf{type}}{\Gamma \vdash c : S(c)}$$

Notice that even though $\Sigma \alpha : k_1. k_2$ is called a dependent sum, it behaves like a product. To avoid the obvious naming confusion here, we will try our best to call $\Sigma \alpha : k_1. k_2$ a "dependent sum" and $\Pi \alpha : k_1. k_2$ a "dependent function space" or sometimes just a "dependent function" because the former is a bit of a mouthful.