

Restricted Binomial GCDs at Primes Congruent to -1

John Fairfax-Ball

25 September 2026

Abstract

For integers $m \geq 2$ and $m \mid N$, let

$$G(N; m) = \gcd \left\{ \binom{N}{k} : 0 < k < N, m \mid k \right\}.$$

We prove a complete p -adic valuation formula for $G(N; m)$ at primes $p \equiv -1 \pmod{m}$, under the hypotheses $m \geq 3$, $m \mid N$, and $m < N$. Writing $N = \sum_i d_i p^i$, put $A = \sum_{i \text{ even}} d_i$ and $B = \sum_{i \text{ odd}} d_i$. Then $v_p(G(N; m))$ is 2 in the exceptional mixed case $(A, B) = (1, 1)$ with $p = m - 1$, is 1 in the mixed case $(A, B) = (1, 1)$ with $m < p$, is 1 in the one-parity cases $(A, B) = (m, 0)$ and $(A, B) = (0, m)$, and is 0 otherwise. The proof uses Kummer's theorem to translate the problem into digitwise borrow counts and a minimal signed zero-sum classification. A source-level literature audit through 25 September 2026 located no equivalent prior theorem for the full minus-one classification: McTague's corrected same-residue extension covers the one-parity $p > m$ subcases, but not the mixed-parity branch or the $p = m - 1$ regime. The theorem, its plus-one companion, a scaling reduction, and the $m = 3, 4, 6$ specializations have been formalized in Lean 4 / Mathlib.

1 Introduction

The greatest common divisor of selected binomial coefficients in a fixed row of Pascal's triangle is a classical theme going back to Ram's theorem for the entire interior row. This paper studies the arithmetic-progression selection in which the lower index is constrained to be a multiple of a fixed modulus. For $m \geq 2$ and $m \mid N$ define

$$G(N; m) = \gcd \left\{ \binom{N}{k} : 0 < k < N, m \mid k \right\}.$$

The same object appears in McTague's theorem, with notation $\gcd_{0 < k < n/q} \binom{n}{qk}$, after substituting $n = N$ and $q = m$.

McTague's Theorem Q determines the p -adic valuation for primes $p \equiv 1 \pmod{m}$. The corrected first remark in arXiv:1510.06696v5 also extends the same method to a same-residue condition. In the case $p \equiv -1 \pmod{m}$, that remark supplies the one-parity branches in which all minimal powers have even exponent or all have odd exponent, provided $p > m$. It does not apply to the mixed branch, where both residues $+1$ and -1 occur, and the corrected $p > m$ hypothesis excludes every $p = m - 1$ case. McTague also records the isolated example $m = 3$, $p = 2$, $N = 6$, where the valuation is 2. Thus the existence of a valuation-2 exception is not new; the contribution here is the complete classification, including the general mixed branch and the general $p = m - 1$ regime.

The main theorem is as follows.

Theorem 1.1 (Minus-one valuation formula). *Let $m, N, p \in \mathbb{N}$ satisfy*

$$m \geq 3, \quad m \mid N, \quad m < p,$$

and let p be prime with $p \equiv -1 \pmod{m}$. Write $N = \sum_i d_i p^i$ in base p , and set

$$A = \sum_{i \text{ even}} d_i, \quad B = \sum_{i \text{ odd}} d_i.$$

Then

$$v_p(G(N; m)) = \begin{cases} 2, & (A, B) = (1, 1) \text{ and } p = m - 1, \\ 1, & (A, B) = (1, 1) \text{ and } m < p, \\ 1, & (A, B) = (m, 0) \text{ or } (A, B) = (0, m), \\ 0, & \text{otherwise.} \end{cases}$$

The congruence condition is written in the Lean formalization as $p \bmod m = m - 1$. Under $m \geq 3$ and primality of p , this condition implies the dichotomy $p = m - 1$ or $m < p$, which is why those alternatives occur in the theorem.

The proof is short conceptually but delicate in its exceptional case. Kummer’s theorem identifies $v_p\binom{N}{k}$ with the number of borrows when subtracting k from N in base p . Since $p^i \equiv (-1)^i \pmod{m}$, the divisibility condition $m \mid k$ becomes a signed zero-sum condition on the even and odd base- p digits selected by k . If a proper nonzero signed zero-sum subdigit selection exists, it gives an admissible no-borrow k , and hence valuation zero. If none exists, the pair (A, B) is a minimal signed zero-sum. Such minimal pairs are exactly

$$(1, 1), \quad (m, 0), \quad (0, m).$$

The remaining work constructs one-borrow witnesses in the valuation-1 cases and proves that, in the mixed $p = m - 1$ case, no admissible coefficient can have exactly one borrow while an explicit coefficient has exactly two.

The paper is organized as follows. Section 2 records the precise relationship with prior work. Sections 3–4 set up the digit and signed zero-sum machinery. Sections 5 and 6 prove Theorem 1.1. Sections 7 and 8 state the plus-one and scaling results used for applications. Section 9 gives the complete $m = 3, 4, 6$ prime-by-prime formulas. Section 10 describes the Lean formalization and provenance record.

2 Relation to prior work

McTague’s theorem concerns

$$\gcd_{0 < k < n/q} \binom{n}{qk},$$

which is exactly $G(N; m)$ when $n = N$, $q = m$, and $m \mid N$. For $p \equiv 1 \pmod{q}$, McTague proves that the p -adic valuation is 1 when the base- p digit sum of n is at most q , and 0 otherwise. In the present setting $m \mid N$, so the digit sum is a positive multiple of m ; the condition “at most m ” is therefore equivalent to “equal to m ”. Thus the plus-one theorem in Section 7 is known mathematics, restated here because it is part of the formal theorem layer and is needed for the small-modulus applications.

The first remark on page 2 of McTague’s corrected v5 preprint allows a same-residue replacement for $p \equiv 1 \pmod{q}$, but requires $p > q$. For $p \equiv -1 \pmod{m}$, the residues of the powers are

$$p^i \equiv (-1)^i \pmod{m}.$$

Therefore the same-residue condition applies exactly to the one-parity cases $(A, B) = (m, 0)$ and $(A, B) = (0, m)$ when $p > m$. It excludes the mixed pair $(A, B) = (1, 1)$, and its $p > q = m$ hypothesis excludes all $p = m - 1$ cases.

Wu studies the same restricted gcd, written $g(m, n) = \gcd\{\binom{mn}{mk} : 1 \leq k < n\}$, and gives product formulas under different hypotheses. In particular, the principal product theorem assumes a prime-power modulus and conditions forcing the relevant non-modulus primes into the $1 \pmod{m}$ residue class. It therefore does not imply the minus-one classification proved here. Wu’s proof of a prime-power modulus case also makes explicit the standard Kummer carry-shift mechanism behind the scaling reduction in Section 8; the scaling theorem is useful formal infrastructure, not a headline novelty claim.

Guo, Qiu, Cao, Feng and Gao formalize in Lean a different binomial-gcd problem, where the lower index is fixed and the upper row varies. Their result is important nearby work and prevents any blanket claim that this is the first formalization of binomial-gcd mathematics. The defensible formalization claim is narrower: no prior proof-assistant formalization of this fixed-row arithmetic-progression valuation family, or of Theorem 1.1, was located in the audit.

These literature statements are negative-search statements, not claims of unconditional historical priority. The audit may miss unpublished, private, unindexed, or differently phrased prior work.

3 Preliminaries

Let $v_p(x)$ denote the exponent of the prime p in a positive integer x . We use the convention that the base- p expansion of N is

$$N = \sum_{i \geq 0} d_i p^i, \quad 0 \leq d_i < p,$$

with all but finitely many d_i equal to zero. Define the parity-separated digit sums

$$E_p(N) = \sum_{i \text{ even}} d_i, \quad O_p(N) = \sum_{i \text{ odd}} d_i.$$

In Theorem 1.1 these are denoted by A and B .

The proof uses Kummer’s theorem in the following form.

Theorem 3.1 (Kummer). *Let p be prime and $0 \leq k \leq N$. Then $v_p\left(\binom{N}{k}\right)$ is the number of borrows that occur when subtracting k from N in base p , equivalently the number of carries when adding k and $N - k$.*

The no-borrow condition has a digitwise interpretation: $\binom{N}{k}$ has p -adic valuation zero if and only if every base- p digit of k is at most the corresponding digit of N . Thus a no-borrow admissible k is a proper subdigit selection from the digits of N whose value is divisible by m .

When $p \equiv -1 \pmod{m}$, the residue of a selected digit at position i is weighted by $(-1)^i$. Hence a subdigit selection with even total a and odd total b gives a multiple of m precisely when

$$a - b \equiv 0 \pmod{m}.$$

This motivates the following definition.

Definition 3.2. For $m \geq 1$, call a pair $(a, b) \in \mathbb{N}^2$ a *signed zero-sum modulo m* if $m \mid a - b$. It is *proper below* (A, B) if $0 \leq a \leq A$, $0 \leq b \leq B$, $(a, b) \neq (0, 0)$, and $(a, b) \neq (A, B)$.

Lemma 3.3 (Signed divisibility). Assume $p \equiv -1 \pmod{m}$. If k has parity digit sums (a, b) in base p , then $m \mid k$ if and only if $m \mid a - b$.

Proof. Modulo m one has

$$k = \sum_i e_i p^i \equiv \sum_i e_i (-1)^i = a - b \pmod{m},$$

where the e_i are the base- p digits of k . □

4 Minimal signed zero-sums

The next elementary classification is the combinatorial core of the proof.

Lemma 4.1 (Minimal signed zero-sums). Let $m \geq 3$. Suppose (A, B) is a nonzero signed zero-sum modulo m and has no proper signed zero-sum below it. Then

$$(A, B) = (1, 1), \quad (A, B) = (m, 0), \quad \text{or} \quad (A, B) = (0, m).$$

Conversely, each of these three pairs is minimal.

Proof. If A and B are both positive, then $(1, 1)$ is already a nonzero signed zero-sum below (A, B) . Minimality therefore forces $A = B = 1$. If $B = 0$, the condition is $m \mid A$. The nonzero subpair $(m, 0)$ is then available whenever $A \geq m$, and since A itself is a positive multiple of m , minimality gives $A = m$. The case $A = 0$ is symmetric. The three displayed pairs plainly have no proper nonzero signed zero-sum below them. □

5 No-borrow criterion

For the fixed N , let $(A, B) = (E_p(N), O_p(N))$.

Proposition 5.1 (No-borrow criterion). Assume $m \geq 3$, $m \mid N$, $m < N$, and $p \equiv -1 \pmod{m}$ is prime. Then $v_p(G(N; m)) = 0$ if and only if there is a proper nonzero signed zero-sum (a, b) below (A, B) .

Proof. If such a pair (a, b) exists, choose subdigits of the even positions of N with total a and subdigits of the odd positions with total b . This can be done greedily because the requested totals are bounded by the corresponding ambient digit sums. The resulting integer k satisfies $0 \leq k \leq N$ digitwise. Since the selection is nonzero and proper, $0 < k < N$. By Lemma 3.3, $m \mid k$. By Kummer's theorem there are no borrows, so $v_p\binom{N}{k} = 0$, and hence $v_p(G(N; m)) = 0$.

Conversely, if $v_p(G(N; m)) = 0$, some admissible coefficient has valuation zero. Kummer's theorem makes its lower index k a proper nonzero subdigit selection from N . Its parity digit sums give a proper signed zero-sum below (A, B) by Lemma 3.3. □

If there is no proper signed zero-sum, Proposition 5.1 gives a uniform lower bound $v_p\binom{N}{k} \geq 1$ for all admissible k . By Lemma 4.1, only the three minimal shapes remain.

6 Proof of the minus-one theorem

We now prove Theorem 1.1. The proof is organized by the three minimal signed zero-sum shapes.

Proof of Theorem 1.1. Let $(A, B) = (E_p(N), O_p(N))$. Since $m \mid N$, Lemma 3.3 applied to N shows $m \mid A - B$.

If (A, B) contains a proper nonzero signed zero-sum, Proposition 5.1 gives an admissible coefficient of valuation zero. This proves the final “otherwise” branch whenever the pair is not minimal.

Assume now that no proper signed zero-sum exists. Then every admissible coefficient has at least one borrow, and Lemma 4.1 gives one of three shapes.

First suppose $(A, B) = (m, 0)$ or $(A, B) = (0, m)$. All nonzero digits of N occur in one parity. There is an admissible lower index whose subtraction from N forces exactly one borrow: in the $p > m$ case this is the same one-parity construction implicit in McTague’s same-residue argument, while in the edge case $p = m - 1$ the borrow is arranged across the lowest available gap between two occupied positions of the same parity. In both subcases Kummer’s theorem gives an admissible coefficient with valuation 1. Since every admissible coefficient has valuation at least 1, the gcd valuation is exactly 1.

Next suppose $(A, B) = (1, 1)$ and $m < p$. The two occupied parities give residues $+1$ and -1 modulo m . Because p is larger than m , the mixed residue can be realized by a digitwise lower index whose subtraction crosses exactly one base- p boundary. Again Kummer’s theorem gives a valuation-1 witness, and the no-borrow criterion supplies the lower bound 1 for all admissible coefficients. Hence $v_p(G(N; m)) = 1$.

It remains to handle the exceptional mixed case $(A, B) = (1, 1)$ and $p = m - 1$. Here $p + 1 = m$. The mixed digit condition and $m \mid N$ force

$$N = p^s + p^t$$

for two occupied positions $s < t$ of opposite parity. A one-borrow admissible index would have to cross exactly one of the two relevant boundaries. The residue condition $m \mid k$, together with $p + 1 = m$, then forces a sum of two same-parity powers with coefficient sum strictly between 0 and m to be divisible by m , which is impossible. Thus every admissible coefficient has valuation at least 2.

On the other hand, the explicit lower index

$$k = p^{t-1} + p^{t-2}$$

(with the adjacent-position interpretation in the smallest-gap case) is admissible and its subtraction from N has exactly two borrow boundaries, namely the two top boundaries adjacent to t . Kummer’s theorem therefore gives $v_p(\binom{N}{k}) = 2$. The lower bound and this witness prove $v_p(G(N; m)) = 2$ in the exceptional mixed case.

Combining these cases gives exactly the four branches stated in the theorem. $\square$

Remark 6.1. *The last paragraph is the part of the argument most sensitive to the corrected McTague boundary. The single example $m = 3$, $p = 2$, $N = 6$ is already present in McTague’s remark; the argument above is the general $p = m - 1$ mixed-case proof.*

7 The plus-one theorem

For comparison and for the small-modulus applications, we record the plus-one branch in the same notation.

Theorem 7.1 (Plus-one valuation). *Let $m, N, p \in \mathbb{N}$ satisfy $0 < m$, $m \mid N$, $m < N$, and let p be prime with $p \equiv 1 \pmod{m}$. Let $s_p(N)$ be the ordinary base- p digit sum of N . Then*

$$v_p(G(N; m)) = \begin{cases} 1, & s_p(N) = m, \\ 0, & s_p(N) \neq m. \end{cases}$$

This is McTague's Theorem Q in the present divisibility regime. The Lean formalization includes it because the reduced moduli $m = 1$ and $m = 2$ arise after scaling the formulas for $m = 3, 4, 6$.

8 Scaling

The following scaling identity removes common trailing base- p zeros from the row and the modulus.

Theorem 8.1 (Scaling). *For prime p and natural numbers c, q, N' , one has*

$$v_p(G(p^c N'; p^c q)) = v_p(G(N'; q)).$$

The proof is a direct Kummer carry-shift argument. The admissible indices in the scaled problem are precisely p^c times the admissible indices in the unscaled problem, and multiplying both lower and upper indices by p^c appends c zero digits in base p , leaving the number of borrows unchanged.

9 Applications: the moduli 3, 4, 6

For the small moduli 3, 4, and 6, every prime not dividing the modulus is congruent to either 1 or -1 modulo the modulus. Primes dividing the modulus are reduced by Theorem 8.1. This yields complete prime-by-prime formulas.

Let $M_-(m, p, N)$ denote the right-hand side of Theorem 1.1 and $M_+(m, p, N)$ denote the right-hand side of Theorem 7.1.

Corollary 9.1 (Modulus 3). *If p is prime, $3 \mid N$, and $3 < N$, then*

$$v_p(G(N; 3)) = \begin{cases} M_+(1, 3, N/3), & p = 3, \\ M_+(3, p, N), & p \equiv 1 \pmod{3}, \\ M_-(3, p, N), & p \equiv -1 \pmod{3}. \end{cases}$$

Corollary 9.2 (Modulus 4). *If p is prime, $4 \mid N$, and $4 < N$, then*

$$v_p(G(N; 4)) = \begin{cases} M_+(1, 2, N/4), & p = 2, \\ M_+(4, p, N), & p \equiv 1 \pmod{4}, \\ M_-(4, p, N), & p \equiv -1 \pmod{4}. \end{cases}$$

Corollary 9.3 (Modulus 6). *If p is prime, $6 \mid N$, and $6 < N$, then*

$$v_p(G(N; 6)) = \begin{cases} M_-(3, 2, N/2), & p = 2, \\ M_+(2, 3, N/3), & p = 3, \\ M_+(6, p, N), & p \equiv 1 \pmod{6}, \\ M_-(6, p, N), & p \equiv -1 \pmod{6}. \end{cases}$$

These corollaries are applications of the theorem layer rather than separate headline novelty claims.

10 Formal verification and provenance

The theorem layer described above has been formalized in Lean 4 / Mathlib. The principal theorem is

`PascalMinusOne.minus_one_valuation.`

The public formal theorem layer also includes

```
PascalMinusOne.plus_one_valuation, PascalMinusOne.scaling_valuation,
PascalMinusOne.modulus_three_valuation, PascalMinusOne.modulus_four_valuation,
PascalMinusOne.modulus_six_valuation.
```

The formalization uses Mathlib infrastructure for binomial coefficients, finite gcds, base- p digits, p -adic valuations, and Kummer’s theorem for binomial coefficients. The project proof development contains zero explicit `sorry`, `admit`, or replacement-axiom placeholders in the Lean sources.

A Palomar provenance record exists for the principal minus-one theorem:

PALOMAR-2026-09-25-000017, version 1.

The record is a verification/provenance artifact for the stated formal theorem; it is not peer review, publication, or evidence of historical priority.

The source repository is

<https://github.com/jfairfaxball-348/pascal-minus-one>

The Palomar record is

<https://palomar-registry.org/entry.html?id=PALOMAR-2026-09-25-000017&version=1>

Acknowledgements

I thank the Mathlib community for the formal infrastructure used here, including Kummer’s theorem for binomial coefficients, digit lemmas, gcd machinery, and p -adic valuation results. I also acknowledge McTague’s and Wu’s closely related work on the same restricted binomial-gcd family, and the Lean formalization by Guo, Qiu, Cao, Feng and Gao of a different binomial-gcd problem.

References

- [1] B. Ram, Common factors of $n!/[m!(n-m)!]$, ($m = 1, 2, \dots, n-1$), *Journal of the Indian Mathematical Club (Madras)* **1** (1909), 39–43.
- [2] E. E. Kummer, *Über die Ergänzungssätze zu den allgemeinen Reciprocitätsgesetzen*, *Journal für die reine und angewandte Mathematik* **44** (1852), 93–146.
- [3] H. Joris, C. Oestreicher and J. Steinig, The greatest common divisor of certain sets of binomial coefficients, *Journal of Number Theory* **21** (1985), no. 1, 101–119. DOI: 10.1016/0022-314X(85)90013-7.
- [4] Andrew Granville, Arithmetic properties of binomial coefficients I: Binomial coefficients modulo prime powers, *CMS Conference Proceedings* **20** (1997), 253–276.
- [5] Gil Kaplan and Dan Levy, GCD of truncated rows in Pascal’s triangle, *Integers* **4** (2004), A14.
- [6] Carl McTague, The Cayley plane and string bordism, *Geometry & Topology* **18** (2014), no. 4, 2045–2078. DOI: 10.2140/gt.2014.18.2045.
- [7] Carl McTague, On the greatest common divisor of binomial coefficients $C(n, q), C(n, 2q), C(n, 3q), \dots$, *American Mathematical Monthly* **124** (2017), no. 4, 353–356. DOI: 10.4169/amer.math.monthly.124.4.353. Corrected preprint: arXiv:1510.06696v5.
- [8] Siao Hong, The greatest common divisor of certain binomial coefficients, *Comptes Rendus Mathématique* **354** (2016), no. 8, 756–761. DOI: 10.1016/j.crma.2016.06.001.
- [9] Jiaqi Xiao, Pingzhi Yuan and Xucan Lin, The greatest common divisor of certain set of binomial coefficients, *Mathematical Theory and Applications* **42** (2022), no. 1, 85–91.
- [10] Sunben Chiu, Pingzhi Yuan and Tao Zhou, On the greatest common divisor of binomial coefficients, *Bulletin of the Korean Mathematical Society* **60** (2023), no. 4, 863–872. DOI: 10.4134/BKMS.b220166.
- [11] Chan-Liang Chung, Tse-Chung Yang and Kanglun Zhou, The greatest common divisor of sets of binomial coefficients with restrictions, *Contemporary Mathematics* **6** (2025), no. 1, 971–985. DOI: 10.37256/cm.6120255017.
- [12] Chai Wah Wu, Computing the greatest common divisor of binomial coefficients $\binom{mn}{mk}$, arXiv:2606.20940v2, 2026.
- [13] Dakai Guo, Ruichen Qiu, Yichuan Cao, Ruyong Feng and Xiao-Shan Gao, A greatest common divisor criterion of certain binomial coefficients, arXiv:2606.22997, 2026.