

AUTOMATED PROOFS OF UNSTABLE ADAMS DIFFERENTIALS

JAKE FRANCIS BAER

ABSTRACT. We present a computer-based approach to computing differentials in the unstable Adams spectral sequence by systematically applying the unstable Leibniz rule and naturality with respect to maps in the EHP sequence. We tabulate our results in tables of upper and lower bounds on the orders of 2-primary unstable homotopy groups of spheres through the unstable 50-stem. We provide examples of proofs for several differentials and give a guide to interpreting the associated unstable Adams charts and flow chart diagrams for differential proofs.

CONTENTS

1. Introduction	1
2. The unstable Adams E_2 -page	3
3. The computation of unstable Adams differentials	5
4. Results	12
References	18

1. INTRODUCTION

The homotopy type of a cell complex is encoded in its attaching maps, which specify how each cell is glued to the lower-dimensional skeleton during inductive construction. Cellular approximation allows such maps to be decomposed into elements of homotopy groups of spheres, so the classification of cell complexes up to homotopy equivalence is governed by the structure of these groups. The explicit computation of unstable homotopy groups of spheres begins with Hopf’s construction of the first non-trivial elements via the Hopf invariant [Hop31] and the low-stem computations of Pontryagin [Pon38] and G. W. Whitehead [Whi50]; Serre’s spectral sequence methods made systematic calculation possible [Ser53]. Toda’s foundational work applied the EHP sequence and bracket arguments, yielding tables of $\pi_{n+k}(S^n)$ for $k \leq 19$ [Tod62]. Mimura and collaborators pushed these computations through roughly the 24-stem [MT63; Mim64; Mim65; MMO75]. Oda extended the 2-primary computations through the 30-stem with partial results in the 31- and 32-stems using refinements of Toda’s methods [Oda76; Oda77]. The remaining ambiguities were resolved by Inoue–Mukai and Miyauchi–Mukai [IM12; MM17]. Separately, Curtis and Mahowald determined the unstable homotopy groups of S^3 through the 52-stem in the only published application of the unstable Adams spectral sequence to computing unstable stems [CM89].

This paper consists of two distinct computations. The first is a modern reimplement of Curtis’ algorithm for computing the homology of the lambda algebra, which is the E_2 -page of the unstable Adams spectral sequence. We compute the unstable Adams E_2 -page for all spheres through total degree 76 as well as all values of the algebraic EHP maps (Proposition 2.1) and all unstable algebraic compositions on the Adams E_2 -page. The second computation uses the algebraic structure of the unstable Adams E_2 -page and input differentials from the stable Adams spectral sequence to propagate unstable Adams differentials using the unstable Leibniz rule and naturality of the EHP maps. This propagator tries all possible combinations of EHP maps and unstable compositions to constrain the possible values of an unstable Adams differential until only one remains. The precise combination of EHP maps and unstable compositions used to compute a differential is recorded in computer-generated flow charts. By resolving unstable Adams differentials and computing the E_∞ -page we prove the following main theorem.

Theorem 1.1. *Upper and lower bounds on the orders of the 2-primary homotopy groups $\pi_{n+k}(S^n)$ for $0 \leq k \leq 50$ are given in Tables 2 to 5.*

2020 *Mathematics Subject Classification.* Primary 55-08; Secondary 55Q40, 55T15.

Key words and phrases. Unstable homotopy groups of spheres, unstable Adams spectral sequence, lambda algebra, EHP sequence, computer-assisted proofs.

Email: francis@wayne.edu.

Through the unstable 32-stem these bounds are consistent with, and largely recover, the previously known orders due to the authors cited above (together with Curtis–Mahowald’s computations for S^3); the uniform machine-generated determination of unstable Adams differentials for all spheres, and the resulting bounds beyond the 32-stem, are new.

1.1. The lambda algebra. In [Section 2](#) we describe our computational approach to the structure of the unstable Adams E_2 -page. We reimplement Curtis’ lambda algebra algorithm from scratch. Our implementation contains nothing conceptually new and the theory underlying this algorithm is well exposed in the many sources we reference in [Section 2](#). By writing our implementation in a fast modern language and optimizing the key inner loops, we compute the homology of the lambda algebra and all algebraic compositions and algebraic EHP maps through total degree 76. While this algorithm for computing algebraic EHP maps and algebraic compositions has existed for several decades, neither of these has been computed exhaustively in a significant range, let alone applied to the computation of unstable Adams differentials. The most conceptually original part of [Section 2](#) consists of our generalization and exhaustive computation of the map of E_2 -pages used by Curtis and Mahowald to compute unstable Adams differentials for S^3 by comparison with the stable Adams spectral sequence for $\mathbb{S}/2$ [[CM89](#)]. While Curtis and Mahowald use this map to compute unstable Adams differentials, they never compute its values explicitly and only indirectly deduce its values by using the fact that this map is an isomorphism in sufficiently high filtration relative to the stem. This map is essential to our results since without it there are many differentials for low-dimensional spheres like S^3 and S^5 that we are unable to compute using EHP naturality and the Leibniz rule alone.

Remark. By running our Curtis algorithm on the Wayne State University high-performance grid, we computed the homology of the lambda algebra additively in degrees $s + f < 90$. The database of cycle representatives for this computation exceeds 50 GB in size, so we did not attempt to compute algebraic compositions between total degrees 76 and 90.

1.2. Unstable Adams differentials. In [Section 3](#) we describe our differential propagation algorithm, which uses the algebraic data from [Section 2](#). Our algorithm systematically applies naturality of unstable Adams differentials with respect to EHP maps and the unstable Leibniz rule, which are stated precisely in [Proposition 2.1](#) and [Proposition 2.2](#) below. Our approach to differential propagation is based on the approach in [[Bea22](#)] to propagating stable Adams d_2 -differentials using the Leibniz rule for the sphere spectrum. Our implementation starts with the space of all possible matrix representations of a d_r -differential considered as a map of $\mathbb{F}_2$ -vector spaces. In order to apply naturality and the Leibniz rule, we rephrase [Proposition 2.1](#) and [Proposition 2.2](#) as statements that certain diagrams have to commute. Our propagation algorithm then iteratively checks for matrices that fail to commute in the relevant diagrams, and removes these from our space of possible d_r -differential values. By repeating this process many times on all bidegrees for all spheres, we can narrow down the space of possible d_r -differentials until there is a unique matrix that respects all of the constraints imposed by [Proposition 2.1](#) and [Proposition 2.2](#). This process of iteratively applying constraints in order to deduce a unique d_r -differential value is recorded in the form of flow charts from which the reader may reconstruct the proof of any differential.

1.3. Results. In [Section 4](#) our results are presented by unstable Adams charts and tables of upper and lower bounds on the 2-primary homotopy groups of spheres through dimension 50. The algebraic data generated in [Section 2](#) and the differential propagation algorithm described in [Section 3](#) allow us to determine all differentials in the unstable Adams spectral sequence for most spheres of dimension at most 50 and in most stems up to 50. We have partial results in some cases. Our results are visualized in interactive Adams charts, rendered with the SeqSee visualization engine [[BI25](#)], displaying the spectral sequences for spheres S^n with $2 \leq n \leq 50$. These charts display proven differentials as well as the remaining possible unknown differentials on each page. The unstable Adams spectral sequence gives us an upper and lower bound on the order of each 2-primary unstable stem. These upper and lower bounds are tabulated in [Section 4.6](#). The homotopy groups whose orders are given in [Table 4](#) and [Table 5](#) occur entirely in the metastable range, i.e. the range $s \leq 3n - 4$. The most interesting entries are therefore located in the purely unstable region in the lower left corner of [Table 2](#). For some spheres such as S^5 , S^9 , S^{11} , S^{13} , S^{17} , S^{18} , and S^{19} , no uncertainties exist and we may read off the exact orders of their 2-primary homotopy groups through at least the 45-stem. For other spheres like S^{12} and S^{24} there are many remaining uncertainties and we can only bound the order of most of their homotopy groups instead of providing an exact value.

1.3.1. *Reading the tables.* Tables 2 to 5 in Section 4.6 display the upper and lower bounds on the orders of 2-primary unstable homotopy groups computed using the differential propagation algorithm in Section 3 and the unstable Adams E_2 -page data from Section 2. To suppress redundant stable information, we indicate the stable range by the vertical lines in the upper part of each table. The 2-primary metastable range is contained between the stable range vertical lines and the pink staircase boundary drawn through Table 2 and Table 3. The entry in column n and row s bounds the order of the 2-primary component of $\pi_{n+s}(S^n)$, recorded through its base-2 logarithm. A single entry e marks a homotopy group with no remaining unknown differentials, whose 2-primary component has order exactly 2^e . When a homotopy group still admits unknown differentials, we can only bracket its order. The entry is then a range e/ℓ , written upper bound first, meaning that the 2-primary component of $\pi_{n+s}(S^n)$ has order 2^m for some $\ell \leq m \leq e$.

1.4. **Relation to prior work.** Our results have been checked against the existing work of Toda, Mimura, Miyauchi, Mukai, Inoue, Mori, and Oda on 2-primary homotopy groups of spheres. Our results are compatible with theirs in the sense that the orders they compute fit between our upper and lower bounds through the unstable 32-stem. There are a few places between the 19-stem and the 32-stem where our computation admits an uncertain differential that could be resolved by referring to the work of these authors. In such places, we choose not to fill in this outside information. By doing this we ensure that our results are entirely machine generated in an attempt to minimize the risk of human error. In forthcoming joint work with Balderrama, Belmont, and Isaksen, we use the unstable Adams spectral sequence in conjunction with the EHP spectral sequence to compute all metastable homotopy groups of spheres whose stabilizations lie in stems 0 through 66 [BBBI].

2. THE UNSTABLE ADAMS E_2 -PAGE

We begin by reviewing the structure of the unstable Adams spectral sequence and describing the computational methods used to determine its E_2 -page. Recall that there is a cofree resolution of $\tilde{H}_*(S^n; \mathbb{F}_2)$ in unstable comodules over the dual Steenrod algebra $\mathcal{A}_*$ that is of finite type in each degree and is presented as $\mathcal{A}_* \otimes \Sigma^n \Lambda(n)$, following Singer [Sin75]. Here $\mathcal{A}_* \otimes N = \bigoplus_j J(j) \otimes N_j$ is the injective unstable comodule on a graded $\mathbb{F}_2$ -vector space N , where $J(j) \subseteq \mathcal{A}_*$ is the span of the monomials $\xi_1^{r_1} \xi_2^{r_2} \cdots$ with $\sum_i r_i \leq j$, and $\Lambda(n)$ is a subcomplex of the lambda algebra Λ , the differential bigraded algebra presented by

$$(2.1) \quad \Lambda = \mathbb{F}_2 \langle \lambda_0, \lambda_1, \lambda_2, \dots \rangle / \sim \quad \begin{aligned} \lambda_i \lambda_{2i+1+m} &\sim \sum_{j \geq 0} \binom{m-1-j}{j} \lambda_{i+m-j} \lambda_{2i+1+j} \\ d(\lambda_n) &= \sum_{j \geq 0} \binom{n-1-j}{j+1} \lambda_{n-1-j} \lambda_j \end{aligned}$$

with the convention that $\binom{a}{b} = 0$ unless $0 \leq b \leq a$. As an $\mathbb{F}_2$ -vector space, Λ has a basis of *admissible* monomials $\lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_f}$, those satisfying $2i_k \geq i_{k+1}$ for all k . We write $\Lambda(n) \subseteq \Lambda$ for the subcomplex spanned by the admissible monomials with $i_1 \leq n-1$, and $\Lambda^{s,f}(n)$ for its subspace spanned by such monomials of length f whose indices sum to s [Sin75]. We write $\alpha \smile \beta$ for the concatenation product of $\alpha, \beta \in \Lambda(n)$, and reserve $\circ$ for the induced composition in homology. We write $[\lambda_I]$ for the class in homology whose cycle representative has leading term λ_I , with respect to the lexicographic order on the index sequences of admissible monomials.

This algebraic resolution can be realized as a filtration in spaces that begets an unstable version of the Adams spectral sequence and has signature

$$E_2^{s,f} = H^{s,f} \Lambda(n) \Rightarrow (\pi_{s+n} S^n)_2^\wedge$$

by work of Massey–Peterson and Harper–Miller [MP67; HM89]. An alternative construction of this spectral sequence is given using the lower central series filtration on simplicial groups [BC70; Cur71; Lie72]; the identification of $\Lambda(n)$ as the E_1 -page of the p -lower central series spectral sequence appears already in [Bou+66, Section 5.4]. For a modern survey of these ideas see [BM24]. Recall that, 2-locally, James' fibration $S^n \rightarrow \Omega S^{n+1} \rightarrow \Omega S^{2n+1}$ gives rise to the long exact EHP sequence relating the homotopy groups of neighboring spheres, in which E is induced by suspension, H by the James–Hopf invariant, and P by Whitehead products; see [MT95] for a survey. When considered as a family, the unstable Adams spectral sequences for all S^n exhibit the following structure:

Proposition 2.1 ([PW19; Sin75; Cur71]). *There are chain maps*

$$\Lambda^{s,f}(n) \xrightarrow{E} \Lambda^{s,f}(n+1) \xrightarrow{H} \Lambda^{s-n,f-1}(2n+1) \xrightarrow{P} \Lambda^{s-1,f+1}(n),$$

which detect E, H, and P in $\pi_(-)$ and commute with unstable Adams differentials.*

Proposition 2.2 ([BK73b; BK73a; IMW18]). *There exists an unstable composition pairing*

$$\begin{aligned} H^{s,f} \Lambda(n) \otimes H^* \Lambda(n+s) &\xrightarrow{-\circ-} H^* \Lambda(n), \\ \alpha \otimes \beta &\mapsto \alpha \smile E^f \beta, \end{aligned}$$

which detects geometric compositions in $\pi_*(-)$ and respects an unstable Leibniz rule

$$d_r(\alpha \circ E\beta) = d_r(\alpha) \circ \beta + \alpha \circ d_r(E\beta).$$

2.1. The Curtis procedure. Filtering $\Lambda(n)$ by the subcomplexes $\Lambda(m)$ for $m \leq n$ yields a spectral sequence known as the *algebraic EHP spectral sequence* [Rav86; CGMM87], whose filtration quotients are identified with shifted copies of the $\Lambda(2m+1)$ by the maps H of Proposition 2.1. The *Curtis procedure* computes the homology of $\Lambda(n)$ by computing the differentials in this spectral sequence [Whi70; Cur71]: each admissible monomial either survives as the leading term of a cycle representative of a homology class or is cancelled by a differential. The Curtis procedure has had numerous well-documented implementations [Han74; Tan85; CGMM87; WX16]. Our implementation is written in Rust (version 1.96) [Bae26]. We have computed the homology of $\Lambda(n)$ additively in all total degrees $s+f < 90$; the dataset released with this paper [Bae26] covers all total degrees $s+f \leq 76$. The data generated by our implementation of the Curtis homology procedure consists of the following:

Data 2.3. The plain-text files `rank.csv` and `names.json` in the `data/` directory of [Bae26] contain

- (1) The dimension of $H^{s,f} \Lambda(n)$ for all $s+f \leq 76$
- (2) For each basis element, the lexicographically leading admissible monomial of a cycle representative

The full cycle representatives, together with the minimal set of differentials in the algebraic EHP spectral sequence used to reduce them, are regenerated on demand by our released implementation; at the full range they are too large to ship.

To operate on a class in $H^{s,f} \Lambda(n)$ we first lift its cycle representative to $\Lambda(n)$ and then operate on it, in our case either by some concatenation product or by the H map in Proposition 2.1. We then take the resulting polynomial in $\Lambda(m)$ and systematically add boundaries until we have expressed it as a linear combination of cycle representatives in the homology of $\Lambda(m)$. This approach has been used by Tangora to compute some products and Massey products in the classical Adams E_2 -page [Tan93; Tan94]. We use this method to compute the following:

Data 2.4. The plain-text table files `E.csv`, `H.csv`, `P.csv`, and `compositions.csv` in the `data/` directory of [Bae26] contain a full description through total degree 76 of the values of the maps in Proposition 2.1 and Proposition 2.2 on every $\mathbb{F}_2$ -linear basis element from Data 2.3.

2.2. The fiber of the double suspension. Write $e_m \Lambda$ for a copy of Λ shifted by a generator e_m in degree m , so that $\mathbb{F}_2\{e_m\} = \Sigma^m \mathbb{F}_2$. The E_2 -page of the Adams spectral sequence for $\Sigma^{2n-1} \mathbb{S}/2$ can be described by taking the homology of $e_{2n} \Lambda \oplus e_{2n-1} \Lambda$ with differential inherited from Λ along with $d(e_{2n}) = e_{2n-1} \lambda_0$ [Pri70]. The following result is implicit in the cited papers, but for completeness we provide a proof sketch along with all relevant references.

Theorem 2.5 ([Mah75; Mah82; MT94]). *For each n there exists the following:*

- (1) a chain map

$$\begin{aligned} \Lambda(2n+1) &\rightarrow e_{2n} \Lambda \oplus e_{2n-1} \Lambda & \lambda_{2n-1} \lambda_I &\mapsto e_{2n-1} \lambda_I \\ & & \lambda_{2n} \lambda_I &\mapsto e_{2n} \lambda_I + \epsilon e_{2n-1} \lambda_{4n+1} \lambda_{I'}, \end{aligned}$$

where $\epsilon = 1$ if $\lambda_I = \lambda_{4n} \lambda_{I'}$ and $\epsilon = 0$ otherwise, and where monomials beginning with λ_i for $i < 2n-1$ are sent to 0;

- (2) a map $\Omega^3 S^{2n+1} \rightarrow \Omega^\infty \Sigma^\infty \Sigma^{4n-4} \mathbb{R}P^2$ which can be covered by a map of unstable resolutions described on the E_1 -page by the chain map in (1).

Proof. Step 1. Write $\text{fib}_n(E^2)$ for the fiber of the double suspension $S^{2n-1} \xrightarrow{E^2} \Omega^2 S^{2n+1}$, and let C_n denote the mapping cone of the chain-level double suspension $E^2: \Lambda(2n-1) \rightarrow \Lambda(2n+1)$ from Proposition 2.1, so that $C_n = \Lambda^{*,*+1}(2n-1) \oplus \Lambda^{*,*}(2n+1)$ as bigraded modules. Following [MT87], we build a resolution for $\text{fib}_n(E^2)$ whose associated algebraic resolution is $\mathcal{A}_* \otimes \Sigma^{2n-2} C_n$. This resolution is constructed so that the map $\Omega^3 S^{2n+1} \rightarrow \text{fib}_n(E^2)$ is covered by a map of resolutions that induces the inclusion of the second summand

$$(2.2) \quad \mathcal{A}_* \otimes \Sigma^{2n-2} \Lambda(2n+1) \hookrightarrow \mathcal{A}_* \otimes \Sigma^{2n-2} C_n.$$

Note that C_n is chain homotopy equivalent to the quotient complex $\Lambda(2n+1)/\Lambda(2n-1)$, which Mahowald [Mah75, Section 2] presents as

$$\begin{aligned} \operatorname{cof}_n(d_1) &:= \kappa_1 \Lambda(4n+1) \oplus \kappa_0 \Lambda(4n-1), & d(\kappa_1) &= \kappa_0 \lambda_0, \\ |\kappa_i| &= (2n-1+i, 1), & d(\kappa_1 \lambda_{4n}) &= \kappa_0 (\lambda_0 \lambda_{4n} + d(\lambda_{4n+1})), \end{aligned}$$

with differential otherwise inherited from $\Lambda(4n+1)$ and $\Lambda(4n-1)$; here $d(\lambda_{4n+1})$ is given by (2.1). Our notation is meant to reflect that this construction is equal to the mapping cone for the algebraic EHP d_1 -differential $\Lambda(4n+1) \xrightarrow{H \circ P} \Lambda(4n-1)$. Composing the inclusion (2.2) with this quotient map, we conclude that the map of resolutions covering $\Omega^3 S^{2n+1} \rightarrow \operatorname{fib}_n(E^2)$ induces the chain map

$$(2.3) \quad \Lambda(2n+1) \rightarrow \operatorname{cof}_n(d_1), \quad \lambda_{2n} \alpha \mapsto \kappa_1 \alpha, \quad \lambda_{2n-1} \alpha \mapsto \kappa_0 \alpha.$$

Step 2. Cohen showed that there is a map $\operatorname{fib}_n(E^2) \rightarrow \Omega^4 \operatorname{fib}_{n+1}(E^2)$ that has degree 1 on the bottom cell and such that the colimit of the sequence of maps

$$\operatorname{fib}_n(E^2) \rightarrow \Omega^4 \operatorname{fib}_{n+1}(E^2) \rightarrow \Omega^8 \operatorname{fib}_{n+2}(E^2) \rightarrow \cdots$$

is homotopy equivalent to $\Omega^\infty \Sigma^\infty \Sigma^{4n-4} \mathbb{R}P^2$ [Coh83, Propositions 1.3 and 1.4]. In algebra we have the following commutative diagram of exact sequences

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & \Lambda(2n-1) & \longrightarrow & \Lambda(2n) & \longrightarrow & \Lambda(4n-1) \rightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \rightarrow & \Lambda(2n-1) & \rightarrow & \Lambda(2n+1) & \longrightarrow & \operatorname{cof}_n(d_1) \rightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & \Lambda(4n+1) = \Lambda(4n+1) & & \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

which shows that the range where $\mathcal{A}_* \otimes \operatorname{cof}_n(d_1)$ is acyclic is equal to the acyclicity range of $\mathcal{A}_* \otimes \Lambda(4n-1)$. The main theorem of Harper–Miller [HM89, Theorem 3.9] states that for a simply connected space of finite type with *very nice* cohomology (one whose $\mathbb{F}_2$ -cohomology is a free unstable algebra $U(M)$ on an unstable module M over the Steenrod algebra), this acyclicity range is equal to the range in which certain structure maps in the associated unstable Adams resolution are surjective on cohomology. This result was generalized to the above resolution for $\operatorname{fib}_n(E^2)$ in [Tho90, Proposition 2.26] (stated there for odd primes; the proof carries over to $p = 2$). Proposition 6.3 and the lambda-algebra estimates from Proposition 6.6 in [MT94] show that any chain map $\mathcal{A}_* \otimes \operatorname{cof}_n(d_1) \rightarrow \mathcal{A}_* \otimes \operatorname{cof}_{n+1}(d_1)$ is uniquely determined up to chain homotopy by its effect in homological degree 0. In particular, since the map $\operatorname{fib}_n(E^2) \rightarrow \Omega^4 \operatorname{fib}_{n+1}(E^2)$ is an isomorphism on $H^{4n-3}(-; \mathbb{Z})$, the induced map on resolutions is chain homotopic to the chain map $\mathcal{A}_* \otimes \operatorname{cof}_n(d_1) \rightarrow \mathcal{A}_* \otimes \operatorname{cof}_{n+1}(d_1)$ induced by the maps

$$(2.4) \quad \operatorname{cof}_n(d_1) \rightarrow \operatorname{cof}_{n+1}(d_1) \rightarrow \operatorname{cof}_{n+2}(d_1) \rightarrow \cdots$$

from [Mah75, Definition 3.2]. Composing the map in (2.3) with the maps in (2.4) gives the result stated in the theorem. $\square$

Data 2.6. The plain-text table file `C2.csv` in the `data/` directory of [Bae26] describes all values of the maps from Theorem 2.5 on E_2 -pages through total degree 76.

In practice, Theorem 2.5 and Data 2.6 give us a map of spectral sequences

$$(2.5) \quad E_r^{s,f}(S^{2n+1}) \xrightarrow{s_{2n+1}} E_r^{s-2n+1,f-1}(\mathbb{S}/2)$$

which we use to prove unstable Adams differentials using the stable Adams spectral sequence for $\mathbb{S}/2$. For $n = 1$ this map was the central tool in [CM89].

3. THE COMPUTATION OF UNSTABLE ADAMS DIFFERENTIALS

In this section we describe our algorithm for exhaustively applying Proposition 2.1 and Proposition 2.2 to the propagation of unstable Adams differentials. Given the degree shifts in Proposition 2.1 and Proposition 2.2, it helps to consider the unstable Adams spectral sequence for all spheres as a single

trigraded spectral sequence where we denote the $\mathbb{F}_2$ -vector space $E_r^{s,f}(S^n)$ by $\mathcal{U}_r^{n,s,f}$. Our strategy begins with the observation that stabilization and [Theorem 2.5](#) give us maps of spectral sequences

$$(3.1) \quad \mathcal{U}_r^{n,s,f} \xrightarrow{E_n^\infty} E_r^{s,f}(\mathbb{S}) \quad \mathcal{U}_r^{2n+1,s,f} \xrightarrow{s_{2n+1}} E_r^{s-2n+1,f-1}(\mathbb{S}/2),$$

where the targets are stable Adams spectral sequences and s_{2n+1} is the map (2.5). The Adams differentials for the sphere spectrum and the cofiber of 2 have been well studied and are completely known at least through the 80-stem [[IWX23](#); [LWX24](#); [LWX25](#)]. The fact that these maps commute with differentials provides the initial unstable Adams differentials, which we then propagate using [Proposition 2.1](#) and [Proposition 2.2](#).

3.1. Affine linear algebra. Our method for propagating unstable Adams differentials—using naturality with respect to EHP maps and the unstable Leibniz rule—is based on the strategy for stable Adams d_2 -differentials presented in [[Bea22](#)]. Throughout this section we write $\mathbf{d}_r$ for the matrix representing the differential d_r on the tridegree in question. Recall that we can express the space of possible values of $\mathbf{d}_r$ as an affine subspace $\delta_r + \mathcal{I}_r$, where the *offset* $\delta_r \in \text{Hom}(\mathcal{U}_r^{n,s,f}, \mathcal{U}_r^{n,s-1,f+r})$ is a chosen matrix representative and the *indeterminacy* $\mathcal{I}_r \subseteq \text{Hom}(\mathcal{U}_r^{n,s,f}, \mathcal{U}_r^{n,s-1,f+r})$ is a linear subspace.

Remark. Typically the Leibniz rule is used to take information about a differential $d_r(x)$ and a product $x \circ y$ and deduce something about the differential $d_r(x \circ y)$. Our implementation instead considers the set of all possible matrix representatives for d_r on some tridegree and narrows down our set of possible differentials by excluding matrices for which certain squares fail to commute. The following description in terms of affine linear algebra is meant to make this idea precise.

We use the affine linear algebra library in SageMath (version 10.4) [[The24](#)] to translate commutative diagrams involving $\mathbf{d}_r$ into linear constraints on its space of possible values by computing intersections of affine spaces and taking images and preimages under fixed linear maps. For each tridegree $\mathcal{U}_r^{n,s,f}$ we initialize its affine subspace of possible differentials $\mathcal{D}_r^{n,s,f}$ as $\mathbf{0} + \text{Hom}_{\mathbb{F}_2}(\mathcal{U}_r^{n,s,f}, \mathcal{U}_r^{n,s-1,f+r})$. For a fixed linear map F and affine subspace $\mathcal{D}$, define the pushforward and pullback operators

$$F_*(\mathcal{D}) = \{F \circ \mathbf{m} : \mathbf{m} \in \mathcal{D}\}, \quad F^*(\mathcal{D}) = \{\mathbf{m} \circ F : \mathbf{m} \in \mathcal{D}\}.$$

Proposition 3.1. *Let F be a linear map on $\mathcal{U}_r^{*,*,*}$ and denote its target tridegree on $\mathcal{U}_r^{n,s,f}$ by $F(n,s,f)$. Naturality of F with respect to $\mathbf{d}_r$ is the equation $F \circ \mathbf{d}_r = \mathbf{d}_r \circ F$. On affine subspaces, naturality with indeterminacy becomes:*

$$\mathbf{d}_r \in \mathcal{D}_r^{n,s,f} \Rightarrow \mathbf{d}_r \in (F^*)^{-1}(F_*(\mathcal{D}_r^{n,s,f})), \quad \mathbf{d}_r \in \mathcal{D}_r^{F(n,s,f)} \Rightarrow \mathbf{d}_r \in F_*^{-1}(F^*(\mathcal{D}_r^{n,s,f})).$$

In each implication the hypothesis concerns the differential on the tridegree appearing in its superscript, and the conclusion constrains the differential on the other tridegree.

Proof. The proof is the same as in [[Bea22](#)]. □

Here and throughout, the arrow $\leftarrow$ denotes algorithmic refinement: the space of possible differentials named on the left is replaced, in place, by its intersection with the space on the right. We therefore refine $\mathcal{D}_r^{*,*,*}$ by intersecting with the subspace of maps respecting naturality:

$$(3.2) \quad \mathcal{D}_r^{n,s,f} \leftarrow F_*^{-1}(F^*(\mathcal{D}_r^{F(n,s,f)})) \quad \mathcal{D}_r^{F(n,s,f)} \leftarrow (F^*)^{-1}(F_*(\mathcal{D}_r^{n,s,f})).$$

The goal is to intersect $\mathcal{D}_r^{n,s,f}$ with constraints until there is no indeterminacy:

$$\mathcal{D}_r^{n,s,f} = \mathbf{d}_r + \{0\}.$$

3.2. Naturality of EHP. In this subsection, we sketch our strategy for exhaustively applying the naturality result in [Proposition 2.1](#). This result implies that the following diagrams commute:

$$\begin{array}{ccc} \mathcal{U}_r^{n,s,f} & \xrightarrow{E} & \mathcal{U}_r^{n+1,s,f} & \mathcal{U}_r^{n,s,f} & \xrightarrow{H} & \mathcal{U}_r^{2n-1,s-n+1,f-1} \\ \mathbf{d}_r \downarrow & & \downarrow \mathbf{d}_r & \mathbf{d}_r \downarrow & & \downarrow \mathbf{d}_r \\ \mathcal{U}_r^{n,s-1,f+r} & \xrightarrow{E} & \mathcal{U}_r^{n+1,s-1,f+r} & \mathcal{U}_r^{n,s-1,f+r} & \xrightarrow{H} & \mathcal{U}_r^{2n-1,s-n,f+r-1} \\ & & \mathcal{U}_r^{2n+1,s,f} & \xrightarrow{P} & \mathcal{U}_r^{n,s+n-1,f+2} & \\ & & \mathbf{d}_r \downarrow & & \downarrow \mathbf{d}_r & \\ & & \mathcal{U}_r^{2n+1,s-1,f+r} & \xrightarrow{P} & \mathcal{U}_r^{n,s+n-2,f+r+2} & \end{array}$$

Terminal Nodes




FIGURE 1.

Internal Nodes




FIGURE 2.

Maps



FIGURE 3.

Products




FIGURE 4.

We can constrain the possible values of a differential on some unstable tridegree by iteratively applying (3.2) where F is the matrix form of the maps E , H , and P computed in Data 2.4. With no additional outside information, this procedure can only compute differentials that are zero. To compute nontrivial unstable Adams differentials, we assume knowledge of all differentials in the Adams spectral sequences for $\mathbb{S}$ and $\mathbb{S}/2$ through total degree 80 [IWX23; LWX24; LWX25] and lift their differentials to the unstable Adams spectral sequence via the maps in (3.1), the second of which is computed in Data 2.6. We present these deductions as flow charts, whose legend is given in Figures 1 to 4. The flow chart for any computed differential can be regenerated from the released data with the `why.py` script in [Bae26].

Algorithm 1 How a differential's proof is recorded as a flow chart. PROPAGATE shrinks the space $\mathcal{D}_r^{n,s,f}$ of still-possible values of $\mathbf{d}_r$ at every tridegree by intersecting it with the naturality and Leibniz constraints, *recording* each refinement together with the neighboring tridegrees it used. WHY then traces these records backward from the target differential to the assumed inputs.

```

1: procedure PROPAGATE( $r$ )
2:    $\mathcal{D}_r^{n,s,f} \leftarrow \text{Hom}(\mathcal{U}_r^{n,s,f}, \mathcal{U}_r^{n,s-1,f+r})$  for all  $(n, s, f)$  ▷ every value still possible
3:   restrict  $\mathcal{D}_r^{n,s,f}$  to the known differential at each assumed input ▷ stable  $\mathbb{S}$  and  $\mathbb{S}/2$ 
4:   repeat
5:     for all maps  $F \in \{E, H, P\}$  and tridegrees  $(n, s, f)$  do ▷ naturality, Proposition 2.1
6:        $\mathcal{D}_r^{n,s,f} \leftarrow \{ \text{maps making the } F\text{-naturality square commute} \}$ 
7:       if this shrank  $\mathcal{D}_r^{n,s,f}$ , record the step  $(F; F(n, s, f))$  at  $(n, s, f)$ 
8:     end for
9:     for all products  $x \cdot y$  landing in  $(n, s+s', f+f')$  do ▷ Leibniz, Proposition 2.2
10:      refine  $\mathcal{D}_r^{n,s+s',f+f'}$  to force  $\mathbf{d}_r(xy) = \mathbf{d}_r(x)y + x\mathbf{d}_r(y)$ 
11:      if this shrank it, record the step  $(x \cdot y; x, y)$  there
12:    end for
13:  until no  $\mathcal{D}_r^{n,s,f}$  changes
14: end procedure

15: function WHY( $w$ ) ▷  $w = (n, s, f)$ ; call on the target to be proved
16:    $\sigma \leftarrow$  the most recent step recorded at  $w$  ▷ empty if  $w$  was never refined
17:   create node  $N_w$ , colored by the state of  $\mathcal{D}_r^w$  at  $\sigma$  ▷ Figure 2
18:   if  $\sigma$  is empty or  $w$  is an assumed input then
19:     return  $N_w$  ▷ terminal: a stable/ $\mathbb{S}/2$  input, or an already-forced group
20:   else if  $\sigma$  is a naturality step for a map  $F$  then
21:     attach WHY( $F(w)$ ) to  $N_w$  on an edge labeled  $F$ 
22:   else  $\sigma$  is the Leibniz step for a product  $x \cdot y$ 
23:     attach WHY( $x$ ) and WHY( $y$ ) to  $N_w$ 
24:   end if
25:   return  $N_w$ 
26: end function

```

Reading a flow chart. The root is the differential being proved and the terminal nodes are the inputs we assume: a stable sphere differential $\mathbb{S}(s, f)$, a differential on the $\mathbb{S}/2$ column $C2(s, f)$, or a group on which $\mathbf{d}_r$ is already forced. Every edge is a single instance of naturality (Proposition 2.1) or the Leibniz rule (Proposition 2.2), so following the labeled maps E, H, P from the terminal nodes up to the root replays exactly the chain of constraints that pins down the value at the root. A node's color records how determined its differential was when it was used—gray unconstrained, blue partially constrained, green once a unique nonzero value is forced, red once it is forced to vanish—and each refined node also points back to its own earlier, less-determined state (the adjacent gray or blue node), so the chart shows $\mathcal{D}_r$ shrinking in place. Because every step is valid and the terminal nodes are assumptions, the chart is a complete, edge-by-edge checkable proof of the differential at its root.

The terminal nodes (Figure 1) mark the places where a proof begins. An orange node is the differential in some bidegree of a stable Adams spectral sequence, for either the sphere spectrum $\mathbb{S}$ or the mod-2 Moore spectrum $\mathbb{S}/2$; these are the inputs we assume. A red terminal node is a differential that vanishes because its target is the trivial vector space, and a gray terminal node is a completely unknown differential—one whose space of possible values is still the full space of linear maps from the source to the target. The internal nodes (Figure 2) record differentials that the propagation has already constrained. A green node is a differential that is fully known and nonzero, and a red node is one that is fully known and zero;

here *fully known* means that the space of possible matrices has been narrowed to a single element. A blue node is a differential that is only partially known, in the sense that some matrices have been ruled out but more than one possibility remains. The diagrams in Figures 3 and 4 record the two kinds of constraint. In the left-hand diagram of Figure 3, a map $F: \mathcal{U}_r^{n,s,f} \rightarrow \mathcal{U}_r^{F(n,s,f)}$ lets us use what we know about the differential on $\mathcal{U}_r^{F(n,s,f)}$ to refine the differential on $\mathcal{U}_r^{n,s,f}$; the right-hand diagram is the mirror image, refining $\mathcal{U}_r^{F(n,s,f)}$ using $\mathcal{U}_r^{n,s,f}$. Figure 4 shows the analogous refinements coming from the Leibniz rule for a product $x \cdot y$.

Example 3.2. Among unstable Adams differentials that stabilize to zero, the one of lowest stem occurs on $\mathcal{U}_2^{5,20,5}$. The commutativity of the following diagram shows how the differential on $\mathcal{U}_2^{5,20,5}$ can be obtained by pulling back the stable Adams differential on h_0f_0 through iterative applications of the maps H and P. Note that all unstable tridegrees in the diagram have a single generator and each horizontal map is nontrivial on that generator.

$$\begin{array}{ccccccc}
 \mathcal{U}_2^{5,20,5} & \xrightarrow{P} & \mathcal{U}_2^{2,21,7} & \xrightarrow{H} & \mathcal{U}_2^{3,20,6} & \xrightarrow{H} & \mathcal{U}_2^{5,18,5} \xrightarrow{E_5^\infty} E_2^{18,5}(\mathbb{S}) \cong \mathbb{F}_2\{h_0f_0\} \\
 \mathbf{d}_2 \downarrow & & \downarrow \mathbf{d}_2 & & \downarrow \mathbf{d}_2 & & \downarrow \mathbf{d}_2 = [1] \\
 \mathcal{U}_2^{5,19,7} & \xrightarrow{P} & \mathcal{U}_2^{2,20,9} & \xrightarrow{H} & \mathcal{U}_2^{3,19,8} & \xrightarrow{H} & \mathcal{U}_2^{5,17,7} \xrightarrow{E_5^\infty} E_2^{17,7}(\mathbb{S}) \cong \mathbb{F}_2\{h_0^3e_0\}
 \end{array}$$

The stable group $E_2^{18,5}(\mathbb{S})$ is generated by h_0f_0 , which supports the known stable differential $d_2(h_0f_0) = h_0^3e_0$. Since the stabilization map E_5^∞ carries the generator of $\mathcal{U}_2^{5,18,5}$ to h_0f_0 and the generator of $\mathcal{U}_2^{5,17,7}$ to $h_0^3e_0$, commutativity of the rightmost square forces the differential on $\mathcal{U}_2^{5,18,5}$ to be nonzero. Repeating this argument square by square from right to left (each of the maps H, H, and P commutes with $\mathbf{d}_2$ and is nonzero on the relevant generator) forces the differentials on $\mathcal{U}_2^{3,20,6}$, on $\mathcal{U}_2^{2,21,7}$, and finally on $\mathcal{U}_2^{5,20,5}$ to be nonzero as well. Using the notational schema presented above, we can abbreviate this commutative diagram as a flow chart given in Figure 5.





FIGURE 5. Flow chart for $\mathcal{D}_2^{5,20,5}$.

The proof in Example 3.2 is a relatively simple example of a multistep proof using EHP naturality. For more complicated arguments, our abbreviated notation allows us to efficiently communicate the inputs, logical steps, and output of a proof in a way that remains readable even when the corresponding commutative diagram becomes too large and convoluted to follow.

Example 3.3. There is a nontrivial differential on $\mathcal{U}_2^{4,20,6}$ that can be computed using stable differentials and applications of the maps E and P. The flow chart for this computation is given in Figure 6. The differential on $\mathcal{U}_2^{9,17,4}$ is completely determined by stable information using the maps E_9^∞ and s_9 . Similarly, there is a nontrivial differential on $\mathcal{U}_2^{3,20,6}$ that is lifted from the nontrivial differential on $E_2^{19,5}(\mathbb{S}/2)$. These nontrivial differentials can be pushed to $\mathcal{U}_2^{4,20,6}$ through the maps P and E, and the remaining possibilities for the differential on $\mathcal{U}_2^{4,20,6}$ can be ruled out using naturality of the stabilization map.

FIGURE 6. Flow chart for $\mathcal{D}_2^{4,20,6}$.

3.3. The unstable Leibniz rule. The unstable Leibniz rule from [Proposition 2.2](#) states that the following diagram commutes, where M is the algebraic composition map from [Proposition 2.2](#)

$$\begin{array}{ccc}
 \mathcal{U}_r^{n,s,f} \otimes \mathcal{U}_r^{n+s-1,s',f'} & \xrightarrow{M \circ (\text{id} \otimes E)} & \mathcal{U}_r^{n,s+s',f+f'} \\
 \left[\begin{array}{c} \mathbf{d}_r \otimes \text{id} \\ \text{id} \otimes (\mathbf{d}_r \circ E) \end{array} \right] \downarrow & & \downarrow \mathbf{d}_r \\
 (\mathcal{U}_r^{n,s-1,f+r} \otimes \mathcal{U}_r^{n+s-1,s',f'}) \oplus (\mathcal{U}_r^{n,s,f} \otimes \mathcal{U}_r^{n+s,s'-1,f'+r}) & \xrightarrow{[M \quad M]} & \mathcal{U}_r^{n,s+s'-1,f+f'+r}
 \end{array}$$

The commutativity of the full Leibniz diagram constrains the possible values of $\mathbf{d}_r$ on $\mathcal{U}_r^{n,s+s',f+f'}$ by updating $\mathcal{D}_r^{n,s+s',f+f'}$ according to

$$\mathcal{D}_r^{n,s+s',f+f'} \leftarrow ((M \circ (\text{id} \otimes E))^*)^{-1} (M_*(\mathcal{D}_r^{n,s,f} \otimes \text{id}) + M_*(\text{id} \otimes E^*(\mathcal{D}_r^{n+s,s',f'}))).$$

These constraints allow us to take information about Adams differentials in earlier stems and propagate it into later stems. We can similarly use the Leibniz rule to take information about Adams differentials in later stems and back-propagate it into earlier stems. If x is a cycle on the unstable Adams E_r -page, the unstable Leibniz rule reduces to the first of the following equations; if Ey is a cycle, it reduces to the second:

$$x \circ \mathbf{d}_r(Ey) = \mathbf{d}_r(x \circ Ey) \qquad \mathbf{d}_r(x) \circ y = \mathbf{d}_r(x \circ Ey).$$

We refer to these two specializations of the unstable Leibniz rule as the *partial Leibniz rule*.

Because algebraic compositions in the unstable Adams spectral sequence are not commutative, we need to keep track of whether we are multiplying on the left or on the right. Given $x \in \mathcal{U}_r^{n,s,f}$ and $y \in \mathcal{U}_r^{n+s,s',f'}$, we define the left and right multiplication maps

$$L_x : \mathcal{U}_r^{n+s,s',f'} \rightarrow \mathcal{U}_r^{n,s+s',f+f'} \qquad R_y : \mathcal{U}_r^{n,s,f} \rightarrow \mathcal{U}_r^{n,s+s',f+f'}.$$

The partial Leibniz rule then implies that the following diagrams commute for $x \in \mathcal{U}_r^{n,s,f}$ and $y \in \mathcal{U}_r^{n+s-1,s',f'}$. Note that y now lives one sphere lower than in the composition pairing, because the partial Leibniz rule gives information about the differential on $Ey \in \mathcal{U}_r^{n+s,s',f'}$ rather than on y itself. The left diagram commutes when x is a cycle, and the right diagram commutes when Ey is a cycle.

$$\begin{array}{ccc}
 \mathcal{U}_r^{n+s-1,s',f'} & \xrightarrow{L_x \circ E} & \mathcal{U}_r^{n,s+s',f+f'} \\
 \mathbf{d}_r \circ E \downarrow & & \downarrow \mathbf{d}_r \\
 \mathcal{U}_r^{n+s,s'-1,f'+r} & \xrightarrow{L_x} & \mathcal{U}_r^{n,s+s'-1,f+f'+r}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathcal{U}_r^{n,s,f} & \xrightarrow{R_{Ey}} & \mathcal{U}_r^{n,s+s',f+f'} \\
 \mathbf{d}_r \downarrow & & \downarrow \mathbf{d}_r \\
 \mathcal{U}_r^{n,s-1,f+r} & \xrightarrow{R_y} & \mathcal{U}_r^{n,s+s'-1,f+f'+r}
 \end{array}$$

On the level of affine subspaces we can express the partial Leibniz rule as a constraint on the space of possible differentials in the following way:

$$\mathcal{D}_r^{n,s,s',f'} \leftarrow ((L_x)_* \circ E^*)^{-1} ((L_x \circ E)^*(\mathcal{D}_r^{n,s+s',f+f'})) \qquad \mathcal{D}_r^{n,s,f} \leftarrow (R_y)_*^{-1} (R_{Ey}^*(\mathcal{D}_r^{n,s+s',f+f'}))$$

These updates back-propagate information from the product tridegree into its factors; forward propagation into the product tridegree is already handled by the full Leibniz update above. The algebraic compositions computed in [Data 2.4](#) provide a full description of the maps M , E , L_x , R_y , and R_{Ey} through total degree 76.

Remark. It is possible to write down commutative diagrams based on the full Leibniz rule that isolate the differentials on x or Ey without assuming that either factor tridegree consists of cycles; the analogous diagrams for the stable Leibniz rule are found in [\[Bea22\]](#). Our implementation applies these constraints

when neither partial rule is available, but the linear algebra involved in writing them down explicitly is more complicated, and we omit the formulas for brevity.

Example 3.4. There is a nontrivial differential on $\mathcal{U}_3^{9,45,11}$ that can be computed using the algebraic unstable compositions

$$[\lambda_8 \lambda_4 \lambda_1^3] \circ [\lambda_{12} \lambda_4 \lambda_5 \lambda_3^3] = [\lambda_8 \lambda_2^2 \lambda_4 \lambda_3^3 \lambda_6^2 \lambda_5 \lambda_3] \quad [\lambda_8 \lambda_4 \lambda_1^3] \circ [\lambda_2^4 \lambda_3 \lambda_5 \lambda_7 \lambda_3^2] = [\lambda_4 \lambda_2^7 \lambda_3^2 \lambda_6^2 \lambda_5 \lambda_3]$$

together with the stable input differential $d_3(\Delta h_2^2) = h_1 d_0^2$, where $\Delta h_2^2 = [\lambda_{12} \lambda_4 \lambda_5 \lambda_3^3]$ and $h_1 d_0^2 = [\lambda_7 \lambda_6 \lambda_2 \lambda_3 \lambda_4^2 \lambda_1^3]$.




FIGURE 7. Flow chart for $\mathcal{D}_3^{9,45,11}$.

By combining deductions using EHP naturality with deductions using the unstable Leibniz rule, we can construct increasingly complex arguments for otherwise challenging differentials.

Example 3.5. The following flow chart illustrates our process of applying constraints to $\mathcal{D}_3^{9,37,7}$. This computation uses constraints coming from both EHP naturality and the unstable Leibniz rule. One of the classes in $\mathcal{U}_3^{9,37,7}$ is a cycle because it is the image under H of a cycle in $\mathcal{U}_3^{5,41,8}$. The other generator of $\mathcal{U}_3^{9,37,7}$ supports a nontrivial d_3 -differential. The input data for this computation consists of nontrivial d_3 -differentials on $E_3^{44,10}(\mathbb{S})$ and $E_3^{52,12}(\mathbb{S}/2)$. The maps E_{17}^∞ and s_3 pull these differentials back to $\mathcal{U}_3^{17,44,10}$ and $\mathcal{U}_3^{3,53,13}$ respectively. By pushing and pulling these differentials through the EHP maps, we can deduce the nontrivial differential on $\mathcal{U}_3^{9,51,11}$. The product

$$[\lambda_7 \lambda_{12} \lambda_4 \lambda_5 \lambda_3^3] \circ [\lambda_6 \lambda_2 \lambda_3^2] = [\lambda_7 \lambda_2 \lambda_4 \lambda_5^3 \lambda_3 \lambda_6^2 \lambda_5 \lambda_3]$$

together with the fact that $[\lambda_6 \lambda_2 \lambda_3^2] \in \mathcal{U}_3^{46,14,4}$ is a cycle gives the differential on $[\lambda_7 \lambda_{12} \lambda_4 \lambda_5 \lambda_3^3] \in \mathcal{U}_3^{9,37,7}$.




FIGURE 8. Flow chart for $\mathcal{D}_3^{9,37,7}$ combining EHP and Leibniz rule.

4. RESULTS

This section presents our results: unstable Adams charts for the spheres S^n with $2 \leq n \leq 50$, displaying proven differentials as well as the remaining possible unknown differentials on each page, and the resulting upper and lower bounds on the orders of the 2-primary unstable homotopy groups, tabulated in [Section 4.6](#). A guide to reading our Adams charts is given in [Section 4.5](#), and the legend for the flow charts we use to present differential proofs is given in [Figures 1 to 4](#).

4.1. Uncertainties. If the d_r -differential on some tridegree is not fully determined, then we can not properly evaluate relations that land in that tridegree on the E_{r+1} -page. We are then unable to make general deductions about the d_{r+1} -differential on that tridegree. As we continue turning the page, ignoring these tridegrees creates a growing family of uncertain differentials around that original uncertain tridegree. In order to avoid this, we implement a much simpler element-wise solver which imposes the following logic:

- (1) If x is a cycle then $F(x)$ is a cycle
- (2) If $F(x)$ is not a boundary then x is not a boundary

This solver is applied at the end of the computation on each page and to tridegrees which were skipped by the main computation due to an uncertain differential value on an earlier page.

4.2. Differentials by contradiction. A handful of differentials can be resolved via proof by contradiction. In future work we will systematically apply this approach throughout all tridegrees. In this project we only compute 17 differentials this way. These particular differentials were chosen because leaving them unresolved makes our Adams charts much harder to read. The proofs are all similar and are easy to check by hand.

TABLE 1. Tridegrees (n, s, f) with d_3 -differentials resolved by proof by contradiction.

(5, 47, 10)	(6, 34, 10)	(6, 37, 10)	(6, 42, 14)	(6, 43, 10)
(6, 45, 14)	(6, 48, 7)	(7, 47, 16)	(10, 38, 10)	(10, 41, 10)
(10, 46, 14)	(10, 47, 10)	(10, 49, 14)	(11, 51, 10)	(13, 51, 9)
(14, 36, 7)	(14, 47, 11)			

Example 4.1. There is a possible d_3 -differential on the unique generator x of $\mathcal{U}_3^{6,37,10}$. If this differential were zero then x would survive to $\mathcal{U}_4^{6,37,10}$ where we would have a d_4 -differential $d_4(H(x)) = H(y)$ on $\mathcal{U}_4^{11,32,9}$ where y is the unique generator of $\mathcal{U}_4^{6,36,14}$. Naturality of H would then imply that $d_4(x) = y$ but this is impossible since $y \circ h_1$ is not a boundary for degree reasons so y can not be a boundary. Therefore our initial assumption that the d_3 -differential on $\mathcal{U}_3^{6,37,10}$ is trivial must be false so $d_3(x)$ is equal to the unique generator of $\mathcal{U}_3^{6,36,13}$.

4.3. Why graphs. Clicking a differential in our interactive Adams charts opens the flow chart of its proof. Zero differentials are not drawn in the charts, but their flow charts can be queried directly with the `why.py` script in [\[Bae26\]](#).

4.4. Higher differentials. The unstable Adams charts in [\[Bae26\]](#) present the results of our computation through the E_8 -page. In our range of interest, there may be nontrivial d_r -differentials for $r > 8$. In practice, the few possibilities for such differentials all target v_1 -periodic elements in the unstable Adams spectral sequence. The behavior of the unstable Adams spectral sequence in this v_1 -periodic region was studied in [\[DM90\]](#), though we have not verified all of the details of their arguments. In forthcoming joint work with Balderrama and Isaksen, we study the 2-primary EHP spectral sequence from the perspective of v_1 -periodic elements [\[BBI\]](#). Both of these perspectives rule out the possibility of long differentials hitting v_1 -periodic elements in the first 50 stems.

4.5. Unstable Adams charts. The results of our computation are displayed graphically as Adams charts in the GitHub repository [Bae26]. The horizontal axis is the stem, the vertical axis is the Adams filtration, and each node in a bidegree is an $\mathbb{F}_2$ -basis element of the corresponding bigraded Ext group. Recall that we write $[\lambda_I]$ for the class whose cycle representative in the homology of the lambda algebra has leading term λ_I . Vertical lines correspond to right multiplication by $[\lambda_0]$, given by unstable algebraic Yoneda composition, carrying a class $[\lambda_I]$ to $[\lambda_I] \circ [\lambda_0]$. The lines of slope 1 and slope 1/3 correspond to algebraic right multiplication by $[\lambda_1]$ and $[\lambda_3]$, respectively (see Figure 9).




FIGURE 9. The structural product lines. From a class $[\lambda_I]$, a vertical line records right multiplication by $[\lambda_0]$, a slope-1 line records right multiplication by $[\lambda_1]$, and a slope-1/3 line records right multiplication by $[\lambda_3]$.

Solid negative-slope lines correspond to unstable Adams differentials, with $d_2, d_3, d_4, d_5, d_6, d_7$, and d_8 drawn in teal, red, green, navy, yellow, peach, and mauve, respectively. Dashed negative-slope lines of the same colors indicate possible values of a d_r -differential that our algorithm was able neither to prove nor to disprove. If a basis element on the E_r -page supports neither a solid line nor a dashed line, then the d_r -differential on that element is proved to be zero.

Our graphical calculus does leave some room for ambiguity. For example, the display in Figure 10 could indicate that either $d_r(x) \in \{0, y + z\}$ or $d_r(x) \in \{y, z\}$. There are other possibilities as well. In such cases the raw CSV data records the possible values of each differential unambiguously.




FIGURE 10. An ambiguous display: the reading may be $d_r(x) \in \{0, y + z\}$ or $d_r(x) \in \{y, z\}$; the raw data resolves which.

4.6. Bounds on the orders of the 2-primary unstable homotopy groups of spheres. Tables 2 to 5 record the bounds on the orders of the 2-primary unstable homotopy groups of spheres proven by our computation, and are read as follows. The entry in column n and row s bounds the order of the 2-primary component of $\pi_{n+s}(S^n)$, recorded through its base-2 logarithm: a single entry e marks a homotopy group with no remaining unknown differentials, whose 2-primary component has order exactly 2^e . When a homotopy group still admits unknown differentials we can only bracket this order, and the entry is a range e/ℓ printed upper bound first: the 2-primary component of $\pi_{n+s}(S^n)$ then has order 2^m for some $\ell \leq m \leq e$, so its order lies between the lower bound 2^ℓ and the upper bound 2^e . An entry of ∞ marks a homotopy group containing a $\mathbb{Z}$ summand. The pink staircase boundary traces the metastable range: the entries lying on or to the right of it, up to the stable line, are in the metastable range $s \leq 3n - 4$, while those to its left are purely unstable. Finally, in the stable range $n \geq s + 2$ every entry in row s records the order of the stable s -stem, so rather than reiterate it we print it once, at $n = s + 2$, and draw the repetitions as a vertical line running down each column.

TABLE 2. Bounds on the order of the 2-primary component of the s -stem of S^n , recorded as a base-2 logarithm (upper/lower), $n = 2, \dots, 14$.

$s \backslash n$	2	3	4	5	6	7	8	9	10	11	12	13	14
0	∞												
1	∞	1											
2	1	1	1										
3	1	2	∞	3									
4	2	1	2	1	0								
5	1	1	2	1	∞	0							
6	1	0	3	1	1	1	1						
7	0	0	0	1	2	3	∞	4					
8	0	1	1	1	4	3	4	3	2				
9	1	2	3	3	3	4	5	4	∞	3			
10	2	3	6	4	4	4	7	4	3	2	1		
11	3	4	7	5	5	4	4	4	3	3	∞	3	
12	4	2	6	3	4	0	0	0	2	1	2	1	0
13	2	1	5	2	1	1	2	1	1	2	2	1	∞
14	1	1	5	2	3	5	9	6	5	5	7	5	4
15	1	1	1	2	3	6	8	7	6	5	5	6	6
16	1	2	3	2	5	4	7	4	5	1	1	1	4
17	2	4	9	4	4	4	5	4	3	3	4	4	4
18	4	4	10	5	5/4	4/3	7/6	4	5	6	10	7	7
19	4	3	7	4	8/6	4/3	4/3	4	4	6	8	6	6
20	3	2	6	3	7/6	3	3	3	6	5	8	6	7
21	2	1	5/4	2	1	2	5	3	3	4	5/4	5	4
22	1	1	5/4	3	5	6	12	7	6/5	6/5	9/7	6	7
23	1	2	5	5	11/10	10/9	13/12	11/10	11/10	9/8	9/8	9	9
24	2	3	5	4	9/8	7/5	12/8	7/6	10	5	4	4	7
25	3	5	11	7	10	8/7	15/12	8	6	6	7	5	5
26	5	5	15/14	8	11	9	14	9	8	7	13/11	7	5
27	5	5	12/10	7	10	7/6	10/9	7	8	8	9/7	7	5
28	5	4	12/11	5	9/8	4/3	7/6	5	7	6	8	4	6
29	4	3	12	6	6/5	7	14	8	9	6	8/7	5/4	3/2
30	3	3	10/9	6	8	9	20	10	10/9	7	9/8	7/6	9/7
31	3	4	8/7	5	10	7	13	8	9/8	5	7	6	10/9
32	4	4	11	5	11	6	12	6	10/9	4	5	5	8
33	4	3	12	4	8	6	11	6	5/4	7	8/7	8	9
34	3	4	11	5	9/8	6	9	6	6	7	14/12	8	9
35	4	4	10	5	11/9	6/5	7/6	6	10	7	10/9	8	8
36	4	3	9	4	6/5	2/1	5/3	2	7	4	7	4	9/8
37	3	2	8	4	4	3	12/9	3	5	4	9/8	5	5/4
38	2	1	7/6	4	9	8	19/17	9	11	9	13/12	10	9
39	1	2	4/3	4	10	10	18/17	11	14	8	9	8	9
40	2	3	6	4	10/9	10/9	19/17	10	12	4	5	4	7
41	3	4	12	6	8/7	9/8	18/16	9	9/8	6	11/10	5	7/6
42	4	6	16	9	11	10	19/17	10	10/9	9/8	17/14	7/6	8/7
43	6	7	17/16	9	14	9	14/13	9	12/11	7/6	12/10	6/5	8/7
44	7	5	14/12	5	8	4	11	5	12/10	5/4	10/9	5	9/8
45	5	3	13/12	4	3/2	3	13	5	7/6	5/4	10/8	6	6/5
46	3	1	10	5	5/4	6	15	9	12/11	8	13/11	11/10	13/11
47	1	1	5	5	10/9	10/9	16/15	12	16/14	11/10	12/9	13/11	19/18
48	1	4	7	6/5	12/11	11/10	17/16	11/10	12/10	8/6	11/9	8/7	15
49	4	6	12/11	7/5	13/10	8/7	15/14	8/6	8/5	8/5	16/12	9/7	11/8
50	6/5	7	17/14	8/6	12/7	5/4	9/8	5/4	10/8	11/9	23/14	13/11	15/12

TABLE 3. Bounds on the order of the 2-primary component of the s -stem of S^n , recorded as a base-2 logarithm (upper/lower), $n = 15, \dots, 26$.

$s \backslash n$	15	16	17	18	19	20	21	22	23	24	25	26
0												
1												
2												
3												
4												
5												
6												
7												
8												
9												
10												
11												
12												
13	0											
14	3	2										
15	6	∞	6									
16	3	4	3	2								
17	5	6	5	∞	4							
18	7	10	7	6	5	4						
19	5	5	5	4	4	∞	4					
20	3	3	3	5	4	5	4	3				
21	3	4	3	3	4	4	3	∞	2			
22	7	11	7	6	6	8	5	4	3	2		
23	11	13	11	10	9	9	9	8	8	∞	8	
24	6	9	6	7	3	2	2	4	3	4	3	2
25	6	7	6	4	3	2	2	2	3	4	3	∞
26	5	8	5	5	5	8	5	5	5	8	5	4
27	3	3	3	3	5	7	5	5	4	4	4	3
28	1	1	1	4	3	6	4	5	1	1	1	3
29	3/2	5/4	2	2	3	4	4	3	2	2	1	1
30	9/6	15/12	8	6	6	9	6	7	7	10/9	6	5
31	11/9	13/11	11	10	8	8	8	8	10	12/11	10	9
32	8/7	12/10	8	11	6	4	4	7	6	9	6	7
33	9/8	13/11	9	7	7	7	5	5	6	7	6	5
34	9/8	13/11	9/8	9	8	14/12	8	6	6	9	6	7
35	9/7	12/9	9/8	10	10	11/9	9	7	5	5	5	5
36	5/4	7/6	5	7	6	8	4	6	1	1	1	4
37	7	9	7	7	5	7	4	3	3	5	3	3
38	10	18	10	8	6	8	6	8/7	8/7	14/13	8	7/6
39	9	11	9	7	7	9	8	11/10	12/11	14/13	12	12/11
40	6	8	6	11/10	7	9	9/8	12/11	12/10	16/14	12	15
41	6	8	7	7/6	9	10/9	10/9	11/10	11/9	15/12	11	9
42	7	10	8	8	9	16/14	10	11	11/10	15/12	11/10	11
43	4	5	4	5	5	7/6	6	7/6	7/5	10/7	7/6	8
44	3	3	3	6	5	7/6	5	11/9	6/5	8/7	7	9
45	7	9	8	9	10	12/11	11	11/10	13	15	14	15
46	13/12	21/19	13/11	12/11	13/12	16/15	14	14	15	23/22	15	14
47	17/16	22/20	17/15	15/14	14/13	15/14	14	15	15	17/16	15	12
48	14	20/19	14	17/16	9/8	8	8	10	9	11	9	13/12
49	12/9	20/15	12/9	10/8	7/6	7/6	6/5	5	5	7	5	5/4
50	12/9	18/13	12/9	10/9	9	16/13	8/7	6	6	9	6	6/5

TABLE 4. Bounds on the order of the 2-primary component of the s -stem of S^n , recorded as a base-2 logarithm (upper/lower), $n = 27, \dots, 38$.

$s \backslash n$	27	28	29	30	31	32	33	34	35	36	37	38
0												
1												
2												
3												
4												
5												
6												
7												
8												
9												
10												
11												
12												
13												
14												
15												
16												
17												
18												
19												
20												
21												
22												
23												
24												
25	2											
26	3	2										
27	3	∞	3									
28	2	3	2	1								
29	2	2	1	∞	0							
30	5	7	4	3	2	1						
31	8	8	8	8	8	∞	8					
32	3	3	3	6	5	6	5	4				
33	5	5	5	5	6	7	6	∞	5			
34	8	11	8	8	8	11	8	7	6	5		
35	7	9	7	7	6	6	6	5	5	∞	5	
36	3	6	4	5	1	1	1	3	2	3	2	1
37	4	5	5	4	3	4	3	3	4	4	3	∞
38	7/6	10/9	7	8	8	12	8	7	7	9	6	5
39	10/9	10/9	10	10	12	14	12	11	10	10	10	9
40	10	9	9	12	11	14	11	12	8	7	7	9
41	9	10	8	8	9	10	9	7	6	5	5	5
42	10	16/14	10	8	8	11	8	8	8	11/10	8	8
43	8	9/7	7	5	3	3	3	3	5	7/6	5	5
44	8	10	6	8	3	3	3	6	5	8	6	7
45	13	15	11	9	9	11	9	9	10	11	11	10
46	12	14	10	11/10	11/9	17/14	11/9	9/8	9/8	12/11	9	10
47	11	13	11	13/12	14/12	16/13	14/12	13/12	11/10	11/10	11	11
48	8	10/9	10	12	12/11	16/15	12	15	10	8	8	11
49	7	8/6	8	9	9/8	13/11	9	7	7	7	5	5
50	7	15/12	8	9/8	9/7	13/9	9/7	9/8	8/7	14/12	8	6

TABLE 5. Bounds on the order of the 2-primary component of the s -stem of S^n , recorded as a base-2 logarithm (upper/lower), $n = 39, \dots, 50$.

$s \backslash n$	39	40	41	42	43	44	45	46	47	48	49	50
0												
1												
2												
3												
4												
5												
6												
7												
8												
9												
10												
11												
12												
13												
14												
15												
16												
17												
18												
19												
20												
21												
22												
23												
24												
25												
26												
27												
28												
29												
30												
31												
32												
33												
34												
35												
36												
37	2											
38	4	3										
39	9	∞	9									
40	8	9	8	7								
41	6	7	6	∞	5							
42	8	11	8	7	6	5						
43	4	4	4	3	3	∞	3					
44	3	3	3	5	4	5	4	3				
45	9	9	8	8	9	9	8	∞	7			
46	10	13/12	9	8	8	10	7	6	5	4		
47	13	15/14	13	12	11	11	11	10	10	∞	10	
48	10	13	10	11	7	6	6	8	7	8	7	6
49	6	7	6	4	3	2	2	2	3	4	3	∞
50	6	9	6	6	6	9/8	6	6	6	9	6	5

REFERENCES

- [Bae26] Jake Francis Baer. *Adams-EHP: automated proofs of unstable Adams differentials*. Zenodo. Version 1.0.0. Code and data; development repository at <https://github.com/jfbaer/adams-ehp>. 2026. DOI: [10.5281/zenodo.22820890](https://doi.org/10.5281/zenodo.22820890).
- [BBBI] Jake Francis Baer, William Balderrama, Eva Belmont, and Dan Isaksen. “The metastable homotopy of spheres in stems 0 through 66”. In preparation.
- [BBI] Jake Francis Baer, William Balderrama, and Dan Isaksen. “The generalized Kervaire problem in algebraic topology”. In preparation.
- [Bea22] Joey Beauvais-Feisthauer. “Automated differential computation in the Adams spectral sequence”. In: *arXiv preprint arXiv:2210.15169* (Oct. 2022). DOI: [10.48550/arXiv.2210.15169](https://doi.org/10.48550/arXiv.2210.15169). arXiv: [2210.15169](https://arxiv.org/abs/2210.15169) [math.AT].
- [BI25] Joey Beauvais-Feisthauer and Dan Isaksen. “SeqSee: A schema-based approach to spectral sequence visualization”. In: *arXiv preprint arXiv:2501.18429* (Jan. 2025). DOI: [10.48550/arXiv.2501.18429](https://doi.org/10.48550/arXiv.2501.18429). arXiv: [2501.18429](https://arxiv.org/abs/2501.18429) [math.AT].
- [BM24] Mark Behrens and Connor Malin. “Unstable homotopy groups and Lie algebras”. In: *arXiv preprint arXiv:2410.18330* (2024). DOI: [10.48550/arXiv.2410.18330](https://doi.org/10.48550/arXiv.2410.18330). arXiv: [2410.18330](https://arxiv.org/abs/2410.18330) [math.AT].
- [BC70] A. K. Bousfield and E. B. Curtis. “A spectral sequence for the homotopy of nice spaces”. In: *Trans. Amer. Math. Soc.* 151.2 (1970), pp. 457–479. ISSN: 0002-9947. DOI: [10.1090/S0002-9947-1970-0267586-7](https://doi.org/10.1090/S0002-9947-1970-0267586-7).
- [Bou+66] A. K. Bousfield, E. B. Curtis, D. M. Kan, D. G. Quillen, D. L. Rector, and J. W. Schlesinger. “The mod $-p$ lower central series and the Adams spectral sequence”. In: *Topology* 5 (1966), pp. 331–342. ISSN: 0040-9383. DOI: [10.1016/0040-9383\(66\)90024-3](https://doi.org/10.1016/0040-9383(66)90024-3).
- [BK73a] A. K. Bousfield and D. M. Kan. “A second quadrant homotopy spectral sequence”. In: *Trans. Amer. Math. Soc.* 177 (1973), pp. 305–318. ISSN: 0002-9947. DOI: [10.1090/S0002-9947-1973-0372859-6](https://doi.org/10.1090/S0002-9947-1973-0372859-6).
- [BK73b] A. K. Bousfield and D. M. Kan. “Pairings and products in the homotopy spectral sequence”. In: *Trans. Amer. Math. Soc.* 177 (1973), pp. 319–343. ISSN: 0002-9947. DOI: [10.1090/S0002-9947-1973-0372860-2](https://doi.org/10.1090/S0002-9947-1973-0372860-2).
- [Coh83] Frederick R. Cohen. “The unstable decomposition of $\Omega^2\Sigma^2X$ and its applications”. In: *Math. Z.* 182.4 (1983), pp. 553–568. ISSN: 0025-5874. DOI: [10.1007/BF01215483](https://doi.org/10.1007/BF01215483). URL: <https://doi.org/10.1007/BF01215483>.
- [Cur71] Edward B. Curtis. “Simplicial homotopy theory”. In: *Adv. Math.* 6.2 (Apr. 1971), pp. 107–209. ISSN: 0001-8708. DOI: [10.1016/0001-8708\(71\)90015-6](https://doi.org/10.1016/0001-8708(71)90015-6).
- [CGMM87] Edward B. Curtis, Paul Goerss, Mark Mahowald, and R. James Milgram. “Calculations of unstable Adams E2 terms for spheres”. In: *Algebraic Topology*. Ed. by Haynes R. Miller and Douglas C. Ravenel. Vol. 1286. Lecture Notes in Mathematics. Berlin, Heidelberg: Springer, 1987, pp. 208–266. ISBN: 978-3-540-47986-4. DOI: [10.1007/BFb0078744](https://doi.org/10.1007/BFb0078744).
- [CM89] Edward B. Curtis and Mark Mahowald. “The unstable Adams spectral sequence for S^3 ”. In: *Algebraic topology (Evanston, IL, 1988)*. Vol. 96. Contemp. Math. Providence, RI: Amer. Math. Soc., 1989, pp. 125–162. ISBN: 0-8218-5102-0. DOI: [10.1090/conm/096/1022678](https://doi.org/10.1090/conm/096/1022678). URL: <https://doi.org/10.1090/conm/096/1022678>.
- [DM90] Donald M. Davis and Mark Mahowald. “ v_1 -periodicity in the unstable Adams spectral sequence”. In: *Math. Z.* 204.3 (1990), pp. 319–339. ISSN: 0025-5874.
- [Han74] William A. Hansen. “A Computer Calculation of the Homology of the Lambda-Algebra”. PhD thesis. Northwestern University, 1974, p. 179.
- [HM89] J. R. Harper and H. R. Miller. “Looping Massey–Peterson towers”. In: *Advances in Homotopy Theory*. Vol. 139. London Math. Soc. Lecture Note Ser. Cambridge University Press, 1989, pp. 69–86.
- [Hop31] Heinz Hopf. “Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche”. In: *Math. Ann.* 104 (1931), pp. 637–665.
- [IM12] Tomohisa Inoue and Juno Mukai. “The 2-components of the 31-stem homotopy groups of the 9 and 10-spheres”. In: *arXiv preprint arXiv:1210.0995* (Oct. 2012). Published as: Inoue, Miyauchi and Mukai, J. Fac. Sci. Shinshu Univ. **46** (2015), 1–19. DOI: [10.48550/arXiv.1210.0995](https://doi.org/10.48550/arXiv.1210.0995). arXiv: [1210.0995](https://arxiv.org/abs/1210.0995) [math.AT].
- [IWX23] Daniel C. Isaksen, Guozhen Wang, and Zhouli Xu. “Stable homotopy groups of spheres: from dimension 0 to 90”. In: *Publ. Math. Inst. Hautes Études Sci.* 137 (2023), pp. 107–243. DOI: [10.1007/s10240-023-00139-1](https://doi.org/10.1007/s10240-023-00139-1).

- [IMW18] Sergei O. Ivanov, Roman Mikhailov, and Jie Wu. “Leibniz rule on higher pages of unstable spectral sequences”. In: *Proc. Edinb. Math. Soc. (2)* 61.1 (Feb. 2018), pp. 265–282. ISSN: 0013-0915. DOI: [10.1017/S0013091517000220](https://doi.org/10.1017/S0013091517000220).
- [Lie72] Martin Liebowitz. “The E1-term of the lower central series spectral sequence for the homotopy of spaces”. Ph.D. thesis. Brandeis University, 1972.
- [LWX24] Weinan Lin, Guozhen Wang, and Zhouli Xu. “Machine proofs for Adams differentials and extension problems among CW spectra”. In: *arXiv preprint arXiv:2412.10876* (2024). DOI: [10.48550/arXiv.2412.10876](https://doi.org/10.48550/arXiv.2412.10876). arXiv: [2412.10876](https://arxiv.org/abs/2412.10876) [math.AT].
- [LWX25] Weinan Lin, Guozhen Wang, and Zhouli Xu. *Machine proofs for Adams differentials and extension problems among CW spectra*. Zenodo dataset. Version v126.3.cw49. Concept DOI [10.5281/zenodo.14272279](https://doi.org/10.5281/zenodo.14272279) resolves to the latest version. 2025. DOI: [10.5281/zenodo.14875701](https://doi.org/10.5281/zenodo.14875701). URL: <https://doi.org/10.5281/zenodo.14875701>.
- [Mah75] Mark Mahowald. “On the double suspension homomorphism”. In: *Trans. Amer. Math. Soc.* 214 (1975), pp. 169–178. ISSN: 0002-9947. DOI: [10.1090/S0002-9947-1975-0438333-5](https://doi.org/10.1090/S0002-9947-1975-0438333-5).
- [Mah82] Mark Mahowald. “The image of J in the EHP sequence”. In: *Ann. of Math. (2)* 116.1 (July 1982), pp. 65–112. ISSN: 0003-486X. DOI: [10.2307/2007048](https://doi.org/10.2307/2007048). URL: <http://www.jstor.org/stable/2007048>.
- [MT87] Mark Mahowald and Robert Thompson. “A commentary on: “Image of J in the EHP sequence” [Ann. of Math. (2) **116** (1982), no. 1, 65–112; MR0662118 (83i:55019)] by Mahowald”. In: *Algebraic topology (Seattle, Wash., 1985)*. Vol. 1286. Lecture Notes in Mathematics. Berlin: Springer, 1987, pp. 298–304. ISBN: 3-540-18481-3. DOI: [10.1007/BFb0078748](https://doi.org/10.1007/BFb0078748).
- [MT94] Mark Mahowald and Robert D. Thompson. “The fiber of the secondary suspension map”. In: *Amer. J. Math.* 116.1 (1994), pp. 179–205. ISSN: 0002-9327. DOI: [10.2307/2374985](https://doi.org/10.2307/2374985). URL: <http://www.jstor.org/stable/2374985>.
- [MT95] Mark Mahowald and Robert D. Thompson. “The EHP sequence and periodic homotopy”. In: *Handbook of algebraic topology*. Ed. by I. M. James. Amsterdam: North-Holland, 1995, pp. 397–423. ISBN: 0-444-81779-4. DOI: [10.1016/B978-044481779-2/50011-0](https://doi.org/10.1016/B978-044481779-2/50011-0). URL: <https://doi.org/10.1016/B978-044481779-2/50011-0>.
- [MP67] W. S. Massey and F. P. Peterson. *The mod 2 cohomology structure of certain fibre spaces*. Mem. Amer. Math. Soc. 74. Providence, RI: American Mathematical Society, 1967.
- [Mim64] Mamoru Mimura. “On the generalized Hopf homomorphism and the higher composition. I”. In: *J. Math. Kyoto Univ.* 4.1 (1964), pp. 171–190. ISSN: 0023-608X. DOI: [10.1215/kjm/1250524712](https://doi.org/10.1215/kjm/1250524712).
- [Mim65] Mamoru Mimura. “On the generalized Hopf homomorphism and the higher composition. II. $\pi_{n+i}(S^n)$ for $i = 21$ and 22 ”. In: *J. Math. Kyoto Univ.* 4.2 (1965), pp. 301–326. ISSN: 0023-608X. DOI: [10.1215/kjm/1250524662](https://doi.org/10.1215/kjm/1250524662).
- [MMO75] Mamoru Mimura, Masamitsu Mori, and Nobuyuki Oda. “Determination of the 2-components of the 23- and 24-stems in homotopy groups of spheres”. In: *Mem. Fac. Sci. Kyushu Univ. Ser. A* 29.1 (1975), pp. 1–42. ISSN: 0373-6385. DOI: [10.2206/kyushumfs.29.1](https://doi.org/10.2206/kyushumfs.29.1).
- [MT63] Mamoru Mimura and Hiroshi Toda. “The $(n + 20)$ -th homotopy groups of n -spheres”. In: *J. Math. Kyoto Univ.* 3.1 (1963), pp. 37–58. ISSN: 0023-608X. DOI: [10.1215/kjm/1250524854](https://doi.org/10.1215/kjm/1250524854).
- [MM17] Toshiyuki Miyauchi and Juno Mukai. “Determination of the 2-primary components of the 32-stem homotopy groups of S^n ”. In: *Bol. Soc. Mat. Mex. (3)* 23.1 (2017), pp. 319–387. ISSN: 1405-213X. DOI: [10.1007/s40590-016-0154-2](https://doi.org/10.1007/s40590-016-0154-2).
- [Oda76] Nobuyuki Oda. “Hopf invariants in metastable homotopy groups of spheres”. In: *Mem. Fac. Sci. Kyushu Univ. Ser. A* 30.2 (1976), pp. 221–246. ISSN: 0373-6385. DOI: [10.2206/kyushumfs.30.221](https://doi.org/10.2206/kyushumfs.30.221).
- [Oda77] Nobuyuki Oda. “On the 2-components of the unstable homotopy groups of spheres I, II”. In: *Proc. Japan Acad. Ser. A Math. Sci.* 53 (1977), pp. 202–205, 215–218.
- [PW19] Fedor Pavutnitskiy and Jie Wu. “A simplicial James–Hopf map and decompositions of the unstable Adams spectral sequence for suspensions”. In: *Algebr. Geom. Topol.* 19.1 (Feb. 2019), pp. 77–108. ISSN: 1472-2747. DOI: [10.2140/agt.2019.19.77](https://doi.org/10.2140/agt.2019.19.77). URL: <http://dx.doi.org/10.2140/agt.2019.19.77>.
- [Pon38] Lev Pontrjagin. “Classification of continuous maps of a complex into a sphere, Communication II”. In: *Dokl. Akad. Nauk SSSR* 19 (1938), pp. 361–363.
- [Pri70] Stewart B. Priddy. “On the homotopy theory of simplicial Lie algebras”. In: *Proc. Amer. Math. Soc.* 25.3 (1970), pp. 513–517. ISSN: 0002-9939. DOI: [10.2307/2036633](https://doi.org/10.2307/2036633). URL: <http://www.jstor.org/stable/2036633>.

- [Rav86] Douglas C. Ravenel. *Complex cobordism and stable homotopy groups of spheres*. Vol. 121. Pure and Applied Mathematics. Orlando, FL: Academic Press, 1986.
- [Ser53] Jean-Pierre Serre. “Groupes d’homotopie et classes de groupes abéliens”. In: *Ann. of Math.* (2) 58 (1953), pp. 258–294. ISSN: 0003-486X.
- [Sin75] William M. Singer. “The algebraic EHP sequence”. In: *Trans. Amer. Math. Soc.* 201 (1975), pp. 367–382. ISSN: 0002-9947. DOI: [10.1090/S0002-9947-1975-0385861-7](https://doi.org/10.1090/S0002-9947-1975-0385861-7).
- [Tan85] Martin C. Tangora. *Computing the homology of the lambda algebra*. Vol. 58. Memoirs of the American Mathematical Society 337. Providence, RI: American Mathematical Society, 1985. ISBN: 978-0-8218-2338-5. DOI: [10.1090/memo/0337](https://doi.org/10.1090/memo/0337).
- [Tan93] Martin C. Tangora. “Some remarks on products in Ext”. In: *Algebraic topology (Oaxtepec, 1991)*. Vol. 146. Contemp. Math. Providence, RI: Amer. Math. Soc., 1993, pp. 419–429. ISBN: 978-0-8218-5162-3. DOI: [10.1090/conm/146/01237](https://doi.org/10.1090/conm/146/01237).
- [Tan94] Martin C. Tangora. “Some Massey products in Ext”. In: *Topology and representation theory (Evanston, IL, 1992)*. Vol. 158. Contemp. Math. Providence, RI: Amer. Math. Soc., 1994, pp. 269–280. ISBN: 0-8218-5165-9. DOI: [10.1090/conm/158/01463](https://doi.org/10.1090/conm/158/01463).
- [The24] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 10.4)*. <https://www.sagemath.org>. 2024.
- [Tho90] Robert D. Thompson. “The v_1 -periodic homotopy groups of an unstable sphere at odd primes”. In: *Trans. Amer. Math. Soc.* 319.2 (1990), pp. 535–559. ISSN: 0002-9947. DOI: [10.2307/2001253](https://doi.org/10.2307/2001253). URL: <https://doi.org/10.2307/2001253>.
- [Tod62] Hirosi Toda. *Composition methods in homotopy groups of spheres*. Annals of Mathematics Studies 49. Princeton, NJ: Princeton University Press, 1962, p. 193. ISBN: 978-0-691-09586-8.
- [WX16] Guozhen Wang and Zhouli Xu. “The algebraic Atiyah-Hirzebruch spectral sequence of real projective spectra”. In: *arXiv preprint arXiv:1601.02185* (Jan. 2016). DOI: [10.48550/arXiv.1601.02185](https://doi.org/10.48550/arXiv.1601.02185). arXiv: [1601.02185](https://arxiv.org/abs/1601.02185) [math.AT].
- [Whi50] George W. Whitehead. “The $(n+2)^{\text{nd}}$ homotopy group of the n -sphere”. In: *Ann. of Math.* (2) 52 (1950), pp. 245–247.
- [Whi70] George W. Whitehead. *Recent advances in homotopy theory*. Vol. 5. CBMS Regional Conference Series in Mathematics. Providence, RI: American Mathematical Society, 1970, p. 82.