			space					well mixed, local mating			well mixed, global mating		
	Calhoun's pattern		$y=0$†	core	short-lived	C1, adults recover	C1	core	short-lived	C1	core	short-lived	C1
B	Socially functional growth	O10b	1	.12 m.71	1	0	.02 m.97	1	1	1	1	.99	1
C	Peak below $m_c\|G\|$ animals		0	1	1	1	1	0	0	1	0	0	1
	Peak below capacity, empty nests	O4	0	1	.85 m.14	1	1	0	0	0	0	0	0
	Adults deteriorate before peak	O10	0	1	1	1	1	.78 m.23	.61 m.39	1	.18 m.82	.13 m.87	1
	Aggregation with free space	O14	1	1	1	1	1	0	0	0 m1	0	0	0 m1
	Crowded and withdrawn coexist	O13	trunc.	0 m.01	0 m.57	1	1	0	0	0 m1	0	0	.46 m.54
	Withdrawn class persists‡	O13p	trunc.	0	0	0 m1	1	0	0	0	0	0	0
	Withdrawn class born late	O13b	trunc.	n.a.	.01 m.10	0 m.04	0 m.01	n.a.	n.a.	0 m.02	n.a.	n.a.	0 m.01
	Gradual stagnation phase§	O3c	trunc.	1	1	1	1	.09	.02	.52	1	1	1
C–D	Late cohorts fail (age 6 wk)	O11	trunc.	0 m.74	0 m.47	.15 m.34	1	1	1	1	1	.19	1
	Late cohorts fail (age a_w)§	$O11_{a_w}$	trunc.	.76 m.24	.43 m.10	0 m.02	.99	1	1	1	1	.20	1
D	Natality ceases near peak	O5	trunc.	.91 m.07	.23 m.22	0	.99	1	1	1	1	.19	1
	No recovery of natality	O6	trunc.	1	.53	0	1	1	1	1	1	.18	1
	Extinction	O7	trunc.	1	.54	0	1	1	1	1	1	.29	1
	Long senescent decline	O8	trunc.	.64 m.23	.01 m.12	n.a.	.62 m.33	.94 m.07	.01	1	.98 m.02	0	1
	Old female-biased remnant	O9	trunc.	.45 m.49	.14 m.16	n.a.	.45 m.39	.41 m.41	.01	.51 m.49	.46 m.49	0	.43 m.56
	Total infant mortality	O15	trunc.	0 m.30	0 m.42	0 m.16	.01 m.99	0 m1	.34 m.66	.02 m.98	.01 m.99	.04 m.96	.01 m.99
all	**All essential patterns**	O17p	trunc.	**0**	**0**	**0**	**.99**	**0**	**0**	**0**	**0**	**0**	**0**

share of runs that pass

0.0 0.5 1.0

m: fraction MARGINAL (one step from the threshold). Hatched: trunc., runs stopped by the cap on N; n.a., criterion undefined (no run reaches the event). ‡: holds by construction. §: added after the preregistration (O3c: phase-C clause of O3). †: every run stopped at weeks 26–29 (median 27) by the cap on N.