

# Answers to Exercises

## CHAPTER 2

- 2.1. b.  $\Pr\{A \cup B\} = 0.7$   
 c.  $\Pr\{A \cap B^C\} = 0.1$   
 d.  $\Pr\{A^C \cap B^C\} = 0.3$
- 2.2. b.  $\Pr\{A\} = 9/31$ ,  $\Pr\{B\} = 15/31$ ,  $\Pr\{A, B\} = 9/31$   
 c.  $\Pr\{A|B\} = 9/15$   
 d. No:  $\Pr\{A\} \neq \Pr\{A|B\}$
- 2.3 a.  $18/22$   
 b.  $22/31$
- 2.4. b.  $\Pr\{E_1, E_2, E_3\} = .000125$   
 c.  $\Pr\{E_1^C, E_2^C, E_3^C\} = .857$
- 2.5. 0.20

## CHAPTER 3

- 3.1. median = 2 mm, trimean = 2.75 mm, mean = 12.95 mm
- 3.2. MAD = 0.4 mb, IQR = 0.8 mb,  $s = 0.88$  mb
- 3.4.  $\gamma_{YK} = 0.273$ ,  $\gamma = 0.877$
- 3.7.  $\lambda = 0$
- 3.9.  $z = 1.36$
- 3.10.  $r_0 = 1.000$ ,  $r_1 = 0.652$ ,  $r_2 = 0.388$ ,  $r_3 = 0.281$

3.12. Pearson: 
$$\begin{bmatrix} 1.000 & 0.703 & -0.830 \\ 0.703 & 1.000 & -0.678 \\ -0.830 & -0.678 & 1.000 \end{bmatrix}$$
 Spearman : 
$$\begin{bmatrix} 1.000 & 0.606 & -0.688 \\ 0.606 & 1.000 & -0.632 \\ -0.688 & -0.632 & 1.000 \end{bmatrix}$$

## CHAPTER 4

- 4.1. 0.168
- 4.2. a. 0.037  
 b. 0.331

- 4.3. a.  $\mu_{\text{drought}} = 0.056$ ,  $\mu_{\text{wet}} = 0.565$   
 b. 0.054  
 c. 0.432
- 4.4. \$280 million, \$2.825 billion
- 4.5. a.  $\mu = 24.8^\circ\text{C}$ ,  $\sigma = 0.98^\circ\text{C}$   
 b.  $\mu = 76.6^\circ\text{F}$ ,  $\sigma = 1.76^\circ\text{F}$
- 4.6. a. 0.00939  
 b.  $22.9^\circ\text{C}$
- 4.7. a.  $\alpha = 3.785$ ,  $\beta = 0.934''$   
 b.  $\alpha = 3.785$ ,  $\beta = 23.7 \text{ mm}$
- 4.8. a.  $q_{30} = 2.41'' = 61.2 \text{ mm}$ ;  $q_{70} = 4.22'' = 107.2 \text{ mm}$   
 b.  $0.30''$ , or  $7.7 \text{ mm}$   
 c.  $\cong 0.05$
- 4.9. a.  $q_{30} = 2.30'' = 58.3 \text{ mm}$ ;  $q_{70} = 4.13'' = 104.9 \text{ mm}$   
 b.  $0.46''$ , or  $11.6 \text{ mm}$   
 c.  $\cong 0.07$
- 4.10. a.  $\beta = 35.1 \text{ cm}$ ,  $\zeta = 59.7 \text{ cm}$   
 b.  $x = \zeta - \beta \ln [-\ln(F)]$ ;  $\Pr\{X \leq 221 \text{ cm}\} = 0.99$
- 4.11. a.  $\mu_{\text{max}} = 31.8^\circ\text{F}$ ,  $\sigma_{\text{max}} = 7.86^\circ\text{F}$ ,  $\mu_{\text{min}} = 20.2^\circ\text{F}$ ,  $\sigma_{\text{min}} = 8.81^\circ\text{F}$ ,  $\rho = 0.810$   
 b. 0.728
- 4.13. a.  $\beta = \Sigma x/n$   
 b.  $-\Gamma^{-1}(\beta) = \hat{\beta}^2/n$
- 4.14.  $x(u) = \beta [-\ln(1-u)]^{1/\alpha}$

## CHAPTER 5

- 5.1. a.  $z = 4.88$ , reject  $H_0$   
 b.  $[1.10^\circ\text{C}, 2.56^\circ\text{C}]$
- 5.2. 6.53 days (Ithaca), 6.08 days (Canandaigua)
- 5.3.  $z = -4.00$   
 a.  $p = 0.000063$   
 b.  $p = 0.000032$
- 5.4.  $|r| \geq 0.366$
- 5.5. a.  $D_n = 0.152$  (reject at 10%, not at 5% level)  
 b. For classes:  $[\leq 2, 2-3, 3-4, 4-5, \geq 5]$ ,  $\chi^2 = 0.33$  (do not reject)  
 c.  $r = 0.971$  (do not reject)
- 5.6.  $A = 21.86$ , reject ( $p < .001$ )
- 5.7. a.  $U_1 = 1$ , reject ( $p < .005$ )  
 b.  $z = -3.18$ , reject ( $p = .0007$ )
- 5.8.  $\approx [1.02, 3.59]$
- 5.9. a. Observed  $(s^2_{\text{E-N}}/s^2_{\text{non-E-N}}) = 329.5$ ; permutation distribution critical value (1%, 2-tailed)  $\approx 141$ , reject  $H_0$  ( $p < 0.01$ )  
 b. 15/10000 members of bootstrap sampling distribution for  $s^2_{\text{E-N}}/s^2_{\text{non-E-N}} \leq 1$ ; 2-tailed  $p = 0.003$

- 5.10. a. Counting method, no (need  $\geq 3$  locally significant); FDR, yes  
 b.  $p=.007$  and  $p=.009$  significant according to FDR

## CHAPTER 6

- 6.1. a.  $\alpha = 14.8$ ,  $\beta = 7.41$   
 b. Beta distribution, with  $\alpha' = 29.8$ ,  $\beta' = 17.4$   
 c.  $\Pr\{X^+ = 0\} = .0094$ ,  $\Pr\{X^+ = 1\} = .0656$ ,  $\Pr\{X^+ = 2\} = .1982$ ,  $\Pr\{X^+ = 3\} = .3248$   
 $\Pr\{X^+ = 4\} = .2895$ ,  $\Pr\{X^+ = 5\} = .1125$ ,  
 6.2. a.  $\beta = 190.8$ ,  $\zeta = 162.3$   
 b.  $\beta = 155.9$ ,  $\zeta = 180.0$   
 c. 1040.0, 897.2  
 6.3. a.  $\alpha = 1.5$ ,  $\beta = 0.1$   
 b. .157  
 6.4. a.  $\mu'_h = 455.6$ ,  $\sigma'_h = 33.3$   
 b.  $\mu_+ = 455.6$ ,  $\sigma_+ = 60.1$   
 6.5. a.  $\mu'_h = 427.4$ ,  $\sigma'_h = 28.6$   
 b.  $\mu_+ = 427.4$ ,  $\sigma_+ = 57.6$   
 6.6. a. 462.3  
 b. 400  
 c. 450

## CHAPTER 7

- 7.1. a.  $a = 959.8^\circ\text{C}$ ,  $b. = -0.925^\circ\text{C}/\text{mb}$   
 c.  $z = -6.33$   
 d. 0.690  
 e. 0.876  
 f. 0.925  
 7.2. a. 3  
 b. 117.9  
 c. 0.974  
 d. 0.715  
 7.3.  $\ln [\bar{y}/(1-\bar{y})]$   
 7.4. a. 1.74 mm  
 b. [0 mm, 13.1 mm]  
 7.5. Range of slopes,  $-0.850$  to  $-1.095$ ;  $\text{MSE} = 0.369$   
 7.6. a.  $-59$  n.m.  
 b.  $-66$  n.m.  
 7.7. a.  $65.8^\circ\text{F}$   
 b.  $52.5^\circ\text{F}$   
 c.  $21.7^\circ\text{F}$   
 d.  $44.5^\circ\text{F}$

- 7.8. a. 0.65  
 b. 0.49  
 c. 0.72  
 d. 0.56
- 7.9.  $f_{\text{MOS}} = 30.8^\circ\text{F} + (0) (\text{Th})$
- 7.10. 0.20
- 7.11. a. 12 mm  
 b. [5 mm, 32 mm], [1 mm, 55 mm]  
 c. 0.625

## CHAPTER 8

- 8.1. a. .0025 .0013 .0108 .0148 .0171 .0138 .0155 .0161 .0177 .0176 .0159 .0189  
 .4087 .0658 .1725 .0838 .0445 .0228 .0148 .0114 .0068 .0044 .0011 .0014  
 b. 0.162
- 8.2. 1644 1330  
 364 9064
- 8.3. a. 0.863  
 b. 0.493  
 c. 0.578  
 d. 0.691  
 e. 0.407
- 8.4. a. 0.074  
 b. 0.097  
 c. 0.761  
 d. 0.406
- 8.5. a. 0.9597 .0127 .0021 .0007  
 .0075 .0043 .0014 .0005  
 .0013 .0013 .0009 .0003  
 .0007 .0006 .0049 .0009  
 b. 0.966  
 c. 0.369  
 d. 0.334
- 8.6. a.  $5.37^\circ\text{F}$   
 b.  $7.54^\circ\text{F}$   
 c.  $-0.03^\circ\text{F}$   
 d. 1.95%
- 8.7. a. 0.1215  
 b. 0.1699  
 c. 28.5%

- 8.8. a. .0415 .0968 .1567 .1428 .1152 .0829 .1060 .0829 .0783 .0553 .0415  
       .3627 .2759 .1635 .0856 .0498 .0230 .0204 .0102 .0051 .0026 .0013  
 c.  $H = .958, .862, .705, .562, .447, .364, .258, .175, .097, .042$   
 $F = .637, .361, .198, .112, .062, .039, .019, .009, .004, .001$   
 d.  $A = 0.831, z = -14.9$
- 8.9. a. 0.298  
 b. 16.4%  
 c. 0.755
- 8.10. a. 30.3  
 b. 5.31 dam<sup>2</sup>  
 c. 46.9%  
 d. 0.726  
 e. 0.714
- 8.11. a. 5 rank 1, 2 rank 2, 3 rank 3, 2 rank 4, 2 rank 5, 6 rank 6  
 b. underdispersed
- 8.12. .352, .509, .673, .598, .504, .426, .343, .275, .195, .128, -.048

## CHAPTER 9

- 9.1. a.  $p_{01} = 0.45, p_{11} = 0.79$   
 b.  $\chi^2 = 3.51, p \approx 0.064$   
 c.  $\pi_1 = 0.682, n_{\bullet 1}/n = 0.667$   
 d.  $r_0 = 1.00, r_1 = 0.34, r_2 = 0.12, r_3 = 0.04$   
 e. 0.624
- 9.2. a.  $r_0 = 1.00, r_1 = 0.40, r_2 = 0.16, r_3 = 0.06, r_4 = 0.03, r_5 = 0.01$   
 a.  $r_0 = 1.00, r_1 = 0.41, r_2 = -0.41, r_3 = -0.58, r_4 = -0.12, r_5 = 0.32$
- 9.3. a. AR(1):  $\phi = 0.80; s^2_\varepsilon = 36.0$   
 AR(2):  $\phi_1 = 0.89, \phi_2 = -0.11; s^2_\varepsilon = 35.5$   
 AR(3):  $\phi_1 = 0.91, \phi_2 = -0.25, \phi_3 = 0.16; s^2_\varepsilon = 34.7$   
 b. AR(1): BIC = 369.6  
 c. AR(1): AIC = 364.4
- 9.4.  $x_1 = 71.5, x_2 = 66.3, x_3 = 62.1$
- 9.5. a. 28.6  
 b. 19.8  
 c. 4.5
- 9.6. a.  $C_1 = 16.92^\circ\text{F}, \phi_1 = 199^\circ; C_2 = 4.16^\circ\text{F}, \phi_2 = 256^\circ$
- 9.7. a.  $82.0^\circ\text{F}$   
 b.  $74.8^\circ\text{F}$
- 9.8. b. 0.990
- 9.9. 56
- 9.10. a. e.g.,  $f_A = 1 - .0508 \text{ mo}^{-1} = .9492 \text{ mo}^{-1}$   
 b.  $\approx$  twice monthly
- 9.12. a. [0.11, 16.3]  
 b.  $C_{11}^2 < 0.921$ , do not reject

**CHAPTER 10**

10.1.  $\begin{bmatrix} 216.0 & -4.32 \\ 135.1 & 7.04 \end{bmatrix}$

10.2.  $([X]^T \mathbf{y})^T = [627, 11475]$ ,  $[X^T X]^{-1} = \begin{bmatrix} .06263 & -.002336 \\ -.002336 & .0001797 \end{bmatrix}$ ,  $\mathbf{b}^T = [12.46, 0.60]$

10.3.  $90^\circ$

10.6. a.  $\begin{bmatrix} 59.5 & 58.1 \\ 58.1 & 61.8 \end{bmatrix}$

b.  $\begin{bmatrix} .205 & -.193 \\ -.193 & .197 \end{bmatrix}$

c.  $\begin{bmatrix} .205 & -.193 \\ -.193 & .197 \end{bmatrix}$

d.  $\begin{bmatrix} 6.16 & 4.64 \\ 4.64 & 6.35 \end{bmatrix}$

e. 1.765

10.7. a.  $\begin{bmatrix} 59.52 & 75.43 & 58.07 & 51.70 \\ 75.43 & 185.47 & 81.63 & 110.80 \\ 58.07 & 81.63 & 61.85 & 56.12 \\ 51.70 & 110.80 & 56.12 & 77.58 \end{bmatrix}$

b.  $\boldsymbol{\mu}_y^T = [21.4, 26.0]$

$[S_y] = \begin{bmatrix} 98.96 & 75.55 \\ 75.55 & 62.92 \end{bmatrix}$

**CHAPTER 11**

11.2. a.  $\boldsymbol{\mu} = [29.87, 13.00]^T$ ,  $[S] = \begin{bmatrix} 4.96 & 0.15 \\ 0.15 & 27.12 \end{bmatrix}$

b.  $N_2(\boldsymbol{\mu}, [\Sigma]); \boldsymbol{\mu} = [-1.90, 5.33]^T$   $[\Sigma] = \begin{bmatrix} 5.23 & 7.01 \\ 7.01 & 50.24 \end{bmatrix}$

11.3.  $r = 0.974 > r_{\text{crit}}(10\%) = 0.970$ ; do not reject

11.4. a.  $T^2 = 68.5 >> 18.421 = \chi^2_2(.9999)$ ; reject

b.  $\mathbf{a} \propto [-.6217, .1929]^T$

11.5. a.  $T^2 = 7.80$ , reject @ 5%

b.  $\mathbf{a} \propto [-.0120, .0429]^T$

**CHAPTER 12**

- 12.1. a. 3.78, 4.51  
 b. 118.8  
 c. 0.979
- 12.2. a. Correlation matrix:  $\Sigma \lambda_k = 3$   
 b. 1, 1, 1  
 c.  $\mathbf{x}_1^T \approx [26.2, 42.6, 1009.6]$
- 12.3. a. [1.51, 6.80], [0.22, 0.98], [0.10, 0.46]  
 b.  $\lambda_2$  and  $\lambda_3$  may be entangled
- 12.4. a.  $\begin{bmatrix} .593 & .332 & .734 \\ .552 & -.831 & -.069 \\ -.587 & -.446 & .676 \end{bmatrix}$   
 b.  $\begin{bmatrix} .377 & .556 & 1.785 \\ .351 & -1.39 & -.168 \\ -.373 & -.747 & 1.644 \end{bmatrix}$
- 12.5. 9.18, 14.34, 10.67

**CHAPTER 13**

- 13.1. Jan 6:  $v_1 = .038$ ,  $w_1 = .433$ ; Jan 7:  $v_1 = .868$ ,  $w_1 = 1.35$
- 13.2. 39.0°F, 23.6°F
- 13.3. a.  $\begin{bmatrix} 1.883 & 0 & 1.838 & -.212 \\ 0 & .927 & .197 & .791 \\ 1.838 & .197 & 1.904 & 0 \\ -.212 & .791 & 0 & .925 \end{bmatrix}$   
 b.  $\mathbf{a}_1 = [.728, .032]^T$ ,  $\mathbf{b}_1 = [.718, -.142]^T$ ,  $r_{C1} = 0.984$   
 $\mathbf{a}_2 = [-.023, 1.038]^T$ ,  $\mathbf{b}_2 = [.099, 1.030]^T$ ,  $r_{C2} = 0.867$

**CHAPTER 14**

- 14.1. b.  $R_1: -1 \leq x \leq 0.25$   
 $R_2: 0.25 < x \leq 1.5$   
 c.  $R_1: -1 \leq x \leq -0.33$   
 $R_2: -0.33 < x \leq 1.5$
- 14.2. a.  $\mathbf{a}_1^T = [0.83, -0.56]$   
 b. 1953  
 c. 1953

- 14.3. a.  $\delta_1 = 38.65$ ,  $\delta_2 = -14.99$ ; Group 3  
 b.  $5.2 \times 10^{-12}$ ,  $2.8 \times 10^{-9}$ , 0.99999997
- 14.4. a. 0.006  
 b. 0.059  
 c. 0.934

## CHAPTER 15

$$15.1. \begin{bmatrix} 0 & & & & & \\ 3.63 & 0 & & & & \\ 2.30 & 1.61 & 0 & & & \\ 3.14 & 0.82 & 0.90 & 0 & & \\ 0.73 & 4.33 & 2.93 & 3.80 & 0 & \\ 1.64 & 2.28 & 0.72 & 1.62 & 2.22 & 0 \end{bmatrix}$$

- 15.2. a. 1967+1970,  $d = 0.72$ ; 1965+1969,  $d = 0.73$ ; 1966+1968,  $d = 0.82$ ;  
 (1967+1970) + (1966+1968),  $d = 1.61$ ; all,  $d = 1.64$ .  
 b. 1967+1970,  $d = 0.72$ ; 1965+1969,  $d = 0.73$ ; 1966+1968,  $d = 0.82$ ;  
 (1967+1970) + (1966+1968),  $d = 2.28$ ; all,  $d = 4.33$ .  
 c. 1967+1970,  $d = 0.72$ ; 1965+1969,  $d = 0.73$ ; 1966+1968,  $d = 0.82$ ;  
 (1967+1970) + (1966+1968),  $d = 1.60$ ; all,  $d = 3.00$ .
- 15.3. a. 1967+1970,  $d = 0.50$ ; 1965+1969,  $d = 0.60$ ; 1966+1968,  $d = 0.70$ ;  
 (1967+1970) + (1965+1969),  $d = 1.25$ ; all,  $d = 1.925$ .  
 b. 1967+1970,  $d = 0.125$ ; 1965+1969,  $d = 0.180$ ; 1966+1968,  $d = .245$ ;  
 (1967+1970) + (1965+1969),  $d = 1.868$ ; all,  $d = 7.053$ .
- 15.4. {1966, 1967}, {1965, 1968, 1969, 1970}; {1966, 1967, 1968}, {1965, 1969, 1970};  
 {1966, 1967, 1968, 1970}, {1965, 1969}.