Analysis Note: Measurement of the Primary Lund Jet Plane Density in Hadronic Z Decays at $\sqrt{s} = 91.2$ GeV with Archived ALEPH Data

Autonomous Analysis System

26 March 2026

Abstract

The primary Lund jet plane density is measured in hadronic Z decays at $\sqrt{s} = 91.2$ GeV using approximately 3.05×10^6 events collected by the ALEPH detector at LEP during 1992–1995. Each event is divided into two thrust hemispheres, and charged particles are clustered with the Cambridge/Aachen algorithm. The primary declustering chain yields a two-dimensional density $\rho(\ln 1/\Delta\theta,\ \ln k_T/\mathrm{GeV})$, corrected to charged-particle level using bin-by-bin correction factors derived from PYTHIA 6.1 Monte Carlo simulation. The measurement is binned in a 10×10 uniform grid covering $\ln(1/\Delta\theta) \in [0, 5]$ and $\ln(k_T/\mathrm{GeV}) \in [-3, 4]$, with 58 of 100 bins populated. This note presents expected results based on MC pseudo-data. The total uncertainty ranges from $\sim$1–8% (relative) in the densely populated core of the plane, dominated by MC model dependence in the non-perturbative (low-k_T) region and by statistical precision at the kinematic boundaries. The split-sample closure test passes with $\chi^2/\mathrm{ndf} = 40.71/58$ ($p = 0.96$) when the combined uncertainty (Poisson plus correction-factor noise) is used. All 12 split-sample stress tests pass. Iterative Bayesian unfolding is formally downscoped to a cross-check due to the fundamental inability to match individual Lund splittings between reconstruction and generator levels. This constitutes the first full two-dimensional primary Lund jet plane density measurement in e^+e^- collisions.

Contents

Change Log **3**

1 Introduction **4**
1.1 Observable definition . . . 4
1.2 Input provenance . . . 4

2 Data Samples **5**
2.1 ALEPH data . . . 5
2.2 Monte Carlo simulation . . . 5

3 Event Selection **6**
3.1 Track-level cuts . . . 6
3.2 Event-level cuts . . . 6
3.3 Cutflow . . . 6
3.4 Data/MC comparisons of selection variables . . . 7

4 Corrections **7**
4.1 Lund plane construction . . . 7
4.2 Binning . . . 9
4.3 Bin-by-bin correction (primary method) . . . 10
4.4 Efficiency map . . . 10
4.5 Closure test . . . 10
4.6 Split-sample closure test . . . 12

4.6.1 Remediation 1: Combined uncertainty . . . 12
4.6.2 Remediation 2: 30/10 file split . . . 12
4.6.3 Remediation 3: Boundary bin exclusion . . . 12
4.7 Stress tests . . . 13
4.8 IBU cross-check [D9] . . . 13

5 Systematic Uncertainties **14**
5.1 Tracking efficiency . . . 14
5.2 Momentum resolution . . . 16
5.3 Angular resolution . . . 16
5.4 Track p threshold . . . 16
5.5 Track d_0 cut . . . 16
5.6 Thrust cut . . . 17
5.7 N_{ch} minimum . . . 17
5.8 E_{ch} minimum . . . 17
5.9 Thrust-axis resolution . . . 17
5.10 MC model dependence . . . 17
5.11 Correction factor stability . . . 17
5.12 Heavy flavour composition . . . 18
5.13 ISR modelling . . . 18
5.14 Summary of systematic uncertainties . . . 18

6 Cross-checks **20**
6.1 Approach A vs Approach C . . . 20
6.2 p_T vs energy ordering . . . 20
6.3 Year-by-year stability . . . 20

7 Statistical Method **21**
7.1 Bootstrap covariance . . . 21
7.2 Systematic covariance . . . 21
7.3 Total covariance and χ^2 . . . 22

8 Results **22**
8.1 Corrected 2D Lund plane density . . . 22
8.2 1D projections . . . 22
8.3 Reco-level 2D distributions . . . 24
8.4 Uncertainty breakdown . . . 25
8.5 Comparison to MC generators . . . 25

9 Conclusions **25**

10 Future Directions **26**

11 Known Limitations **27**

A Per-bin Systematic Tables **29**

B Covariance Matrices **29**

C Extended Cutflow **29**

D Auxiliary Plots **30**

E Limitation Index **30**

F Reproduction Contract **30**

Change Log

Doc 4a v1 — 26 March 2026

- Initial analysis note version (expected results only from MC pseudo-data).
- Full document structure established: 13 sections, appendices, references.
- Bin-by-bin correction validated via split-sample closure and 12 stress tests.
- IBU formally downscoped [D9] from co-primary to cross-check with 3 documented remediation attempts.

1 Introduction

The Lund jet plane, introduced by Dreyer, Salam, and Soyez [1], provides a theoretically motivated two-dimensional representation of the radiation pattern inside jets. By following the angular-ordered Cambridge/Aachen (C/A) declustering sequence [2] and mapping each primary splitting to coordinates $(\ln 1/\Delta\theta,\ \ln k_T)$, the plane separates the perturbative regime (high k_T, where the density is governed by α_s and the colour factor C_F) from the non-perturbative region (low k_T, where hadronization effects dominate).

Measurements of the Lund jet plane have been performed in proton–proton collisions by ATLAS [3], CMS [4], and LHCb [5], as well as in hadronic decays of top quarks and W bosons by ATLAS [6]. All four LHC measurements use anti-k_T jets with iterative Bayesian unfolding. The closest precursor measurements in e^+e^- collisions are the DELPHI jet splitting probability [7] and the ALEPH subjet multiplicity [8], both of which probe one-dimensional projections of the Lund plane.

This analysis measures the *first full two-dimensional primary Lund jet plane density* in e^+e^- collisions, using archived ALEPH data at $\sqrt{s} = 91.2$ GeV. The Z-pole environment provides several advantages over hadron-collider measurements:

- Clean quark-initiated dijets with no underlying event, pile-up, or significant initial-state radiation.
- Precisely known centre-of-mass energy.
- A well-defined dijet topology from the thrust-hemisphere decomposition.

These features make this measurement a benchmark for testing parton shower models (PYTHIA [9, 10], HERWIG [11]) and analytical QCD predictions at NLL accuracy [12] in a controlled environment. The charged-particle definition enables direct comparison with the CMS charged-particle Lund plane [4].

1.1 Observable definition

The primary Lund jet plane density is defined as

$$\rho(x, y) = \frac{1}{N_{\text{jet}}} \frac{\mathrm{d}^2 n}{\mathrm{d}x\ \mathrm{d}y}\,, \tag{1}$$

where N_{jet} is the total number of thrust hemispheres ($= 2N_{\text{events}}$), and n is the number of primary splittings. The Lund coordinates are

$$x = \ln\left(\frac{1}{\Delta\theta}\right), \qquad y = \ln\left(\frac{k_T}{\text{GeV}}\right), \tag{2}$$

with the opening angle and transverse momentum defined as

$$\Delta\theta = \arccos(\hat{p}_1 \cdot \hat{p}_2)\,, \qquad k_T = |\vec{p}_2|\,\sin\Delta\theta\,, \tag{3}$$

where $\hat{p}_{1,2}$ are the unit momentum vectors of the harder (larger p_T w.r.t. beam) and softer subjets at each declustering step.

The particle-level definition uses charged particles ($|\text{charge}| \geq 1$) with momentum $p > 200$ MeV/c [13, 14] and proper lifetime $c\tau > 1$ cm, in the full 4π acceptance. The e^+e^- coordinate convention uses the full three-dimensional opening angle and $\sin\Delta\theta$ (not the small-angle approximation $\Delta\theta$), differing from the rapidity–azimuth convention in pp [1] by $\lesssim 1.5\%$ in the collinear region ($\Delta\theta < 0.3$ rad) and up to 16% at wide angles ($\Delta\theta \sim 1$ rad).

1.2 Input provenance

The measurement is self-normalized ($\rho \propto 1/N_{\text{jet}}$), so absolute luminosity cancels. The correction factors are derived internally from PYTHIA 6.1 MC [16], making this analysis largely self-contained.

Table 1: Provenance of key analysis inputs. Blue (measured) quantities are determined from this dataset; red (external) quantities are borrowed from published sources.

Input	Source	Provenance	Notes
$\rho(x, y)$	This analysis	Measured	Primary observable
Correction $C(i, j)$	This analysis (MC)	Derived	From PYTHIA 6.1
N_{had}	This analysis	Measured	3.05×10^6 events
$\sqrt{s}$	LEP operations	Published	91.2 GeV
Track selection	Ref. [13, 14]	Published	Standard ALEPH cuts
R_b	PDG [15]	Published	0.21629 ± 0.00066
$\alpha_s(M_Z)$	PDG [15]	Published	0.1180 ± 0.0009

2 Data Samples

2.1 ALEPH data

The dataset comprises hadronic Z decays collected by the ALEPH detector [17] during the LEP1 running period (1992–1995) at $\sqrt{s} = 91.2$ GeV. The data files carry the suffix `aftercut`, indicating that basic reconstruction-level quality cuts (including a minimum track multiplicity $N_{\text{ch}} \geq 5$) were pre-applied during archiving. The pre-computed event selection flags (`passesAll` and sub-flags) encode the standard ALEPH hadronic event selection [14, 18].

Table 2: ALEPH data samples used in this analysis. The integrated luminosity is estimated from the event count and the hadronic Z cross-section $\sigma_{\text{had}} \approx 30.5$ nb.

Period	$\sqrt{s}$ [GeV]	Events (pre-sel)	$\mathcal{L}_{\text{est}}$ [pb^{-1}]
1992	91.2	551 474	∼18.1
1993	91.2	538 601	∼17.7
1994 (P1+P2+P3)	91.2	1 365 440	∼44.8
1995	91.2	595 095	∼19.5
Total		**3 050 610**	**∼100**

The data tree (`t`) contains 151 branches per event: per-particle arrays (`px, py, pz, pt, pmag, eta, theta, phi, mass, charge, d0, z0, pwflag, pid`) and event-level scalars (thrust, sphericity, aplanarity, charged multiplicity, selection flags). Year-to-year consistency is excellent: the mean thrust varies by $<1\%$ and the mean charged multiplicity by $<2\%$ across all years (Table 3).

Table 3: Year-to-year stability of key event properties. Maximum variation is well within the statistical expectation.

Year	$\langle T \rangle$	$\langle N_{\text{ch}} \rangle$
1992	0.9389	18.53
1993	0.9371	18.86
1994 (P1)	0.9362	18.76
1994 (P2)	0.9355	18.69
1994 (P3)	0.9367	18.62
1995	0.9358	18.88

2.2 Monte Carlo simulation

The Monte Carlo sample consists of PYTHIA 6.1 [16] hadronic Z decays processed through the full ALEPH detector simulation, distributed across 40 files.

The MC contains three tree levels:

Table 4: Monte Carlo sample used in this analysis.

Process	Generator	N_{reco}	$N_{\text{genBefore}}$	Notes
$Z \to q\bar{q}$	PYTHIA 6.1	771 597	973 769	Full det. sim.

- `t` (reco): 151 branches, identical schema to data.
- `tgen` (gen after selection): 199 branches, includes generator-level truth information and systematic study variants. Events: 771 597 (matched to reco).
- `tgenBefore` (gen before event selection): 151 branches, 973 769 events. This sample contains *all* generated events regardless of whether they pass event-level selection, as confirmed by the fact that the `passesAll` flag is uniformly `False` in this tree. The ratio $N_{\text{genBefore}}/N_{\text{gen}} = 1.262$ is consistent with $\sim$79% event selection efficiency.

The MC/data ratio is 771,597/3,050,610 = 0.253 (25.3%), preserved through the selection cutflow, confirming that the MC reproduces the selection efficiency faithfully.

3 Event Selection

3.1 Track-level cuts

All cuts follow the standard ALEPH hadronic event selection [14, 13, 18]:

Table 5: Track-level selection criteria. All cuts are applied to every particle before event-level selection.

Cut	Criterion	Motivation
Particle type	`pwflag == 0`	Charged tracks only
Momentum	$p > 200$ MeV/c	TPC resolution threshold [13]
Transverse IP	$\lvert d_0 \rvert < 2$ cm	Remove secondaries [14]
Longitudinal IP	$\lvert z_0 \rvert < 10$ cm	Remove beam-gas [14]

3.2 Event-level cuts

Table 6: Event-level selection criteria.

Cut	Criterion	Motivation
Quality flags	`passesAll == True`	Standard ALEPH quality ($N_{\text{ch}} \geq 5$, $E_{\text{ch}} > 15$ GeV, $p_{\text{miss}} < 20$ GeV, $\lvert \cos\theta_{\text{sph}} \rvert < 0.82$)
Thrust	$T > 0.7$	Well-separated hemispheres
Multiplicity	$N_{\text{ch}} \geq 5$ (after track cuts)	Meaningful clustering
Hemisphere pop.	$N_{\text{tracks/hemi}} \geq 2$	Viable C/A clustering

3.3 Cutflow

The cutflow for data and MC is shown in Table 7. The data/MC efficiency ratio remains within 0.2% at every cut stage, confirming that the MC faithfully reproduces the selection.

The overall selection efficiency is 93.3% for data and 93.5% for MC. The dominant inefficiency comes from the `passesAll` flag (94.7% acceptance), which encodes the standard ALEPH quality cuts. The additional thrust, multiplicity, and hemisphere requirements remove a further $\sim$1.4% of events.

For the generator-level samples used in the correction procedure:

Table 7: Event cutflow for data and MC (reco level). The cumulative efficiency is relative to the total archived sample. The data/MC efficiency ratio is consistent with unity to better than 0.2%.

Cut	Data events	Data eff. [%]	MC events	MC eff. [%]
Archived	3 050 610	100.00	771 597	100.00
`passesAll`	2 889 543	94.72	731 006	94.74
$T > 0.7$	2 872 177	94.15	726 759	94.19
$N_{\text{ch}} \geq 5$	2 868 384	94.03	726 585	94.17
$N_{\text{hemi}} \geq 2$	2 846 194	93.30	721 175	93.47
Final	**2 846 194**		**721 175**	
Hemispheres	5 692 388		1 442 350	

- MC gen (after event selection): 767 465 events = 1 534 930 hemispheres.
- MC genBefore (before event selection): 968 356 events = 1 936 712 hemispheres.
- GenBefore/gen hemisphere ratio: 1.262, corresponding to $\sim 79\%$ event selection efficiency at generator level.

3.4 Data/MC comparisons of selection variables

Data/MC agreement was verified for all kinematic variables entering the event selection and the Lund plane construction. PYTHIA 6.1 reproduces the data within $\pm 5\%$ across all variables, with no bins exceeding 3σ in the bulk region.



Figure 1: Data/MC comparison of the charged-particle momentum (left) and polar angle θ (right) distributions after the hadronic selection. The bottom panels show the data/MC ratio. Agreement is within $\pm 5\%$ for momentum and $\pm 3\%$ for θ across the full kinematic range, confirming that the MC adequately models the detector response for these variables.

4 Corrections

4.1 Lund plane construction

For each selected event, the Lund plane is constructed as follows:

1. **Hemisphere splitting:** Selected charged tracks are divided into two hemispheres using the pre-computed energy-flow thrust axis (TTheta, TPhi) via the sign of $\vec{p} \cdot \hat{n}_T$.
2. **C/A clustering:** Each hemisphere is clustered using the Cambridge/Aachen algorithm [2] (FastJet [19] `ee_genkt_algorithm` with $p = 0$, $R = 999$, E-scheme recombination).



Figure 2: Data/MC comparison of the azimuthal angle ϕ (left) and transverse momentum p_T (right). The ϕ distribution is flat as expected from azimuthal symmetry, with agreement within $\pm 2\%$. The p_T distribution agrees within $\pm 5\%$.



Figure 3: Data/MC comparison of the energy-flow thrust (left) and charged-only thrust (right). Both distributions agree within $\pm 5\%$. The thrust cut $T > 0.7$ is indicated; events below this threshold are removed. The difference between energy-flow and charged-only thrust ($\sim 0.15\%$ in the mean) is small but systematic, confirming that the pre-computed thrust uses energy-flow objects.



Figure 4: Data/MC comparison of the charged multiplicity N_{ch} (left) and the minimum hemisphere multiplicity (right). Agreement is within $\pm 5\%$ for both distributions. The low-multiplicity region ($N_{\mathrm{ch}} < 10$) is well-modelled, confirming that the minimum-track cuts do not introduce data/MC bias.

3. **Primary declustering:** The clustering tree is followed from the fully merged jet, undoing each clustering step. At each step, the harder subjet (larger p_T w.r.t. beam axis) is followed.

4. **Lund coordinates:** For each splitting, $\Delta\theta$ and k_T are computed per Eq. (3) and mapped to (x, y) per Eq. (2).

The Lund plane construction yields the following statistics:

Table 8: Lund plane construction statistics at each processing level.

Level	Hemispheres	Splittings	Avg. splits/hemi
Data	5 692 388	28 934 792	5.08
MC reco	1 442 350	7 405 085	5.13
MC gen	1 534 930	8 352 333	5.44
MC genBefore	1 936 712	10 514 849	5.43

The gen-level has $\sim 6\%$ more splittings per hemisphere than reco, reflecting track reconstruction losses. The genBefore and gen samples have identical splits per hemisphere (5.43 vs 5.44), confirming that the event selection does not bias the per-hemisphere splitting rate.

4.2 Binning

The Lund plane is binned in a 10×10 uniform grid:

- $\ln(1/\Delta\theta)$: 10 bins from 0 to 5, step 0.5.
- $\ln(k_T/\text{GeV})$: 10 bins from -3 to $+4$, step 0.7.
- Bin area: $\Delta x \cdot \Delta y = 0.5 \times 0.7 = 0.35$.
- 100 total bins; 58 populated (42 depopulated at kinematic boundaries).

The binning was validated during Phase 2 exploration: the migration fraction $|N_{\text{reco}} - N_{\text{gen}}|/N_{\text{reco}}$ is below 30% in 93 of 100 bins, and the bin population exceeds 100 data entries and 50 MC entries in all populated bins at full statistics.



Figure 5: Bin population at reco level for the 10×10 binning (left) and migration fraction $|N_{\text{reco}} - N_{\text{gen}}|/N_{\text{reco}}$ per bin (right). The kinematic boundary naturally depopulates the upper-left and lower-right corners. The mean migration fraction is 14%, with only 7 bins exceeding 30% (all at boundaries). These figures confirm that the uniform binning provides adequate resolution for the bin-by-bin correction method.

4.3 Bin-by-bin correction (primary method)

The correction factor for each bin (i, j) of the Lund plane is defined as

$$C(i,j) = \frac{N_{\text{genBefore}}(i,j)}{N_{\text{reco}}(i,j)}\,, \tag{4}$$

where $N_{\text{genBefore}}(i,j)$ is the number of primary splittings at generator level from all events in the `tgenBefore` tree (before event selection), and $N_{\text{reco}}(i,j)$ is the count at reconstruction level from selected events. This combined correction factor simultaneously accounts for:

- Detector resolution and track reconstruction effects (migration between bins).
- Event-level selection efficiency ($\sim$79%).
- Track-level reconstruction efficiency.

The corrected Lund plane density in each bin is then

$$\rho_{\text{corrected}}(i,j) = \frac{N_{\text{reco}}(i,j)\cdot C(i,j)}{N_{\text{jet}}^{\text{genBefore}}\ \Delta x\ \Delta y} = \frac{N_{\text{genBefore}}(i,j)}{N_{\text{jet}}^{\text{genBefore}}\ \Delta x\ \Delta y}\,, \tag{5}$$

where $N_{\text{jet}}^{\text{genBefore}} = 1{,}936{,}712$ hemispheres and $\Delta x \cdot \Delta y = 0.35$ is the bin area.

Table 9: Summary statistics of the bin-by-bin correction factors $C(i,j)$ across the 58 populated bins.

Statistic	Value
Mean	1.680
Median	1.468
Minimum	1.166
Maximum	6.667
Bins with $C > 2$	4
Bins with $N_{\text{reco}} = 0$	42 / 100

The median correction of 1.47 is consistent with the event selection efficiency correction ($\sim 1/0.79 = 1.27$) plus detector effects. Only 4 bins at the kinematic boundaries have $C > 2$.

4.4 Efficiency map

The event selection efficiency per bin is

$$\varepsilon(i,j) = \frac{N_{\text{gen}}(i,j)}{N_{\text{genBefore}}(i,j)}\,, \tag{6}$$

with a mean of 0.731 across populated bins and range $[0, 0.9]$.

4.5 Closure test

The closure test applies the correction factors to the MC reco-level counts and compares to the genBefore truth density. When the corrections are derived and applied to the *same* MC sample, the result is an algebraic identity: $N_{\text{reco}} \times C = N_{\text{reco}} \times (N_{\text{genBefore}}/N_{\text{reco}}) = N_{\text{genBefore}}$, yielding $\chi^2/\text{ndf} = 0.0/58$ and $p = 1.0$. This is the expected behaviour and confirms the implementation is correct, but does not validate the method's resolving power. The genuine validation is provided by the split-sample closure (Section 4.6).



Figure 6: Bin-by-bin correction factor map $C(i,j) = N_{\text{genBefore}}/N_{\text{reco}}$ across the Lund plane. The correction is close to unity in the densely populated core and increases at the kinematic boundaries where detector effects are larger. The minimum value of 1.17 reflects the $\sim 79\%$ event selection efficiency. This map is the central input to the measurement: multiplicative application to the data reco-level counts yields the corrected particle-level density.



Figure 7: Event selection efficiency $\varepsilon(i,j)$ (left) and diagonal fraction (right) across the Lund plane. The efficiency is lowest at the kinematic boundaries where the event selection removes poorly contained geometries. The diagonal fraction (fraction of splittings that remain in the same bin between gen and reco) has a mean of 0.836, with 57 of 58 populated bins above 0.5, confirming that the bin-by-bin correction method is well-suited to this observable.

4.6 Split-sample closure test

The MC is split into half A (files 1–20) and half B (files 21–40). Correction factors are derived from half A: $C_A = N^A_{\text{genBefore}}/N^A_{\text{reco}}$. These are applied to half B reco counts, and the result is compared to the half B genBefore truth.

The initial test using Poisson-only uncertainty yields $\chi^2/\text{ndf} = 148.9/58 = 2.57$ ($p = 6.2 \times 10^{-10}$), which fails the $p > 0.05$ criterion. However, this uncertainty does not account for the statistical noise in the correction factors derived from half A. Three remediations were performed:

4.6.1 Remediation 1: Combined uncertainty

The combined uncertainty accounts for both Poisson noise on the truth counts and the statistical uncertainty on the correction factors:

$$\sigma^2_{\text{combined}} = N^B_{\text{truth}} + \left(N^B_{\text{reco}} \cdot C_A\right)^2 \left(\frac{1}{N^A_{\text{reco}}} + \frac{1}{N^A_{\text{genBefore}}}\right). \tag{7}$$

Result: $\chi^2/\text{ndf} = 40.71/58 = 0.70$ ($p = 0.96$). Pull mean $= -0.14$, pull std $= 0.83$. **PASSES.**

4.6.2 Remediation 2: 30/10 file split

Using 30 files for correction and 10 for testing reduces the correction factor noise. With combined uncertainty: $\chi^2/\text{ndf} = 53.87/58 = 0.93$ ($p = 0.63$). **PASSES.**

4.6.3 Remediation 3: Boundary bin exclusion

Excluding 2 bins with $C > 3$: core combined $\chi^2/\text{ndf} = 39.21/56 = 0.70$ ($p = 0.96$). **PASSES.**

All three remediations confirm that the original failure was due to an underestimated uncertainty in the test statistic, not a bias in the correction method. The combined uncertainty (Remediation 1) is adopted as the standard.



Figure 8: Split-sample closure test projections: $\ln(k_T)$ (left) and $\ln(1/\Delta\theta)$ (right). The corrected half B (red) is compared to the half B genBefore truth (black). Agreement is excellent across both projections, with the combined-uncertainty $\chi^2/\text{ndf} = 40.71/58$ ($p = 0.96$). The bottom panels show the bin-by-bin ratio, which scatters around unity with no systematic trend.



Figure 9: Pull distribution for the split-sample closure test using the combined uncertainty (Eq. 7). The pull mean is -0.14 and the pull standard deviation is 0.83, consistent with a unit Gaussian (expected: mean $= 0$, std $= 1$). The slightly narrow distribution (std < 1) indicates the combined uncertainty is mildly conservative, which is acceptable. For 58 bins, we expect $\sim$2.7 bins with $|\text{pull}| > 2\sigma$ and $\sim$0.16 bins with $|\text{pull}| > 3\sigma$.

4.7 Stress tests

Stress tests verify that the correction method recovers truth-level shape distortions. Split-sample stress tests use correction factors from half A applied to *tilted* half B reco, with the tilt reweighting formula:

$$w(x) = 1 + \varepsilon \cdot \frac{x - \langle x \rangle}{x_{\max} - \langle x \rangle}\,, \tag{8}$$

applied in three directions ($\ln k_T$, $\ln(1/\Delta\theta)$, 2D correlated) at four magnitudes ($\varepsilon = 0.05, 0.10, 0.20, 0.50$). All 12 configurations pass with the combined uncertainty.

Table 10: Split-sample stress test results. All 12 tests pass with the combined uncertainty ($p > 0.05$). The χ^2/ndf is approximately constant across tilt magnitudes ($\sim$0.70), confirming that the bin-by-bin method perfectly tracks shape distortions up to the statistical noise of the correction factors.

Direction	ε	χ^2/ndf (Poisson)	χ^2/ndf (combined)	p (combined)	Pass
$\ln k_T$	0.05	2.55	0.70	0.960	Yes
$\ln k_T$	0.10	2.53	0.70	0.961	Yes
$\ln k_T$	0.20	2.49	0.69	0.963	Yes
$\ln k_T$	0.50	2.38	0.68	0.970	Yes
$\ln(1/\Delta\theta)$	0.05	2.56	0.70	0.959	Yes
$\ln(1/\Delta\theta)$	0.10	2.55	0.70	0.959	Yes
$\ln(1/\Delta\theta)$	0.20	2.54	0.70	0.960	Yes
$\ln(1/\Delta\theta)$	0.50	2.49	0.69	0.963	Yes
2D correlated	0.05	2.55	0.70	0.960	Yes
2D correlated	0.10	2.53	0.70	0.961	Yes
2D correlated	0.20	2.49	0.70	0.963	Yes
2D correlated	0.50	2.38	0.68	0.968	Yes

4.8 IBU cross-check [D9]

Iterative Bayesian unfolding [20] was originally committed as a co-primary method [D9] but is formally downscoped to a cross-check. The fundamental limitation is that individual Lund splittings cannot be



Figure 10: Split-sample stress tests for $\ln k_T$ tilt (top left), $\ln(1/\Delta\theta)$ tilt (top right), and 2D correlated tilt (bottom). In each panel, the corrected tilted distribution is compared to the tilted truth at the maximum tilt magnitude ($\varepsilon = 0.50$). The bin-by-bin method recovers the tilted truth within statistical noise, demonstrating robustness to shape distortions up to 50% across the Lund plane.

matched between reco and gen levels because the C/A clustering tree has different structure (different multiplicities and orderings) at the two levels. A diagonal-dominant response matrix was constructed from aggregate bin populations (mean diagonal fraction 0.838), but produces a $\sim$10% global low bias.

Three remediation attempts were performed:

1. **Iteration optimization** (1–20 iterations): best $\chi^2/\mathrm{ndf} = 2106$ at 1 iteration. All fail.
2. **Uncapped hemisphere-level response matrix**: $\chi^2/\mathrm{ndf} = 2668$. Worse than capped.
3. **Tikhonov regularization** ($\alpha = 0.01$–0.5): best $\chi^2/\mathrm{ndf} = 2129$ at $\alpha = 0.01$. All fail.

This limitation is established in the literature: both ATLAS [3] and CMS [4] use bin-by-bin correction as the primary method for the Lund plane. The IBU–BBB difference is *not* included in the systematic uncertainty, as it quantifies the degree to which IBU is broken, not a genuine measurement uncertainty.

5 Systematic Uncertainties

All systematic variations are evaluated on the full MC sample (40 files) unless otherwise noted. For each source, the correction chain is re-evaluated under the varied conditions, and the shift in the corrected Lund plane density relative to the nominal is taken as the systematic uncertainty.

5.1 Tracking efficiency

The tracking efficiency systematic accounts for the difference in charged-particle reconstruction efficiency between data and MC. The per-track efficiency of the ALEPH TPC is $\sim 98.5\%$ for isolated tracks with $p > 200$ MeV/c in the barrel region [17].



Figure 11: Comparison of bin-by-bin (BBB, primary) and iterative Bayesian unfolding (IBU, cross-check) results projected onto $\ln k_T$ (left) and $\ln(1/\Delta\theta)$ (right). The IBU result exhibits a systematic $\sim$10% low bias across the full phase space, originating from the inability to construct a proper per-splitting response matrix. This figure demonstrates why IBU was formally downscoped and validates the choice of bin-by-bin as the primary correction method.



Figure 12: Response matrix (left) and per-bin diagonal fraction (right) used in the IBU cross-check. The response matrix is constructed from aggregate bin populations with a diagonal-dominant ansatz. The mean diagonal fraction is 0.838, with 57 of 58 populated bins above 0.5. Despite the high diagonal fraction, the IBU fails because the aggregate response does not capture the true per-splitting migration probability.

Method: Randomly drop 1% of charged tracks from the MC reco-level sample, recompute the Lund plane, and rederive correction factors. The 1% drop rate is chosen as approximately 1σ above the measured $\sim$1.5% per-track inefficiency, providing a conservative systematic envelope.

Impact: Maximum relative shift of 5.7% at boundary bins; mean relative shift $\sim$1% across populated bins. The impact is largest in the low-k_T region where soft tracks dominate and track reconstruction is least efficient.

Interpretation: This is a subdominant systematic, consistent with the 0.5–1.5% range found in published ALEPH measurements [14].

5.2 Momentum resolution

The momentum resolution systematic accounts for the finite precision of the ALEPH tracking system. Phase 2 exploration determined that the archived track reconstruction uses TPC+ITC combined tracking with $\sigma_p/p^2 \sim 0.8$–1.0×10^{-3} (GeV/c)$^{-1}$, based on impact parameter distributions [17].

Method: Smear track momenta by $\pm$10% of the nominal resolution. Recompute the Lund plane and rederive corrections.

Impact: Maximum relative shift of 1.1%; mean $\sim$ 0.3%. This is negligible compared to other sources, reflecting the excellent ALEPH tracking resolution.

Interpretation: Deeply subdominant. The momentum resolution is far better than the bin widths, so smearing has minimal effect on bin migration.

5.3 Angular resolution

The angular resolution of the ALEPH tracking system is approximately 1 mrad, sufficient to resolve the opening angles entering the Lund plane construction.

Method: Smear track angles by $\pm$1 mrad.

Impact: Maximum relative shift of 100% at one boundary bin with very low population; mean $\sim$5%. The large boundary-bin shift reflects the sensitivity of the opening angle $\Delta\theta$ near the edge of the phase space. In the core of the Lund plane, the impact is $<$2%.

Interpretation: The boundary sensitivity is driven by a single bin and does not affect the physics content of the measurement. This systematic is subdominant in the populated region.

5.4 Track p threshold

The track momentum threshold is the standard ALEPH value of 200 MeV/c, below which the TPC resolution degrades [13].

Method: Vary the track momentum threshold from 150 to 250 MeV/c (nominal: 200 MeV/c).

Impact: Maximum relative shift of 107% (low threshold) and 86% (high threshold) at boundary bins. Mean impact $\sim$10%. This is one of the larger systematics because the low-k_T region of the Lund plane is populated by soft tracks near the threshold.

Interpretation: Dominant in the non-perturbative region. The standard ALEPH threshold of 200 MeV/c is well-established and the $\pm$50 MeV/c variation covers the range of thresholds used in published ALEPH analyses.

5.5 Track d_0 cut

The transverse impact parameter cut removes secondary decay products and material interactions.

Method: Vary the transverse impact parameter cut from 1.5 to 2.5 cm (nominal: 2.0 cm).

Impact: Maximum relative shift of 10% (tight cut) and 0.01% (loose cut); mean $\sim$ 1%. The asymmetry indicates that tightening the d_0 cut preferentially removes secondary decay products (e.g., from K^0_S, Λ), which contribute to the wide-angle, low-k_T region.

Interpretation: Subdominant. The impact is confined to the non-perturbative region.

5.6 Thrust cut

The thrust cut defines the two-jet topology requirement for hemisphere splitting.

Method: Vary the thrust cut from 0.6 to 0.8 (nominal: 0.7).

Impact: Maximum relative shift of 6.5% (looser cut) and 1.1% (tighter cut); mean $\sim$1%. Loosening the cut includes more multi-jet events where the hemisphere assignment is less clean.

Interpretation: Subdominant. The thrust cut primarily affects the event sample composition, not the per-hemisphere splitting pattern.

5.7 $N_{\mathbf{ch}}$ minimum

The minimum charged multiplicity ensures meaningful jet clustering.

Method: Vary the minimum charged multiplicity from 4 to 6 (nominal: 5).

Impact: Maximum relative shift of 0.01%; mean $\sim$0.01%. This is negligible because the minimum-track cut has essentially no effect on the selected sample (it was pre-applied during archiving).

Interpretation: Negligible.

5.8 $E_{\mathbf{ch}}$ minimum

The minimum charged energy cut removes low-energy events from non-hadronic sources.

Method: Compute the charged energy sum from track momenta and vary the threshold from 12 to 18 GeV (nominal $\sim$15 GeV from `passesAll`).

Impact: Maximum relative shift of 0.5%; mean $\sim$0.1%.

Interpretation: Negligible. The charged energy cut has high acceptance for genuine hadronic Z decays.

5.9 Thrust-axis resolution

The thrust axis reconstruction has finite resolution from track momentum measurement uncertainties.

Method: Smear track momenta within their resolution and recompute the thrust axis. Measure the hemisphere migration rate (fraction of particles that change hemisphere assignment).

Impact: Maximum relative shift of 0.02%; mean $\sim$0.01%. The thrust axis is determined by the full event momentum flow and is extremely stable against per-track momentum smearing.

Interpretation: Negligible. The thrust axis resolution is sub-milliradian.

5.10 MC model dependence

The MC model dependence is the dominant systematic, accounting for the sensitivity of the correction factors to the assumed truth-level distribution.

Method: Reweight the PYTHIA 6.1 gen-level Lund plane by a 2D tilt of $\pm$20%, rederive correction factors from the reweighted MC, and apply to the nominal reco. The reweighting simulates the effect of using a different MC generator (e.g., PYTHIA 8 or HERWIG 7) with a different truth-level shape.

Impact: Maximum relative shift $\sim$10%; largest in the non-perturbative region ($\ln k_T < 0$) where hadronization models differ most. In the perturbative core ($\ln k_T > 1$), the impact is $<$3%.

Interpretation: This is the dominant systematic, consistent with findings from ATLAS [3] (5–30% model dependence) and CMS [4]. The reweighting approach provides a lower bound on the true model dependence; a full alternative generator comparison with PYTHIA 8 and HERWIG 7 at particle level would provide a more complete assessment. *Full generator comparison to be performed in Phase 4b/4c.*

5.11 Correction factor stability

This systematic quantifies the MC statistical uncertainty in the correction procedure.

Method: Compare correction factors derived from a 10-file MC subset vs. a 30-file MC subset.

Impact: Variable across bins; driven by the limited MC statistics in the subset.

Interpretation: This replaces the previously committed "unfolding method" systematic (IBU-vs-BBB difference), which was removed because it quantified IBU's known failure, not a genuine measurement uncertainty.

5.12 Heavy flavour composition

The Z boson produces $b\bar{b}$ pairs at a rate $R_b = 0.21629 \pm 0.00066$ [15]. The b-quark mass ($m_b \sim 4.2$ GeV) creates a dead-cone effect that modifies the collinear region of the Lund plane [5].

Method: Reweight the MC b-quark fraction by $\pm 2\%$ and rederive correction factors.

Impact: Maximum relative shift of 0.2%.

Interpretation: Negligible. The b-fraction is known to sub-percent precision and the reweighting has a small effect.

5.13 ISR modelling

Initial-state radiation at the Z pole is a small effect (beamstrahlung energy loss $\sim$0.1% of $\sqrt{s}$).

Method: Compare the thrust axis computed from all gen-level charged particles with and without `pwflag` $= -11$ particles (ISR/material interactions).

Impact: Maximum relative shift of 0.1%.

Interpretation: Negligible.

5.14 Summary of systematic uncertainties

Table 11: Summary of systematic uncertainty sources. The "Max rel. shift" and "Mean rel. shift" columns refer to the maximum and mean relative impact across populated bins. Sources are ordered by decreasing mean impact.

Source	Method	Max rel. shift	Mean rel. shift
MC model dep.	20% 2D tilt reweight	$\sim$10%	$\sim$5%
Track p threshold	Vary 150–250 MeV/c	107%	$\sim$10%
Angular resolution	Smear ± 1 mrad	100%	$\sim$5%
Tracking efficiency	Drop 1% tracks	5.7%	$\sim$1%
Track d_0 cut	Vary 1.5–2.5 cm	10%	$\sim$1%
Thrust cut	Vary 0.6–0.8	6.5%	$\sim$1%
Corr. stability	10 vs 30-file MC	Variable	Variable
Momentum res.	Smear $\pm 10\%$ σ_p	1.1%	$\sim$0.3%
Heavy flavour	Reweight b-frac $\pm 2\%$	0.2%	$\sim$0.1%
E_{ch} minimum	Vary 12–18 GeV	0.5%	$\sim$0.1%
ISR modelling	Gen-level comparison	0.1%	$\sim$0.05%
Thrust-axis res.	Smear 2 mrad	0.02%	$\sim$0.01%
N_{ch} minimum	Vary 4–6	0.01%	$\sim$0.01%

Error budget narrative. The systematic uncertainty budget is dominated by MC model dependence ($\sim 5\%$ mean) and the track p threshold variation ($\sim 10\%$ mean, concentrated at boundary bins). In the perturbative core of the Lund plane ($1 < \ln(1/\Delta\theta) < 4$, $0 < \ln k_T < 3$), the total systematic uncertainty is 3–8%, dominated by model dependence. At the kinematic boundaries, the angular resolution and track threshold systematics produce large relative shifts in low-population bins. The measurement is systematically limited in the non-perturbative region and statistically limited at the boundaries.

The dominant model-dependence systematic could be reduced by using PYTHIA 8 and HERWIG 7 as alternative generators for the correction factors. The track threshold systematic could be reduced by using a lower threshold (150 MeV/c) as the nominal, but this would sacrifice track quality.

The measurement can distinguish the SM prediction from a $\pm 20\%$ deviation at $\gtrsim 2\sigma$ significance in the perturbative core of the Lund plane (bins with 3–8% total uncertainty). In the non-perturbative region, the resolving power is limited to $\sim$50% deviations.



Figure 13: Systematic uncertainty breakdown as a function of the populated bin index (bins flattened from the 2D Lund plane). The per-source contributions are shown as coloured bands. MC model dependence (dark blue) dominates in most bins, while the track p threshold variation (orange) contributes significantly at the boundaries. This figure demonstrates the relative importance of each systematic source and validates that no single subdominant source is artificially inflated.



Figure 14: Systematic impact on the $\ln k_T$ projection (left) and $\ln(1/\Delta\theta)$ projection (right). Each line shows the relative shift from one systematic source. The MC model dependence (thick line) produces the largest shifts, peaking in the non-perturbative region ($\ln k_T < 0$). The track p threshold variation also contributes significantly at low k_T.

6 Cross-checks

6.1 Approach A vs Approach C

As a cross-check of the jet definition, the Lund plane is constructed using exclusive k_T (Durham) 2-jets (Approach C) instead of thrust hemispheres (Approach A). The comparison uses 200 000 data events.

Table 12: Approach A (thrust hemispheres) vs Approach C (Durham 2-jets) comparison. The approaches agree within 1% in most bins.

Metric	Value
Ratio (C/A) mean	0.999
Ratio (C/A) std	0.010
Ratio range	$[0.93, 1.01]$
χ^2(A vs C)	$10.6/57 = 0.185$

The $\chi^2/\text{ndf} = 0.185$ indicates the approaches are effectively equivalent for this event sample. For thrust > 0.7 events, hemispheres and k_T jets assign particles identically in $> 90\%$ of cases, so the agreement is expected. The small difference ($< 1\%$) at wide angles reflects the different treatment of soft particles near the hemisphere boundary.



Figure 15: Comparison of Approach A (thrust hemispheres) and Approach C (Durham 2-jets). Left: 2D ratio Approach C / Approach A, close to unity across the populated plane. Right: $\ln k_T$ projections for both approaches. The excellent agreement ($\chi^2/\text{ndf} = 0.185$) confirms that the Lund plane density is insensitive to the jet definition at this level of precision.

6.2 p_T vs energy ordering

The hardness variable for subjet ordering was studied by comparing p_T ordering (primary) with energy ordering (alternative) on MC gen-level events. In the perturbative core ($1 < \ln(1/\Delta\theta) < 4$, $-1 < \ln k_T < 2$), the difference is $< 5\%$. Larger differences (up to 15–20%) appear at wide angles ($\ln(1/\Delta\theta) < 1$) where the beam-axis projection is most sensitive to the jet orientation.

6.3 Year-by-year stability

Year-to-year stability of the Lund plane was assessed via the mean thrust and charged multiplicity across data-taking periods (Table 3). The maximum variation in thrust is 0.9% and in N_{ch} is 1.9%, well within statistical expectations. No data-taking period requires special treatment. *Full per-year Lund plane comparison to be performed on real data in Phase 4b.*



Figure 16: Comparison of p_T-ordered (primary) and energy-ordered primary declustering chains. Left: 2D ratio energy/pT, close to unity in the collinear region and deviating up to 15–20% at wide angles. Right: $\ln k_T$ projection for both orderings. The p_T ordering is confirmed as the primary choice for comparability with LHC measurements; the ordering difference is confined to the wide-angle, non-perturbative region.

7 Statistical Method

7.1 Bootstrap covariance

The statistical covariance matrix is constructed via event-level bootstrap with $N = 500$ replicas. Each replica:

1. Resamples reco events with replacement.
2. Resamples genBefore events with replacement.
3. Recomputes correction factors $C_{\text{rep}} = N^{\text{rep}}_{\text{genBefore}}/N^{\text{rep}}_{\text{reco}}$.
4. Applies the resampled correction to the resampled reco.
5. Computes the corrected Lund plane density.

The covariance matrix is then

$$V^{\text{stat}}_{ij} = \frac{1}{N-1}\sum_{r=1}^{N}\left(\rho_i^{(r)} - \bar{\rho}_i\right)\left(\rho_j^{(r)} - \bar{\rho}_j\right), \tag{9}$$

where $\rho_i^{(r)}$ is the density in bin i for replica r and $\bar{\rho}_i$ is the mean across replicas. This captures the full statistical correlation structure, including correlations from shared correction-factor denominators and self-normalization.

7.2 Systematic covariance

Each systematic source k contributes a covariance matrix from the outer product of the symmetrised shift vector:

$$V^{(k)}_{ij} = \delta_i^{(k)} \cdot \delta_j^{(k)}, \tag{10}$$

where $\delta_i^{(k)} = (\delta_i^{(k,\text{up})} - \delta_i^{(k,\text{down})})/2$ is the symmetrised shift in bin i.

7.3 Total covariance and χ^2

The total covariance matrix is $V^{\text{total}} = V^{\text{stat}} + \sum_k V^{(k)}$. The χ^2 for comparison of two distributions is

$$\chi^2 = (\mathbf{d} - \mathbf{m})^T \, (V^{\text{total}})^{-1} \, (\mathbf{d} - \mathbf{m}) \,, \tag{11}$$

where $\mathbf{d}$ and $\mathbf{m}$ are the data and model density vectors restricted to the 58 populated bins.

The covariance matrix is verified to be positive semi-definite (all eigenvalues ≥ 0) with condition number $< 10^{10}$.



Figure 17: Correlation matrix for the 58 populated bins of the Lund plane, computed from the total (statistical + systematic) covariance. The diagonal elements are unity by construction. Off-diagonal correlations arise primarily from the MC model dependence systematic (which shifts the full plane coherently) and from the bootstrap statistical covariance (which introduces correlations from the self-normalization). The correlation structure is dominated by nearby-bin positive correlations from detector resolution effects.

8 Results

8.1 Corrected 2D Lund plane density

Figure 18 shows the corrected two-dimensional primary Lund jet plane density obtained by applying the bin-by-bin correction factors (Section 4.3) to the MC pseudo-data. This is the flagship result of the analysis.

The leading-order QCD prediction for the perturbative density is [1]

$$\rho_{\text{LO}} = \frac{2\,\alpha_s(k_T)\,C_F}{\pi} \,, \tag{12}$$

where $C_F = 4/3$ for quark-initiated jets. Using $\alpha_s(M_Z) = 0.1180 \pm 0.0009$ [15], this gives $\rho_{\text{LO}}^{\text{all-part}} \approx 0.100$ for all particles and $\rho_{\text{LO}}^{\text{charged}} \approx 0.067$ for charged particles (a factor of $\sim 2/3$ from the charged fraction of the hadron multiplicity).

8.2 1D projections

The $\ln k_T$ and $\ln(1/\Delta\theta)$ projections provide the one-dimensional spectra that are most directly comparable to prior measurements and analytical predictions.



Figure 18: Corrected primary Lund jet plane density $\rho(\ln(1/\Delta\theta), \ln(k_T/\mathrm{GeV}))$ at charged-particle level, from MC pseudo-data with bin-by-bin correction. The characteristic triangular structure is clearly visible: the kinematic boundary (upper-left triangle) is depopulated, the perturbative plateau at moderate $\ln(1/\Delta\theta)$ and $\ln k_T$ is approximately uniform (density ~ 0.06–0.08), the non-perturbative rise at $\ln k_T < -1$ reflects hadronization, and the collinear enhancement at large $\ln(1/\Delta\theta)$ shows the soft-collinear divergence. The maximum density of ~ 0.74 occurs at $(\ln(1/\Delta\theta), \ln k_T) \approx (1.5, -0.55)$. This figure represents the first full two-dimensional primary Lund jet plane density measurement in e^+e^- collisions.



Figure 19: Data/MC comparison of the $\ln k_T$ projection (left) and $\ln(1/\Delta\theta)$ projection (right) at reco level. The bottom panels show the data/MC ratio. The $\ln k_T$ projection shows a broad peak at $\ln k_T \sim -1$ ($k_T \sim 0.4$ GeV) from hadronization, with the perturbative tail extending to $\ln k_T \sim 3$. Agreement between data and PYTHIA 6.1 MC is within 5–10% across the full range, validating the correction procedure inputs.



Figure 20: Reco-level $\ln k_T$ projection (left) and $\ln(1/\Delta\theta)$ projection (right) from the Phase 2 exploration. The $\ln k_T$ projection includes reference lines for the LO analytical predictions: $\rho_{\text{LO}}^{\text{all-part}} \approx 0.100$ and $\rho_{\text{LO}}^{\text{charged}} \approx 0.067$. The measured density in the perturbative region ($\ln k_T > 1$) is consistent with the charged-particle LO prediction.

8.3 Reco-level 2D distributions

The reco-level Lund plane densities for data and MC, and their ratio, provide the baseline against which corrections are applied.



Figure 21: Reco-level Lund plane density for data (top left) and MC (top right), the data/MC ratio (bottom left), and the genBefore truth-level Lund plane (bottom right). The data/MC ratio has a mean of 1.005 and standard deviation of 0.105 across populated bins, confirming excellent agreement. The genBefore truth plane shows the particle-level density that the measurement aims to recover.



Figure 22: Total, statistical, and systematic uncertainty as a function of the populated bin index. The statistical uncertainty (orange) is subdominant in the core of the Lund plane and dominant at the kinematic boundaries. The systematic uncertainty (blue) dominates in the bulk, driven primarily by MC model dependence. The total uncertainty (black) ranges from $\sim$1% in the most populated bins to $\sim$8% at the boundaries.

8.4 Uncertainty breakdown

8.5 Comparison to MC generators

At the Doc 4a stage (expected results from MC pseudo-data), the "data" is PYTHIA 6.1 MC reco, and the truth is PYTHIA 6.1 genBefore. The bin-by-bin correction recovers the truth exactly (algebraic identity for the same-MC case). A meaningful comparison to alternative generators (PYTHIA 8, HERWIG 7) requires application of the correction to real data.

Comparison to PYTHIA 8 Monash and HERWIG 7 particle-level predictions will be performed in Phase 4b/4c.

Quantitative comparison to published ATLAS [3] and CMS [4] Lund plane data via χ^2 with full covariance will be included at Doc 4c.

9 Conclusions

This analysis note presents the expected results for the measurement of the primary Lund jet plane density in hadronic Z decays at $\sqrt{s} = 91.2$ GeV using archived ALEPH data. The measurement uses approximately 3.05 million events (5.69 million thrust hemispheres), corrected to charged-particle level using bin-by-bin correction factors derived from PYTHIA 6.1 Monte Carlo.

The key findings are:

1. The bin-by-bin correction method is validated by the split-sample closure test ($\chi^2/\text{ndf} = 40.71/58$, $p = 0.96$) and all 12 split-sample stress tests.

2. Iterative Bayesian unfolding is formally downscoped from co-primary to cross-check [D9] due to the fundamental inability to match per-splitting migrations.

3. The total uncertainty ranges from $\sim$1% to $\sim$8% (relative) across populated bins, dominated by MC model dependence in the non-perturbative region.

4. The Lund plane density shows the expected triangular structure with a perturbative plateau consistent with the LO QCD prediction $\rho_{\text{LO}}^{\text{charged}} \approx 0.067$.

5. The result is insensitive to the jet definition (thrust hemispheres vs. Durham jets: $\chi^2/\text{ndf} = 0.185$) and to the hardness variable ordering ($<5\%$ in the perturbative core).



Figure 23: Left: reco/gen 1D comparison from Phase 2 exploration, showing the detector-level distortion of the Lund plane projections. Right: correction factor preview ($N_{\text{gen}}/N_{\text{reco}}$, post-selection only), ranging from 0.8 to 1.2 in the core. These demonstrate that the detector response is close to unity and the bin-by-bin correction is well-motivated.

This constitutes the first full two-dimensional primary Lund jet plane density measurement in e^+e^- collisions.

10 Future Directions

The following improvements are genuinely infeasible within the current analysis framework and would require substantial additional resources:

1. **Alternative generator MC production.** Full detector simulation with PYTHIA 8 or HERWIG 7 would provide independent correction factors and a robust model-dependence assessment. This requires access to the ALEPH detector simulation framework, which is not available in the archived dataset.

2. **Secondary Lund plane.** Measuring the secondary declustering branch requires separate validation infrastructure (different correction factors, different migration patterns) and would approximately double the analysis scope.

3. **NLL analytical comparison.** The NLL calculation of the primary Lund plane density by Lifson, Salam, and Soyez [12] provides 5–7% precision predictions for pp jets. Adapting these predictions to e^+e^- quark jets requires input from the theory authors.

4. **Flavour-separated measurement.** Separating b-quark and light-quark hemispheres using the available lifetime information would test the dead-cone effect [5] at the Z pole. This requires a dedicated b-tagging strategy not developed in this analysis.

11 Known Limitations

1. **IBU failure [D9].** The iterative Bayesian unfolding method fails due to the inability to match individual Lund splittings between reco and gen levels. Three remediation attempts were made (iteration optimization, uncapped response matrix, Tikhonov regularization); all failed with $\chi^2/\text{ndf} > 2000$. *Impact:* No independent unfolding cross-check with an alternative method. *Mitigation:* The split-sample closure and stress tests validate the bin-by-bin method directly; published ATLAS and CMS analyses also use bin-by-bin as primary. *What would fix it:* Proper per-splitting matching between reco and gen C/A trees, which requires event-level reco/gen particle matching before clustering.

2. **Single MC generator.** Only PYTHIA 6.1 is available with full detector simulation. The model-dependence systematic is evaluated via reweighting, which assumes locally linear detector response. *Impact:* The model systematic may be underestimated in the non-perturbative region where different generators disagree most. *Mitigation:* Phase 4b/4c comparison with standalone PYTHIA 8 and HERWIG 7 at particle level. *What would fix it:* Full detector simulation with alternative generators.

3. **Track-quality branch (`ntpc`) inaccessible.** The number of TPC hits per track is not accessible in the archived data, preventing the standard variation of the TPC hit requirement. *Impact:* One element of the track selection systematic cannot be evaluated. *Mitigation:* The momentum threshold and impact parameter variations cover the primary track quality effects. *What would fix it:* Access to the full ALEPH reconstruction output with all track quality variables.

4. **Systematic evaluation on MC subset.** Detector-level and selection-cut systematics were evaluated on a 10-file MC subset. *Impact:* Systematic shifts may fluctuate by $\sim 2\times$ due to the reduced statistics. *Mitigation:* Full-sample evaluation in Phase 4b/4c. *What would fix it:* Processing all 40 MC files for every variation.

5. **Hardness variable [D13].** Energy ordering as a systematic variation was not evaluated; only the cross-check comparison was performed. *Impact:* The systematic from the ordering choice is unquantified. *Mitigation:* Phase 4b/4c implementation. *What would fix it:* Full correction chain with energy-ordered declustering.

References

[1] Frédéric A. Dreyer, Gavin P. Salam, and Grégory Soyez. The Lund Jet Plane. *JHEP*, 12:064, 2018.

[2] Yu. L. Dokshitzer, G. D. Leder, S. Moretti, and B. R. Webber. Better jet clustering algorithms. *JHEP*, 08:001, 1997.

[3] ATLAS Collaboration. Measurement of the Lund Jet Plane in the Search for Physics Beyond the Standard Model. *Phys. Rev. Lett.*, 124:222002, 2020.

[4] CMS Collaboration. Measurement of the primary Lund jet plane density in proton-proton collisions at $\sqrt{s} = 13$ TeV. *JHEP*, 05:116, 2024.

[5] LHCb Collaboration. Measurements of the Lund jet plane for light- and beauty-quark jets. *Phys. Rev. D*, 112:072015, 2025.

[6] ATLAS Collaboration. Measurement of the Lund jet plane using charged particles in hadronic decays of top quarks and W bosons. *Eur. Phys. J. C*, 2025.

[7] DELPHI Collaboration. A study of the energy evolution of event shape distributions and their means with the DELPHI detector at LEP. *Eur. Phys. J. C*, 55:1–48, 2008.

[8] ALEPH Collaboration. Subjet structure of b and light-quark jets at LEP. ALEPH internal note, cds_388806.

[9] Torbjorn Sjöstrand et al. An introduction to PYTHIA 8.2. *Comput. Phys. Commun.*, 191:159–177, 2015.

[10] Peter Skands, Stefano Carrazza, and Juan Rojo. Tuning PYTHIA 8.1: the Monash 2013 Tune. *Eur. Phys. J. C*, 74:3024, 2014.

[11] Johannes Bellm et al. Herwig 7.0/Herwig++ 3.0 release note. *Eur. Phys. J. C*, 76:196, 2016.

[12] Alexander Lifson, Gavin P. Salam, and Grégory Soyez. Calculating the primary Lund Jet Plane density. *JHEP*, 10:170, 2020.

[13] ALEPH Collaboration. Determination of α_s from hadronic event shapes at $\sqrt{s} = 91.2$ GeV. inspire:322679.

[14] ALEPH Collaboration. Measurements of α_s from hadronic event shapes. ALEPH internal note, CERN-PPE/97-002, cds_2876991.

[15] Particle Data Group. Review of Particle Physics. *Phys. Rev. D*, 110:030001, 2024.

[16] Torbjorn Sjöstrand et al. High-energy physics event generation with PYTHIA 6.1. *Comput. Phys. Commun.*, 135:238–259, 2001.

[17] ALEPH Collaboration. Performance of the ALEPH detector at LEP. *Nucl. Instrum. Meth. A*, 360:481–506, 1995.

[18] ALEPH Collaboration. A measurement of the QCD colour factors and a limit on the light gluino. *Z. Phys. C*, 76:1–14, 1997. inspire:430544.

[19] Matteo Cacciari, Gavin P. Salam, and Grégory Soyez. FastJet User Manual. *Eur. Phys. J. C*, 72:1896, 2012.

[20] Giulio D'Agostini. A multidimensional unfolding method based on Bayes' theorem. *Nucl. Instrum. Meth. A*, 362:487–498, 1995.

A Per-bin Systematic Tables

The full per-bin systematic shifts are available in machine-readable format in `results/systematics.json`. Table 13 shows a representative sample of the bin-by-bin systematic shifts for the three dominant sources in five representative bins.

Table 13: Per-bin systematic shifts for three dominant sources in five representative bins across the Lund plane. Values are the absolute shift in density $\Delta\rho$. The full 58-bin tables for all 13 sources are provided in the machine-readable JSON.

i (x-bin)	j (y-bin)	ρ_{nom}	$\Delta\rho_{\text{model}}$	$\Delta\rho_{\text{p-thr}}$	$\Delta\rho_{\text{track}}$
3	3	0.572	±0.040	±0.010	±0.001
4	4	0.744	±0.032	±0.011	±0.001
5	3	0.600	±0.026	±0.010	±0.001
7	2	0.240	±0.007	±0.001	±0.001
9	1	0.046	±0.001	<0.001	<0.001

B Covariance Matrices

The statistical covariance is constructed from 500 bootstrap replicas with full correction chain resampling. The systematic covariance is the sum of rank-1 matrices from each source's symmetrised shift vector. The total covariance matrix (58×58 for populated bins) is positive semi-definite with condition number $< 10^{10}$.

The off-diagonal correlations arise primarily from two sources: (1) the MC model dependence systematic, which shifts the entire plane coherently and produces large-scale positive correlations, and (2) the self-normalization constraint ($\rho \propto 1/N_{\text{jet}}$), which introduces anticorrelations between bins. The statistical covariance from the bootstrap captures both effects.

The correlation matrix is shown in Figure 17. The maximum off-diagonal correlation $|\rho_{ij}|$ is driven by the model-dependence systematic.

Recommendation for downstream use: The full covariance matrix should be used for any χ^2-based comparison with the measured Lund plane. The boundary bins (first and last populated bins in each row of the 2D grid) have larger uncertainties and correlations; fits should consider restricting to the core region for maximum sensitivity.

The full covariance matrices (statistical, per-source systematic, and total) are provided in machine-readable JSON format at `results/covariance.json`.

C Extended Cutflow

The cutflow in Table 7 (Section 3.3) presents the main data and MC event counts. For completeness, the generator-level cutflow is:

Table 14: Generator-level cutflow for the MC sample. The genBefore sample contains all generated events before event selection; the gen sample applies the same particle-level cuts used at reco level.

Cut	gen events	gen eff. [%]	genBefore events	genBefore eff. [%]
Total	771 597	100.0	973 769	100.0
$N_{\text{ch}} \geq 5$	771 468	99.98	973 584	99.98
$N_{\text{hemi}} \geq 2$	767 465	99.46	968 356	99.44
Final hemis.	1 534 930		1 936 712	

The `passesAll` flag in the genBefore tree is uniformly `False`, confirming that this sample contains all generated events regardless of event selection. The gen/genBefore hemisphere ratio (1,534,930/1,936,712 = 0.793) gives the event selection efficiency at generator level.

D Auxiliary Plots

Additional data/MC comparisons and validation plots from the exploration phase are collected here for reference.



Figure 24: Left: momentum distribution on logarithmic scale, showing data/MC agreement across four orders of magnitude. Right: reco-level Lund plane density from Phase 2 exploration (100 000 events, 200 000 hemispheres). The characteristic triangular structure with the perturbative plateau is clearly visible.



Figure 25: Self-closure test (same-MC) projections: $\ln k_T$ (left) and $\ln(1/\Delta\theta)$ (right). The corrected MC reco (red) recovers the genBefore truth (black) exactly ($\chi^2/\mathrm{ndf} = 0.0/58$), as expected for the algebraic identity. This confirms the implementation is correct.

E Limitation Index

F Reproduction Contract

The full analysis can be reproduced with the following sequence of commands from the analysis root directory (`analyses/lund_jet_plane/`):



Figure 26: Pull distribution for the self-closure test. All pulls are identically zero, confirming the algebraic identity of the bin-by-bin correction when applied to the same MC sample.



Figure 27: Full-MC (non-split) stress tests: $\ln k_T$ tilt (left) and $\ln(1/\Delta\theta)$ tilt (right). The full-MC stress tests are algebraically trivial (same as self-closure with reweighting), so perfect recovery is expected. The split-sample stress tests (Figure 10) provide the genuine validation.



Figure 28: Full-MC stress test with 2D correlated tilt. Same comment as Figure 27: the full-MC test is algebraically exact.



Figure 29: Left: closure pull distribution from the Phase 4a initial computation (prior to split-sample re-analysis). Right: correlation matrix from the initial computation. Both were superseded by the updated analysis (Figures 9 and 17), which includes bootstrap with full correction chain resampling.



Figure 30: Per-systematic impact on $\ln k_T$ (left) and $\ln(1/\Delta\theta)$ (right) projections from the initial Phase 4a computation (prior to removal of the IBU-vs-BBB systematic). The updated systematic impact plots (Figure 14) supersede these.



Figure 31: Initial systematic breakdown (left) and uncertainty summary (right) from the Phase 4a initial computation. Superseded by Figures 13 and 22.

Table 15: Complete registry of constraints [A], limitations [L], and decisions [D] from the analysis strategy, with their current status.

Label	Type	Description	Status
[A]	Constraint	Particle-level: charged, $p > 200$ MeV/c, $c\tau > 1$ cm	Implemented
[D1]	Decision	C/A clustering for Lund plane	Implemented
[D2]	Decision	Primary declustering chain	Implemented
[D3]	Decision	10×10 uniform binning	Implemented
[D4]	Decision	`pwflag == 0` (charged only)	Implemented
[D5]	Decision	Thrust cut $T > 0.7$	Implemented
[D6]	Decision	MVA not used (no S/B needed)	Justified
[D7]	Decision	Durham 2-jets as Approach C	Implemented
[D8]	Decision	BBB as co-primary correction	Primary
[D9]	Decision	IBU downscoped to cross-check	Downscoped
[D10]	Decision	ISR/FSR treatment	Implemented
[D11]	Decision	Bootstrap covariance, 500 rep.	Implemented
[D12]	Decision	Bin-level response matrix	Implemented
[D13]	Decision	p_T ordering primary	Partial
[D14]	Decision	Sherpa comparison deferred	Deferred
[L]	Limitation	Luminosity from event count	Documented
[L]	Limitation	Single MC (PYTHIA 6.1)	Documented
[L]	Limitation	`ntpc` inaccessible	Documented

```
pixi install           # Environment setup

# Phase 2: Exploration
pixi run py phase2_exploration/src/01_explore_data.py
pixi run py phase2_exploration/src/02_data_mc_comparison.py
pixi run py phase2_exploration/src/03_lund_plane_reco.py
pixi run py phase2_exploration/src/04_binning_optimization.py
pixi run py phase2_exploration/src/05_ordering_comparison.py

# Phase 3: Selection
pixi run py phase3_selection/src/01_apply_selection.py
pixi run py phase3_selection/src/02_correction_factors.py
pixi run py phase3_selection/src/03_closure_test.py
pixi run py phase3_selection/src/04_data_mc_lund.py
pixi run py phase3_selection/src/05_approach_comparison.py

# Phase 4a: Inference (expected)
pixi run py phase4_inference/4a_expected/src/01_corrected_lund.py
pixi run py phase4_inference/4a_expected/src/02_systematics.py
pixi run py phase4_inference/4a_expected/src/03_ibu.py
pixi run py phase4_inference/4a_expected/src/04_covariance.py
pixi run py phase4_inference/4a_expected/src/05_split_closure.py
pixi run py phase4_inference/4a_expected/src/06_stress_tests.py

# Analysis note compilation
cd analysis_note && tectonic ANALYSIS_NOTE_doc4a_v1.tex
```

Data path: The `.analysis_config` file must contain the path to the archived ALEPH data directory. The 6 data files and 40 MC files total approximately 25 GB.

Runtime estimate: Phase 2: ∼5 min. Phase 3: ∼30 min (8-worker parallel processing). Phase 4a: ∼45 min. Total: ∼1.5 hours on a modern multi-core machine.