

SUPPORTING INFORMATION

Optimizing Vanadium-Catalyzed ESA Reactions: Machine Learning-Driven Yield Predictions and Data Augmentation

José Ferraz-Caetano¹, Filipe Teixeira², M. Natália D. S. Cordeiro¹

1) LAQV-REQUIMTE – Department of Chemistry and Biochemistry – Faculty of Sciences,
University of Porto - Rua do Campo Alegre, S/N, 4169-007 Porto, Portugal

2) CQUM – Centre of Chemistry, University of Minho, Campus de Gualtar, 4710-057 Braga,
Portugal

*Corresponding Authors: **José Ferraz-Caetano** (jose.caetano@fc.up.pt);

M. Natália D. S. Cordeiro (ncordeir@fc.up.pt)

INDEX

Annex S1 – Model Database	2
Annex S2 – Data Augmentation Protocol	3
Annex S3 – Data Augmentation Distribution Analysis.....	5
Annex S4 – Results and Model Optimization Features.....	7
Annex S5 – RDKit Full Descriptor List	8
Annex S6 – Experimental Descriptor List.....	11
Annex S7 – Sample Code for ML Model Design	12
References	14

Annex S1 – Model Database

Complete database, featuring individual bibliographical source, and the list of calculated and experimental descriptors is available at the following repository: <https://gitfront.io/r/ESA-Vanadium/biMx4obXpmkj/ESA-Vanadium-Yield/>. In the database file, we include individual digital identification of all bibliographical sources used to populate the dataset. The database is also presented in our previous article¹: <https://doi.org/10.1039/D3NJ05784D>.

The digital representation (mol file) of each chemical structure involved is also presented in the repository, in the folder “Mol Files”. The chemical species are sorted in each descriptor name: ‘_Cat’ for Catalyst, ‘_Sol’ for Solvent, ‘_Lig’ for Ligand and ‘_Sub’ for Substrate.

We also provide code for descriptor generation in the online repository.

Annex S2 – Data Augmentation Protocol

The data augmentation protocol expands the dataset while ensuring that the generated synthetic data remains within experimentally observed boundaries. It is designed to conform to chemical plausibility and enhance the performance of the ML model. The full script is on the article's [online repository](#).

1. Dataset Preprocessing and Constraint Definition

The dataset is first cleaned by filling missing values and converting all numeric variables to the correct format. The numeric variables are:

- Temperature (Temp_K)
- Yield (%)
- Enantiomeric Excess (EE, %)
- Substrate Quantity (Substrate_quant_mmol)
- Catalyst Quantity (Catalyst_quant_mmol)
- Ligand Quantity (Ligand_quant_mmol)
- Oxidant Quantity (Oxidant_quant_mmol)
- Additive Quantity (Additive_quant_mL)
- Solution Volume (Solution_vol_mL)
- Reaction Time (Time_h)

The dataset is then grouped by Laboratory, and for each numeric variable, the minimum and maximum values are determined within each lab's dataset. These values will be the experimental bounds that guarantee synthetic data will not exceed beyond the limits of real experimentation observations.

2. Synthetic Data Generation with Controlled Variation

For each laboratory, 20 synthetic rows are generated by randomly selecting an existing real data point and applying a variation of up to 10% to its numeric values. This variation is constrained so that no synthetic value exceeds the observed minimum or maximum range for that laboratory. For Yield (%) and Enantiomeric Excess (EE, %), additional restrictions ensure that values remain between 0% and 100% to maintain chemical relevance. Each synthetic data point is labeled with a new column ('is_synthetic = 1'), distinguishing it from the original dataset.

3. Addressing Underrepresented Yield Ranges

To ensure a balanced dataset, the Yield (%) variable is divided into 10 discrete bins based on its distribution. The number of data points in each bin is compared to the average count per bin, and bins with fewer than 50% of the average count are identified as underrepresented. 20 additional synthetic data points are generated for each underrepresented bin to create a more even distribution.

4. One-Hot Encoding of Categorical Variables

To prepare the dataset for machine learning, categorical variables (Catalyst Structure, Laboratory, Catalyst, Substrate, Ligand, Oxidant, and Solvent) are transformed using one-hot encoding. Any missing categorical values are replaced with "missing" before encoding.

5. Final Data Storage

The final dataset, now containing both real and synthetic data, is saved as for further modeling.

Annex S3 – Data Augmentation Distribution Analysis

To support the argument of Data Augmentation (DA) for our ESA dataset, t-SNE (t-distributed Stochastic Neighbor Embedding) plots can be instrumental in visualizing how the augmented data represents the original dataset. The t-SNE plot is a powerful, by its non nonlinear method for the visualization of high-dimensional data by projecting back into two-dimensional or three-dimensional plots. They are useful for revealing hidden relations, patterns, and clusters in complex datasets that are usually very hard to understand. By comparing the t-SNE plots of the datasets before and after DA, we will assess whether the augmentation succeeded in extending the coverage of the datasets and filling in underrepresented areas. Figure A1 represents the t-SNE plots before and after DA. A two-component separation was used as it proved to be the best score on data distribution for the DA dataset (2 components = 0.6917; 3 components = 0.6449).

This chart provides evidence that supports data augmentation in our model. In the original dataset, there is a clear clustering of certain catalysts on the t-SNE plots, like VO(acac)₂ and VO(OiPr)₃, suggesting that the dataset may be biased towards particular catalysts, with some regions of chemical space underrepresented. After applying data augmentation, the second plot demonstrates a more balanced and expanded coverage of the chemical space, with the catalyst structures more evenly distributed. Although the KNN preservation score drops in the augmented dataset, this suggests that the model sacrifices precision in nearest-neighbor relationships for a broader representation of the chemical space. Such trade-offs render the model better capable of generalizing across other catalysts.




Figure A1. ESA data distribution t-SNE plots before and after DA using 2 components.

Annex S4 – Results and Model Optimization Features

Best hyper-parameters for each algorithm for model optimization:

Random Forest Regressor = n_estimators= 150, min_samples_split= 3, min_samples_leaf= 1, max_features='sqrt', max_depth=None, bootstrap= False, random_state=47

Gradient Boosting Regressor = n_estimators=150, min_samples_split=4, min_samples_leaf= 3, max_depth=40, random_state=47

Cross validation scoring result

CV = RepeatedKFold(n_splits=2, n_repeats=3, random_state=47)

Scoring Mode = 'neg_mean_absolute_error', cv=cv, n_jobs=-1, error_score='raise'

Cross Validation STD Score RF Optimized Model = 1.7 ± 0.8 (Average MAE = 4.7 ± 0.3)

Cross Validation STD Score GB Optimized Model = 1.9 ± 0.9 (Average MAE = 4.3 ± 0.7)

The scripts for algorithm development and hyperparameter optimization are in the article's [online repository](#), in the “Scripts” folder. The full statistical results are available in the “Results” folder of the [online repository](#).

Annex S5 – RDKit Full Descriptor List

Descriptors used in Machine Learning (ML) model development (Table S1) retrieved from RDKit software. The respective bibliographical foundation of each described is described in detail in the RDKit WebBook Documentation². The descriptor calculations were made by converting each SMILES string (representing each molecular entry), running on top of Python version 3.9³ and using RDKit package, version 2022.09.4⁴. The Chemical information Type is presented in Table S1, and the chemical species are sorted in the database file in the online repository.

Table S1

List of RDKit Descriptors used in ML model development.

Descriptor	Chemical information Type	Meaning
BalabanJ	Structural	Distance sum of the two end-vertex for each edge. BalabanJ index has been proven to be relevant to network branching.
BertzCT	Structural	Complexity index, taking into account both the variety of kinds of bond connectivities and atom types; information contents related to bond connectivity and atom type diversity.
$0\chi, 1\chi$	Structural	This descriptor signifies a retention index (zero order) derived directly from gradient retention times.
$0\chi_n - 4\chi_n$	Structural	This descriptor signifies a retention index (zero order) derived directly from gradient retention times.
$0\chi_v - 4\chi_v$	Structural	This descriptor signifies atomic valence connectivity index (order 0).
VSA_EState1 – VSA_EState11	Surface Area	MOE-type (QSAR model) descriptors using EState indices and surface area contributions.
Hall Kier α	Structural	Descriptor fingerprint that displays the difference between active and inactive molecules.

HeavyAtomCount	Structural	The number of heavy atoms in the molecule.
HeavyAtomMolWt	Structural	The average molecular weight of the molecule ignoring hydrogens.
$\kappa_1, \kappa_2, \kappa_3$	Structural	This descriptor signifies # κ shape index: $(n-1) \times 2 / m^2$
<i>Max and Min AbsEStateIndex</i>	Electronic	Maximum and minimum Absolute E-State
<i>Max and Min AbsPartialCharge</i>	Electronic	Maximum and minimum Absolute Partial Charge
<i>Max and Min EStateIndex</i>	Electronic	Maximum and minimum E-State
<i>Max and Min PartialCharge</i>	Electronic	Maximum and minimum Partial Charge
Mol logP	Structural	Wildman-Crippen logP value.
MolMR	Electronic	Wildman-Crippen molar refractivity.
MolWt	Structural	The average molecular weight of the molecule.
NHOH Count	Structural	The number of NHs or OHs.
NO Count	Structural	The number of Nitrogens and Oxygens.
<i>n_{alcarb}</i>	Structural	The number of aliphatic carbocycles.
<i>n_{alihet}</i>	Structural	The number of aliphatic heterocycles.
<i>n_{alirig}</i>	Structural	The number of aliphatic rings.
<i>n_{arocarb}</i>	Structural	The number of aromatic carbocycles.
<i>n_{arohet}</i>	Structural	The number of aromatic heterocycles.
<i>n_{arorig}</i>	Structural	The number of aromatic rings.
<i>n_{Ha}</i>	Structural	The number of Hydrogen Bond Acceptors.
<i>n_{Hd}</i>	Structural	The number of Hydrogen Bond Donors.
<i>n_{het}</i>	Structural	The number of Heteroatoms.
<i>n_{radele}</i>	Structural	The number of radical electrons.

<i>n_{rot}</i>	Structural	The number of Rotatable Bonds.
<i>n_{satrig}</i>	Structural	The number of saturated rings.
<i>n_{ele}</i>	Structural	The number of valence electrons.
PEOE_VSA1 – PEOE_VSA14	Surface Area	MOE-type (QSAR model) descriptors using partial charges and surface area contributions.
SMR_VSA1 – SMR_VSA10	Surface Area	MOE-type (QSAR model) descriptors using MR contributions and surface area contributions.
SlogP_VSA1 – SlogP_VSA12	Surface Area	MOE-type (QSAR model) descriptors using logP contributions and surface area contributions.
TPSA	Surface Area	The total polar surface area of a molecule based upon fragment calculations.
VSA_EState1 – VSA_EState10	Surface Area	MOE-type (QSAR model) descriptors using EState indices and surface area contributions.

Annex S6 – Experimental Descriptor List

Experimental descriptors used in ML model development (Table S2) retrieved from each bibliographical source. Specific solvent parameters were retrieved from the Minnesota Solvent Descriptor Database⁵.

Table S2. List of experimental descriptors used in machine learning model development.

Descriptor	Chemical information Type	Meaning
Temp_K	Experimental	Reaction temperature in Kelvin.
Yield	Experimental	Reaction yield.
EE	Experimental	Reaction enantiomeric excess.
Configuration	Experimental	Product geometrical configuration.
Substrate_quant_mmol	Experimental	Quantity of reaction substrate in mmol.
Catalyst_quant_mmol	Experimental	Quantity of reaction catalyst in mmol.
Ligand_quant_mmol	Experimental	Quantity of reaction catalyst ligand in mmol.
Oxidant_quant_mmol	Experimental	Quantity of reaction oxidant in mmol.
Additive_quant_mL	Experimental	Volume of reaction additive in mL.
Solution_vol_mL	Experimental	Volume of reaction solution in mL.
Time_h	Experimental	Reaction duration in hours.
Solvent	Experimental	Reaction solvent.
Solv_opt_freq20	Experimental	Solvent index of refraction at optical frequencies at 293 K.
Solv_opt_freq25	Experimental	Solvent index of refraction at optical frequencies at 298 K.
Solv_Hbond_ac	Experimental	Solvent Abraham's hydrogen bond acidity.
Solv_Hbond_bs	Experimental	Solvent Abraham's hydrogen bond basicity.
Solv_surf_tens	Experimental	Solvent surface tension.
Solv_diele_cst	Experimental	Solvent dielectric constant at 298 K.

Annex S7 – Sample Code for ML Model Design

The scripts for algorithm development are in the articles [online repository](#), in the folder Scripts.

Scripts Overview

1. mol-conversion.py

This script provides scripts molecular data processing and conversion. It transforms molecular structures contained within the dataset into formats applicable for ML models. Key functionalities include:

- Converting molecular data (e.g., SMILES strings) into descriptor formats suitable for model input.
- Handling molecule-based transformations, ensuring that molecular representations are standardized and compatible with the model.
- The output of this script is typically used as input features for subsequent model training and evaluation.

2. model_preliminary_run.py

This script provides initial evaluations of ML models to deliver baseline performance metrics. The model evaluation is performed without hyperparameter tuning on a set of algorithms as a starting point to gauge how well the models perform on the initial dataset. Functions include:

- Loading the dataset and preparing it for modeling.
- Running baseline models: Random Forest, Gradient Boosting, Support Vector Machines, Neural Networks.
- Outputting preliminary performance metrics (R^2 , MAE, RMSE) to help guide future model selection and optimization steps.

3. ESA_data_augmentation.py

This script is used for **data augmentation** on the ESA-Vanadium dataset. It enriches the dataset by generating synthetic data points or applying transformations to existing data. Key functions of this script include:

- Handling missing values or filling gaps in the dataset.
- Augmenting the dataset to expand feature space, enabling robust model training.
- Ensures that augmented data remains representative of the original chemical reaction conditions, focusing on maintaining chemical consistency.

4. ESA-hyperparameter-opt.py

This script performs hyperparameter optimization for a Random Forest model using the RandomizedSearchCV function from scikit-learn. The goal is to optimize the model's performance by searching over a predefined range of hyperparameters. The script:

- Loads the ESA-Vanadium dataset, excluding certain irrelevant features.
- Conducts Random Forest regression, applying a hyperparameter search over parameters like the number of estimators, depth, and minimum samples split.
- Evaluates model performance using metrics such as R^2 , mean absolute error (MAE), and root mean squared error (RMSE) over multiple iterations.
- Saves the optimized model results and the best hyperparameters to a CSV file.

5. ESA_model_opt_run.py

This script focuses on the **execution and evaluation** of the optimized RF ML model on the augmented ESA-Vanadium dataset. It is responsible for:

- Collecting results from each model's training and testing phases.
- Recording performance metrics, including training and test R^2 , MAE, and RMSE for each model and iteration.
- Saving the evaluation metrics and model-specific results for further analysis or comparison.

6. partial_dependency_plots.py

This script generates partial dependence plots (PDPs) to interpret model behavior. PDPs show how each feature impacts the predicted outcome, allowing for better understanding of the model's decision-making process. The script:

- Loads the trained model and dataset.
- Generates plots that display the relationship between individual features and the model's predictions.
- Visualizes feature importance and dependencies, helping to uncover key drivers in the reaction outcomes.

7. weight_analysis.py

The script records optimized RF model performance across 10 iterations with varying synthetic data weights. These weights, ranging from 0.1 to 0.9, adjust the contribution of synthetic reactions in the training data, simulating different levels of experimental noise.

References

- (1) Ferraz-Caetano, J.; Teixeira, F.; Cordeiro, M. N. D. S. Navigating epoxidation complexity: building a data science toolbox to design vanadium catalysts. *New Journal of Chemistry* **2024**, 48 (12), 5097-5100, 10.1039/D3NJ05784D. DOI: 10.1039/D3NJ05784D.
- (2) *RDKit: Open-source cheminformatics - Descriptor Webbook*. <https://www.rdkit.org/docs/GettingStartedInPython.html#list-of-available-descriptors> (accessed January 18th, 2024).
- (3) *Python Software Foundation - Python Language Reference, version 3.9.8*. <http://www.python.org> (accessed January 18th, 2024).
- (4) Landrum, G. *RDKit: Open-source cheminformatics 2022_09_4 (Q3 2022) Release - January 16, 2023*. 2023. <http://www.rdkit.org/> (accessed January 18th, 2024).
- (5) Marenich, A. V. K., Casey P; Thompson, Jason D; Hawkins, Gregory D; Chambers, Candee C; Giesen, David J; Winget, Paul; Cramer, Christopher J; Truhlar, Donald G. *Minnesota Solvation Database (MNSOL) version 2012*. Retrieved from the Data Repository for the University of Minnesota. 2020. <https://doi.org/10.13020/3eks-j059> (accessed January 18th, 2024)