

Synthetic-Inductive Category Theory

Jacob Neumann

University of Nottingham & Reykjavik University

TYPES 2025, Glasgow, Scotland

09 June 2025

This talk corresponds to [Neu25, Chapter 4],
draft available at

jacobneu.com/PhD

Martin L^öf Type Theory: Synthetic groupoid theory

Synthetic groupoid structure of identity types

$t : A$



Synthetic groupoid structure of identity types




$$\frac{t : A}{\text{refl}_t : \text{Id}(t, t)}$$

Synthetic groupoid structure of identity types




$$p: \text{Id}(t, t')$$

Synthetic groupoid structure of identity types




$$p: \text{Id}(t, t')$$
$$\underline{q: \text{Id}(t', t'')}$$

Synthetic groupoid structure of identity types



$$\frac{p: \text{Id}(t, t') \quad q: \text{Id}(t', t'')}{p \cdot q: \text{Id}(t, t')}$$


Synthetic groupoid structure of identity types



$$\frac{p: \text{Id}(t, t') \quad q: \text{Id}(t', t'')}{p \cdot q: \text{Id}(t, t')}$$


$$p \cdot q := J_{\text{Id}(t, _)} p (t'', q)$$

Synthetic groupoid structure of identity types




$$\underline{p: \text{Id}(t, t')}$$

Synthetic groupoid structure of identity types



$$\frac{p : \text{Id}(t, t')}{p^{-1} : \text{Id}(t', t)}$$

Synthetic groupoid structure of identity types




$$\frac{p : \text{Id}(t, t')}{p^{-1} : \text{Id}(t', t)}$$

$$p^{-1} := J_{\text{Id}(_, t)} \text{ refl}_t (t', p)$$


Directed TT: Synthetic *category* theory

Synthetic category structure of hom-types




$$\frac{t : A}{\text{refl}_t : \text{Hom}(t, t)}$$

Synthetic category structure of hom-types



$$\frac{t : A}{\text{refl}_t : \text{Hom}(t, t)}$$


Synthetic category structure of hom-types



$$\frac{p: \text{Hom}(t, t') \quad q: \text{Hom}(t', t'')}{p \cdot q: \text{Hom}(t, t'')}$$

$$p \cdot q := J_{\text{Hom}(t, _)} p (t'', q)$$


Synthetic category structure of hom-types



$$\frac{p: \text{Hom}(t, t') \quad q: \text{Hom}(t', t'')}{p \cdot q: \text{Hom}(t, t'')}$$

$$p \cdot q := J_{\text{Hom}(t, _)} p (t'', q)$$


Synthetic category structure of hom-types



$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(t', t)}$$

$$p^{-1} := J_{\text{Hom}(_, t)} \text{ refl}_t (t', p)$$

Synthetic category structure of hom-types



$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(t', t)}$$

$$p^{-1} := J_{\text{Hom}(_, t)} \text{ refl}_t (t', p)$$

Problem of Directed
TT: Make symmetry
unprovable

Problem of Directed TT: Make symmetry unprovable

*Synthetic categories that aren't (necessarily)
synthetic groupoids*

Polarized and Directed type theory

We have **polarity annotations** on our types to mark co- or contra-variance

$$\frac{\Delta \vdash A \text{ type}}{\Delta \vdash A^- \text{ type}} \qquad \overline{\Delta \vdash (A^-)^- = A}$$

Polarized and Directed type theory

We have **polarity annotations** on our types to mark co- or contra-variance

$$\frac{\Delta \vdash A \text{ type}}{\Delta \vdash A^- \text{ type}} \quad \overline{\Delta \vdash (A^-)^- = A}$$

Types are categories, A^- is the opposite category of A

Polarized and Directed type theory

We have **polarity annotations** on our types to mark co- or contra-variance

$$\frac{\Delta \vdash A \text{ type}}{\Delta \vdash A^- \text{ type}} \quad \overline{\Delta \vdash (A^-)^- = A}$$

Types are categories, A^- is the opposite category of A

The polarity annotations allow us to properly state the variance of hom-sets [Nor19]:

$$\frac{\Delta \vdash t : A^- \quad \Delta \vdash t' : A}{\Delta \vdash \text{Hom}(t, t') \text{ type}}$$

For **closed** terms, we can coerce between A and A^- [NA25, Neu25]:

$$\frac{t : A^-}{-t : A} \quad \frac{}{- - t = t}$$

Closed term coercion

For **closed** terms, we can coerce between A and A^- [NA25, Neu25]:

$$\frac{t : A^-}{-t : A} \quad \overline{- - t = t}$$

A category and its opposite have the same objects

Closed term coercion

For **closed** terms, we can coerce between A and A^- [NA25, Neu25]:

$$\frac{t : A^-}{-t : A} \quad \frac{}{- - t = t}$$

Key Point Cannot (in general) negate open terms

$$x : A^- \vdash -x : A$$

Closed term coercion

For **closed** terms, we can coerce between A and A^- [NA25, Neu25]:

$$\frac{t : A^-}{-t : A} \quad \frac{}{- - t = t}$$

Key Point Cannot (in general) negate open terms

$$x : A^- \vdash -x : A$$

For **closed** terms, we can coerce between A and A^- [NA25, Neu25]:

$$\frac{t : A^-}{-t : A} \quad \frac{}{- - t = t}$$


For **closed** terms, we can coerce between A and A^- [NA25, Neu25]:

$$\frac{t : A^-}{-t : A} \quad \frac{}{- - t = t}$$

The coercions make it possible to introduce refl:


$$\frac{t : A^-}{\text{refl}_t : \text{Hom}(t, -t)}$$

Synthetic category structure of hom-types



$$\frac{t : A}{\text{refl}_t : \text{Hom}(t, t)}$$

Synthetic category structure of hom-types



$$\frac{t : A^-}{\text{refl}_t : \text{Hom}(t, t)}$$

Synthetic category structure of hom-types



$$\frac{t : A^-}{\text{refl}_t : \text{Hom}(t, -t)}$$

Synthetic category structure of hom-types



$$\frac{t : A^-}{\text{refl}_t : \text{Hom}(t, -t)}$$

Coslice Path Induction


$$t: A^-$$




Coslice Path Induction


$$t: A^-$$
$$x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type}$$

Coslice Path Induction


$$t : A^-$$
$$x' : A, u : \text{Hom}(t, x') \vdash M(x', u) \text{ type}$$


Coslice Path Induction



$t: A^-$
 $x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type}$


$x': A, u: \text{Hom}(t, x') \vdash J_M^+ m (x', u): M(x', u)$

Coslice Path Induction




$$\begin{array}{c}
 t: A^- \\
 x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type} \\
 m: M(-t, \text{refl}_t) \\
 \hline
 x': A, u: \text{Hom}(t, x') \vdash J_M^+ m (x', u): M(x', u)
 \end{array}$$

Coslice Path Induction




$$\begin{array}{c}
 t: A^- \\
 x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type} \\
 m: M(-t, \text{refl}_t) \\
 \hline
 x': A, u: \text{Hom}(t, x') \vdash J_M^+ m (x', u): M(x', u)
 \end{array}$$

Coslice Path Induction



$$\begin{array}{c}
 t: A^- \\
 x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type} \\
 m: M(-t, \text{refl}_t) \\
 \hline
 x': A, u: \text{Hom}(t, x') \vdash J_M^+ m (x', u): M(x', u)
 \end{array}$$


Coslice Path Induction



$$\begin{array}{c}
 t: A^- \\
 x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type} \\
 m: M(-t, \text{refl}_t) \\
 \hline
 x': A, u: \text{Hom}(t, x') \vdash J_M^+ m (x', u): M(x', u)
 \end{array}$$

$$J_M^+ m (-t, \text{refl}_t) = m$$

Coslice Path Induction




$$\begin{array}{c}
 t: A^- \\
 x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type} \\
 m: M(-t, \text{refl}_t) \\
 \hline
 x': A, u: \text{Hom}(t, x') \vdash J_M^+ m (x', u): M(x', u)
 \end{array}$$

$$J_M^+ m (-t, \text{refl}_t) = m$$

refl_t is the “universal coslice” under t


**This solves the
fundamental problem
of directed TT**

Synthetic category structure of hom-types




$$\frac{p: \text{Hom}(t, t') \quad q: \text{Hom}(t', t'')}{p \cdot q: \text{Hom}(t, t'')}$$

Synthetic category structure of hom-types



$$\frac{p: \text{Hom}(t, t') \quad q: \text{Hom}(\text{---}t', t'')}{p \cdot q: \text{Hom}(t, t'')}$$


Synthetic category structure of hom-types



$$\frac{p: \text{Hom}(t, t') \quad q: \text{Hom}(-t', t'')}{p \cdot q: \text{Hom}(t, t'')}$$

$$x'': A, u: \text{Hom}(-t', x'') \vdash \\ J_{\text{Hom}(t, x'')}^+ p (x'', u): \text{Hom}(t, x'')$$


Synthetic category structure of hom-types



$$\frac{p: \text{Hom}(t, t') \quad q: \text{Hom}(-t', t'')}{p \cdot q: \text{Hom}(t, t'')}$$


$$x'': A, u: \text{Hom}(-t', x'') \vdash \\ J_{\text{Hom}(t, x'')}^+ p (x'', u): \text{Hom}(t, x'')$$

Synthetic category structure of hom-types




$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(t', t)}$$

Synthetic category structure of hom-types




$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(\text{---}t', t)}$$

Synthetic category structure of hom-types



$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(\text{---}t', \text{---}t)}$$


Synthetic category structure of hom-types



$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(\text{---}t', \text{---}t)}$$

$$x' : A, u : \text{Hom}(t, x') \vdash \\ J^+ \text{ refl}_t (x', u) : \text{Hom}(\text{---}x', \text{---}t)$$


Synthetic category structure of hom-types



$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(\text{---}t', \text{---}t)}$$

$$x' : A, u : \text{Hom}(t, x') \vdash \\ J^+ \text{ refl}_t (x', u) : \text{Hom}(\text{---}x', \text{---}t)$$

Synthetic category structure of hom-types



$$\frac{p : \text{Hom}(t, t')}{p^{-1} : \text{Hom}(\text{---}t', \text{---}t)}$$

$$x' : A, u : \text{Hom}(t, x') \vdash \\ J^+ \text{refl}_t (x', u) : \text{Hom}(\text{---}x', \text{---}t)$$

Semantic Proof:


**Symmetry *can't* be
proved in general**

*Category theory is concerned with
universal mapping properties*

*Category theory is concerned with
universal mapping properties*

*Type-theoretic constructs are introduced with
principles of induction*

Coslice Path Induction



$$\begin{array}{c} t: A^- \\ x': A, u: \text{Hom}(t, x') \vdash M(x', u) \text{ type} \\ m: M(-t, \text{refl}_t) \\ \hline x': A, u: \text{Hom}(t, x') \vdash J_M^+ m(x', u): M(x', u) \end{array}$$

$$J_M^+ m(-t, \text{refl}_t) = m$$

refl_t is the “universal coslice” under t

A **coproduct** of $s, t: A^-$ consists of terms

t ●

s ●

A **coproduct** of $s, t: A^-$ consists of terms

- $Q: A^-$

t ●

Q ●

s ●

A **coproduct** of $s, t: A^-$ consists of terms


- $Q: A^-$
- $\iota_1: \text{Hom}(s, -Q)$ and $\iota_2: \text{Hom}(t, -Q)$



A **coproduct** of $s, t: A^-$ consists of terms

- $Q: A^-$
- $\iota_1: \text{Hom}(s, -Q)$ and
 $\iota_2: \text{Hom}(t, -Q)$

such that




$x': A, u: \text{Hom}(s, x'), v: \text{Hom}(t, x') \vdash M$ **type**

A **coproduct** of $s, t: A^-$ consists of terms

- $Q: A^-$
- $\iota_1: \text{Hom}(s, -Q)$ and $\iota_2: \text{Hom}(t, -Q)$

such that

$x': A, u: \text{Hom}(s, x'), v: \text{Hom}(t, x') \vdash M \text{ type}$




A **coproduct** of $s, t: A^-$ consists of terms

- $Q: A^-$
- $\iota_1: \text{Hom}(s, -Q)$ and $\iota_2: \text{Hom}(t, -Q)$

such that

$$\frac{x': A, u: \text{Hom}(s, x'), v: \text{Hom}(t, x') \vdash M \text{ type} \quad m: M(-Q, \iota_1, \iota_2)}{}$$



A **coproduct** of $s, t: A^-$ consists of terms


- $Q: A^-$
- $\iota_1: \text{Hom}(s, -Q)$ and $\iota_2: \text{Hom}(t, -Q)$

such that

$x': A, u: \text{Hom}(s, x'), v: \text{Hom}(t, x') \vdash M \text{ type}$

$m: M(-Q, \iota_1, \iota_2)$


$x', u, v \vdash \text{elim}_M m (x', u, v): M(x', u, v)$



A **coproduct** of $s, t: A^-$ consists of terms

- $Q: A^-$
- $\iota_1: \text{Hom}(s, -Q)$ and $\iota_2: \text{Hom}(t, -Q)$

such that



$x': A, u: \text{Hom}(s, x'), v: \text{Hom}(t, x') \vdash M$ **type**

$m: M(-Q, \iota_1, \iota_2)$

$x', u, v \vdash \text{elim}_M m (x', u, v): M(x', u, v)$

$\text{elim}_M m (-Q, \iota_1, \iota_2) = m$

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

- Define the identity functor on A , $I_A := \lambda x.x$.

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

- Define the identity functor on A , $I_A := \lambda x.x$.
- Given $F: A \rightarrow B$ and $f: \text{Hom}_A(t, t')$, define map $F f: \text{Hom}(F(t), F(t'))$:

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

- Define the identity functor on A , $I_A := \lambda x.x$.
- Given $F: A \rightarrow B$ and $f: \text{Hom}_A(t, t')$, define map $F f: \text{Hom}(-F(-t), F(t'))$:

$$x': A, u: \text{Hom}_A(t, x') \vdash J^+ \text{ refl}_{-F(-t)}: \text{Hom}(-F(-t), F(x'))$$

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

- Define the identity functor on A , $I_A := \lambda x.x$.
- Given $F: A \rightarrow B$ and $f: \text{Hom}_A(t, t')$, define map $F f: \text{Hom}(-F(-t), F(t'))$:

$$x': A, u: \text{Hom}_A(t, x') \vdash J^+ \text{ refl}_{-F(-t)}: \text{Hom}(-F(-t), F(x'))$$

A term $\alpha: \text{Hom}_{A \rightarrow B}(F, G)$ is a natural transformation

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

- Define the identity functor on A , $I_A := \lambda x.x$.
- Given $F: A \rightarrow B$ and $f: \text{Hom}_A(t, t')$, define map $F f: \text{Hom}(-F(-t), F(t'))$:

$$x': A, u: \text{Hom}_A(t, x') \vdash J^+ \text{ refl}_{-F(-t)}: \text{Hom}(-F(-t), F(x'))$$

A term $\alpha: \text{Hom}_{A \rightarrow B}(F, G)$ is a natural transformation

By Coslice Path Induction,

$$\frac{F: (A \rightarrow B)^- \quad G: A \rightarrow B \quad \alpha: \text{Hom}(F, G) \quad t': A}{\alpha @ t': \text{Hom}(-((-F) t'), G(t'))}$$

Synthetic functors and natural transformations

A term $F: A \rightarrow B$ is a functor from A to B

- Define the identity functor on A , $I_A := \lambda x.x$.
- Given $F: A \rightarrow B$ and $f: \text{Hom}_A(t, t')$, define map $F f: \text{Hom}(-F(-t), F(t'))$:

$$x': A, u: \text{Hom}_A(t, x') \vdash J^+ \text{ refl}_{-F(-t)}: \text{Hom}(-F(-t), F(x'))$$

A term $\alpha: \text{Hom}_{A \rightarrow B}(F, G)$ is a natural transformation

By Coslice Path Induction, $\text{refl}@t' := \text{refl}_{-((-F) t')}$ yields

$$\frac{F: (A \rightarrow B)^- \quad G: A \rightarrow B \quad \alpha: \text{Hom}(F, G) \quad t': A}{\alpha@t': \text{Hom}(-((-F) t'), G(t'))}$$

A **left adjoint** of $U: B \rightarrow A$ consists of

Adjoints

A **left adjoint** of $U: B \rightarrow A$ consists of

- $F: A \rightarrow B$

Adjoints

A **left adjoint** of $U: B \rightarrow A$ consists of

- $F: A \rightarrow B$
- $\eta: \text{Hom}(-, I_A, U \circ F)$

Adjoint

A **left adjoint** of $U: B \rightarrow A$ consists of

- $F: A \rightarrow B$
- $\eta: \text{Hom}(-, I_A, U \circ F)$

such that

$$t: A^-$$

Adjoints

A **left adjoint** of $U: B \rightarrow A$ consists of

- $F: A \rightarrow B$
- $\eta: \text{Hom}(-I_A, U \circ F)$

such that

$$\begin{array}{c} t: A^- \\ z': A, u: \text{Hom}(t, U(z')) \vdash M(z', u) \text{ type} \end{array}$$

Adjoints

A **left adjoint** of $U: B \rightarrow A$ consists of

- $F: A \rightarrow B$
- $\eta: \text{Hom}(-I_A, U \circ F)$

such that

$$\begin{array}{c} t: A^- \\ z': A, u: \text{Hom}(t, U(z')) \vdash M(z', u) \text{ type} \\ m: M(F(-t), \eta @ (-t)) \end{array}$$

Adjoint

A **left adjoint** of $U: B \rightarrow A$ consists of

- $F: A \rightarrow B$
- $\eta: \text{Hom}(-I_A, U \circ F)$

such that

$$\frac{\begin{array}{c} t: A^- \\ z': A, u: \text{Hom}(t, U(z')) \vdash M(z', u) \text{ type} \\ m: M(F(-t), \eta @ (-t)) \end{array}}{z': A, u: \text{Hom}(t, U(z')) \vdash \text{elim}_M m (z', u): M(z', u)}$$

$$\text{elim}_M m (F(-t), \eta @ (-t)) = m$$

Apprehended

- Initial and Terminal Objects
- (Co)Products
- Pullbacks and Pushouts
- Left and Right Adjoints
- Applying a natural transformation
- Definition of Yoneda embedding
 - ▶ Exponentials
 - ▶ All limits of shape I

Still at large

- Lambda-abstraction rule for natural transformations
 - ▶ Proof by directed path induction that natural transformations are natural
 - ▶ Internal proof of Yoneda Lemma
 - ▶ (Co)limits in presheaf categories
- Monomorphisms/Epimorphisms (coinductive characterization?)

- [NA25] **Jacob Neumann and Thorsten Altenkirch.**
Synthetic 1-categories in directed type theory.
In Rasmus Ejlers Møgelberg and Benno van den Berg, editors, *30th International Conference on Types for Proofs and Programs (TYPES 2024)*, Leibniz International Proceedings in Informatics (LIPIcs), 2025.
- [Neu25] **Jacob Neumann.**
A Generalized Algebraic Theory of Directed Equality.
PhD thesis, University of Nottingham, 2025.
- [Nor19] **Paige Randall North.**
Towards a directed homotopy type theory.
Electronic Notes in Theoretical Computer Science, 347:223–239, 2019.

- Track polarities
- Limit coercions to closed terms
- Fail to prove symmetry
- Phrase universal mapping properties as principles of induction

Thank you!