

# Linear Regression Review

FW8051 Statistics for Ecologists

Department of Fisheries, Wildlife and Conservation Biology



# Objectives

Review important statistical concepts within the context of linear regression using simulated data:

- Sampling Distributions
- T-tests for regression coefficients
- Confidence intervals
- P-values
- How to check assumptions

# Sustainable trophy hunting of African Lions

Whitman et al. 2014 Nature 428:175-178

Important to know the age of male lions to help manage trophy hunting

- Removing males over 6 has little effect on social structure
- Removing younger males is more disruptive

# Sustainable trophy hunting of African Lions

Whitman et al. 2014 Nature 428:175-178

Important to know the age of male lions to help manage trophy hunting

- Removing males over 6 has little effect on social structure
- Removing younger males is more disruptive

How, from afar, can we tell the age of a lion?

## Its in the Nose!



After about 3 years of age, the fleshy part of the nose begins to freckle or become liver spotted. As the lion ages, more pigmentation appears until the nose is entirely black by about 8 years. A general rule of thumb is that by 6 years noses are >50% black.

Data are contained in abd library of Program R:


```
install.packages("abd") # only if not installed (do once)
```

```
library(abd) # Each time you want to access the data  
data(LionNoses)  
head(LionNoses)
```

	age	proportion.black
1	1.1	0.21
2	1.5	0.14
3	1.9	0.11
4	2.2	0.13
5	2.6	0.12
6	3.2	0.13

# Lion's Noses

```
library(ggplot2)
ggplot(LionNoses, aes(proportion.black, age)) +
  geom_point() +
  geom_smooth(method = "lm", formula = y ~ x) + xlab("Proportion Black")
ylab("Age")
```



# Linear Regression Assumptions

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

Assumptions (HILE Gauss):



# Linear Regression Assumptions

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

Assumptions (HILE Gauss):

- **Homogeneity of variance** (constant scatter about the line);  
 $var(\epsilon_i) = \sigma^2$

# Linear Regression Assumptions

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

Assumptions (HILE Gauss):

- **Homogeneity of variance** (constant scatter about the line);  
 $var(\epsilon_i) = \sigma^2$
- **Independence**:  $Correlation(\epsilon_i, \epsilon_j) = 0$

# Linear Regression Assumptions

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

Assumptions (HILE Gauss):

- **Homogeneity of variance** (constant scatter about the line);  
 $\text{var}(\epsilon_i) = \sigma^2$
- **Independence**:  $\text{Correlation}(\epsilon_i, \epsilon_j) = 0$
- **Linearity**:  $E[Y_i | X_i] = \beta_0 + X_i \beta_1$

# Linear Regression Assumptions

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$
$$\epsilon_i \sim N(0, \sigma^2)$$

Assumptions (HILE Gauss):

- **Homogeneity of variance** (constant scatter about the line);  
 $\text{var}(\epsilon_i) = \sigma^2$
- **Independence**:  $\text{Correlation}(\epsilon_i, \epsilon_j) = 0$
- **Linearity**:  $E[Y_i | X_i] = \beta_0 + X_i \beta_1$
- **Existence** (we observe random variables that have finite variance; we won't worry about this one)

# Linear Regression Assumptions

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$


$$\epsilon_i \sim N(0, \sigma^2)$$

Assumptions (HILE Gauss):

- **Homogeneity of variance** (constant scatter about the line);  
 $var(\epsilon_i) = \sigma^2$
- **Independence**:  $Correlation(\epsilon_i, \epsilon_j) = 0$
- **Linearity**:  $E[Y_i | X_i] = \beta_0 + X_i \beta_1$
- **Existence** (we observe random variables that have finite variance; we won't worry about this one)
- **Gauss**:  $\epsilon_i$  come from a Normal (Gaussian) distribution


# Regression Assumptions

- We specify a probability distribution for  $Y_i|X_i \sim N(\mu_i, \sigma^2)$
- We have a model for the mean  $\mu_i = \beta_0 + \beta_1 X_i$



# Regression Assumptions

- We specify a probability distribution for  $Y_i|X_i \sim N(\mu_i, \sigma^2)$
- We have a model for the mean  $\mu_i = \beta_0 + \beta_1 X_i$



- How are these assumptions reflected in the figure? How can we evaluate the assumptions with our data?

# Residuals Versus Fitted




Figure 10.2

© John Wiley & Sons, Inc. All rights reserved.




# Graphical Check

```
lm.nose <- lm(age ~ proportion.black, data = LionNoses)
performance::check_model(lm.nose,
  check = c("linearity", "homogeneity", "qq", "normality"))
```


Linearity

Reference line should be flat and horizontal




Homogeneity of Variance

Reference line should be flat and horizontal




Normality of Residuals

Dots should fall along the line



Normality of Residuals

Distribution should be close to the normal curve



## Interpretation: Intercept, Slope, t-tests and p-values, Residual Standard Error ( $\hat{\sigma}$ ), $R^2$

```
summary(lm.nose)
```

Call:

```
lm(formula = age ~ proportion.black, data = LionNoses)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.5449	-1.1117	-0.5285	0.9635	4.3421

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.8790	0.5688	1.545	0.133
proportion.black	10.6471	1.5095	7.053	7.68e-08 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.669 on 30 degrees of freedom

Multiple R-squared: 0.6238, Adjusted R-squared: 0.6113

F-statistic: 49.75 on 1 and 30 DF, p-value: 7.677e-08

## Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65\text{Proportion.black}$$

Intercept:

## Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65\text{Proportion.black}$$

Intercept: Estimate of the average age of lions that have no black pigmentation on their noses ( $E[Y|X = 0]$ ).

# Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65\text{Proportion.black}$$

Intercept: Estimate of the average age of lions that have no black pigmentation on their noses ( $E[Y|X = 0]$ ).

Slope =

# Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65\text{Proportion.black}$$

Intercept: Estimate of the average age of lions that have no black pigmentation on their noses ( $E[Y|X = 0]$ ).

Slope = Predicted change in age per unit increase in proportion black pigmentation

# Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65 \text{Proportion.black}$$

Intercept: Estimate of the average age of lions that have no black pigmentation on their noses ( $E[Y|X = 0]$ ).

Slope = Predicted change in age per unit increase in proportion black pigmentation

$$10.65 = \frac{\Delta \text{age}}{\Delta \text{Proportion.Black}}$$

But, proportion black  $< 1$  for all lions.

# Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65\text{Proportion.black}$$

Intercept: Estimate of the average age of lions that have no black pigmentation on their noses ( $E[Y|X = 0]$ ).

Slope = Predicted change in age per unit increase in proportion black pigmentation

$$10.65 = \frac{\Delta \text{age}}{\Delta \text{Proportion.Black}}$$

But, proportion black < 1 for all lions.

$$? = \frac{\Delta \text{age}}{\Delta 0.1 \text{Proportion.Black}} =$$



# Interpretation: Intercept and Slope

$$\widehat{age} = 0.879 + 10.65\text{Proportion.black}$$

Intercept: Estimate of the average age of lions that have no black pigmentation on their noses ( $E[Y|X = 0]$ ).

Slope = Predicted change in age per unit increase in proportion black pigmentation

$$10.65 = \frac{\Delta \text{age}}{\Delta \text{Proportion.Black}}$$

But, proportion black  $< 1$  for all lions.

$$? = \frac{\Delta \text{age}}{\Delta 0.1 \text{Proportion.Black}} = 1.065.$$

$H_0 : \beta_1 = 0$  vs.  $H_A : \beta_1 \neq 0$ ?

```
summary(lm.nose)
```

Call:

```
lm(formula = age ~ proportion.black, data = LionNoses)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.5449	-1.1117	-0.5285	0.9635	4.3421

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.8790	0.5688	1.545	0.133
proportion.black	10.6471	1.5095	7.053	7.68e-08 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.669 on 30 degrees of freedom

Multiple R-squared: 0.6238, Adjusted R-squared: 0.6113

F-statistic: 49.75 on 1 and 30 DF, p-value: 7.677e-08

## SE, t-value, p-value

Need to understand the concept of a **Sampling Distribution** of a statistic.

## SE, t-value, p-value

Need to understand the concept of a **Sampling Distribution** of a statistic.

A **sampling distribution** is the distribution of sample statistics computed for different samples of the same size from the same population.

## SE, t-value, p-value

Need to understand the concept of a **Sampling Distribution** of a statistic.

A **sampling distribution** is the distribution of sample statistics computed for different samples of the same size from the same population.

A sampling distribution shows us how the sample statistic varies from sample to sample

## SE, t-value, p-value

Need to understand the concept of a **Sampling Distribution** of a statistic.

A **sampling distribution** is the distribution of sample statistics computed for different samples of the same size from the same population.

A sampling distribution shows us how the sample statistic varies from sample to sample

See: AP stats guy videos in Ch. 1 of the textbook

SE  $\hat{\beta}$

Think of many repetitions of:

- Collecting a new data set (of the same size)

SE  $\hat{\beta}$

Think of many repetitions of:

- Collecting a new data set (of the same size)
- Fitting the same regression model



SE  $\hat{\beta}$

Think of many repetitions of:

- Collecting a new data set (of the same size)
- Fitting the same regression model
- Calculating  $\hat{\beta}$

# SE $\hat{\beta}$

Think of many repetitions of:

- Collecting a new data set (of the same size)
- Fitting the same regression model
- Calculating  $\hat{\beta}$

**Sampling distribution** of  $\hat{\beta}$  is the distribution of  $\hat{\beta}$  values across the different samples.

# SE $\hat{\beta}$

Think of many repetitions of:

- Collecting a new data set (of the same size)
- Fitting the same regression model
- Calculating  $\hat{\beta}$

**Sampling distribution** of  $\hat{\beta}$  is the distribution of  $\hat{\beta}$  values across the different samples.

**SE** = standard deviation of the sampling distribution!

# SE $\hat{\beta}$

Think of many repetitions of:

- Collecting a new data set (of the same size)
- Fitting the same regression model
- Calculating  $\hat{\beta}$

**Sampling distribution** of  $\hat{\beta}$  is the distribution of  $\hat{\beta}$  values across the different samples.

**SE** = standard deviation of the sampling distribution!

Lets explore through simulation!

# Simulation

Lets first generate a single data set consistent with our fitted model using the following code.

```
# Sample size of simulated observations
n<-32

# Use the observed proportion.black to simulate obs.
p.black<-LionNoses$proportion.black

# Use the estimated parameters to simulate data.
# - can get these from the regression output
# sigma<-summary(lm.nose)$sigma # residual variation about the line
# betas<-coef(lm.nose) # Regression coefficients
sigma<-1.67 # residual variation
betas<-c(0.88, 10.65) #betas

# Create random errors (epsilons) and random responses
epsilon<-rnorm(n,0, sigma) # Errors
y<-betas[1] + p.black*betas[2] + epsilon # Response
```

# Linear regression using lm function

```
lmfit<-lm(y~p.black)
summary(lmfit)
```

Call:

```
lm(formula = y ~ p.black)
```

Residuals:

Min	1Q	Median	3Q	Max
-3.2401	-0.8812	-0.3871	0.9053	3.2192

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	1.7028	0.5627	3.026	0.00505 **
p.black	8.9392	1.4934	5.986	1.45e-06 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.651 on 30 degrees of freedom

Multiple R-squared: 0.5443, Adjusted R-squared: 0.5291

F-statistic: 35.83 on 1 and 30 DF, p-value: 1.45e-06

# Sampling Distribution

Use a for loop to:

- Generate 5000 data sets using the same code
- Fit a linear regression model to each data set
- For each fit, store  $\hat{\beta}$

In-class exercise

# Sampling distribution

When conducting hypothesis tests or constructing confidence intervals, we will work with the distribution of:

$$\frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$



## Sampling distribution of the t-statistic

$$\frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$

# Sampling distribution of the t-statistic

$$\frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$

Think of many repetitions of:

- Collecting a new data set (of the same size)
- Fitting the regression model
- Calculating:  $t = \frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)}$

A histogram of the different  $t$  values should be well described by a Student's  $t$ -distribution with  $n - 2$  degrees of freedom.

# Sampling Distribution

Use a for loop to:

- Generate 5000 data sets using the same code
- Fit a linear regression model to each data set
- For each fit, store  $\hat{\beta}$  and calculate:  $t = \frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)}$

# Sampling Distribution

Use a for loop to:

- Generate 5000 data sets using the same code
- Fit a linear regression model to each data set
- For each fit, store  $\hat{\beta}$  and calculate:  $t = \frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)}$

Helpful hints:

- $\beta_1$  = true value used to simulate the data,  
`coef(lm.nose)[2] = 10.6471`
- $\hat{\beta}_1$  is extracted using: `coef(lmfit)[2]`
- $\widehat{SE}(\hat{\beta}_1)$  is extracted using `sqrt(vcov(lm.temp)[2,2])`

In-class exercise


My code

# My code

```
nsims<-5000 # number of simulations
tsamp.dist<-matrix(NA, nsims,1) # matrix to hold results
for(i in 1:5000){
  epsilon<-rnorm(n,0,sigma) # Errors
  y<-betas[1] + betas[2]*p.black + epsilon # Response
  lm.temp<-lm(y~p.black) # lm

  # Here is our statistic, calculated for each sample
  tsamp.dist[i]<-(coef(lm.temp)[2]-betas[2])/sqrt(vcov(lm.temp)[2,2])
}
# Plot results
hist(tsamp.dist, xlab="",
     main=expression(t==frac(hat(beta)-beta, se(hat(beta)))), freq=FALSE)
tvalues<-seq(-3,3, length=1000) # xvalues to evaluate t-distribution
lines(tvalues,dt(tvalues, df=30)) # overlay t-distribution
```

# Sampling distribution of t-statistic



# Confidence Interval



# Confidence Interval

A **confidence interval** for a parameter is an interval computed from sample data by a method that will capture the parameter for a specified proportion of all samples

# Confidence Interval

A **confidence interval** for a parameter is an interval computed from sample data by a method that will capture the parameter for a specified proportion of all samples

- The success rate (proportion of all samples whose intervals contain the parameter) is known as the **confidence level** or **coverage rate**

# Confidence Interval

A **confidence interval** for a parameter is an interval computed from sample data by a method that will capture the parameter for a specified proportion of all samples

- The success rate (proportion of all samples whose intervals contain the parameter) is known as the **confidence level** or **coverage rate**
- A 95% confidence interval should contain the true parameter for 95% of all samples


# Confidence Interval

A **confidence interval** for a parameter is an interval computed from sample data by a method that will capture the parameter for a specified proportion of all samples

- The success rate (proportion of all samples whose intervals contain the parameter) is known as the **confidence level** or **coverage rate**
- A 95% confidence interval should contain the true parameter for 95% of all samples
- The parameter is *fixed*, but the endpoints of the interval are *random*

# Confidence Interval

$$\frac{\hat{\beta}_1 - \beta_1}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$



Note:

$$t_{0.025, n-2} = \text{qt}(p=0.025, \text{df}=30) = -2.04$$

$$t_{0.975, n-2} = \text{qt}(p=0.975, \text{df}=30) = 2.04$$

# Confidence Interval

$$P(t_{0.025,n-2} < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < t_{0.975,n-2}) = 0.95$$

# Confidence Interval

$$P(t_{0.025,n-2} < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < t_{0.975,n-2}) = 0.95$$

$$P(-2.04 < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < 2.04) = 0.95$$

# Confidence Interval

$$P(t_{0.025, n-2} < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < t_{0.975, n-2}) = 0.95$$

$$P(-2.04 < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < 2.04) = 0.95$$

$$P(-2.04\widehat{SE}(\hat{\beta}) < \hat{\beta} - \beta < 2.04\widehat{SE}(\hat{\beta})) = 0.95$$



# Confidence Interval

$$P(t_{0.025, n-2} < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < t_{0.975, n-2}) = 0.95$$

$$P(-2.04 < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < 2.04) = 0.95$$

$$P(-2.04\widehat{SE}(\hat{\beta}) < \hat{\beta} - \beta < 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

$$P(-\hat{\beta} - 2.04\widehat{SE}(\hat{\beta}) < -\beta < -\hat{\beta} + 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

# Confidence Interval

$$P(t_{0.025, n-2} < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < t_{0.975, n-2}) = 0.95$$

$$P(-2.04 < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < 2.04) = 0.95$$

$$P(-2.04\widehat{SE}(\hat{\beta}) < \hat{\beta} - \beta < 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

$$P(-\hat{\beta} - 2.04\widehat{SE}(\hat{\beta}) < -\beta < -\hat{\beta} + 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

$$P(\hat{\beta} + 2.04\widehat{SE}(\hat{\beta}) > \beta > \hat{\beta} - 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

## Confidence Interval

$$P(t_{0.025, n-2} < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < t_{0.975, n-2}) = 0.95$$

$$P(-2.04 < \frac{\hat{\beta} - \beta}{\widehat{SE}(\hat{\beta})} < 2.04) = 0.95$$

$$P(-2.04\widehat{SE}(\hat{\beta}) < \hat{\beta} - \beta < 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

$$P(-\hat{\beta} - 2.04\widehat{SE}(\hat{\beta}) < -\beta < -\hat{\beta} + 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

$$P(\hat{\beta} + 2.04\widehat{SE}(\hat{\beta}) > \beta > \hat{\beta} - 2.04\widehat{SE}(\hat{\beta})) = 0.95$$

So, take  $(\hat{\beta} - 2.04\widehat{SE}(\hat{\beta}), \hat{\beta} + 2.04\widehat{SE}(\hat{\beta}))$  as the the 95% confidence interval.

## confint function

```
confint(lm.nose)
```

	2.5 %	97.5 %
(Intercept)	-0.2826733	2.040686
proportion.black	7.5643082	13.729931

What is wrong with the following interpretation?

$$P(7.56 \leq \beta \leq 13.72) = 0.95$$

What is wrong with the following interpretation?

$$P(7.56 \leq \beta \leq 13.72) = 0.95$$

- $\beta$  is either in this particular interval ( $P = 1$ ) or it is not ( $P = 0$ )

What is wrong with the following interpretation?

$$P(7.56 \leq \beta \leq 13.72) = 0.95$$

- $\beta$  is either in this particular interval ( $P = 1$ ) or it is not ( $P = 0$ )
- $P(L \leq \beta \leq U) = 0.95$ , where  $L$  and  $U$  are random variables that determine the upper and lower limits of the 95% confidence interval

What is wrong with the following interpretation?

$$P(7.56 \leq \beta \leq 13.72) = 0.95$$

- $\beta$  is either in this particular interval ( $P = 1$ ) or it is not ( $P = 0$ )
- $P(L \leq \beta \leq U) = 0.95$ , where  $L$  and  $U$  are random variables that determine the upper and lower limits of the 95% confidence interval

We are 95% sure that the true slope (relating proportion of nose that is black and age) falls between 7.56 and 13.73.



# Explore CIs through simulation

Simulate another 5000 data sets in R:

Determine 95% confidence limits for each data set and examine whether or not the CI contains the true  $\beta$ .

## My Code

# My Code


```
nsims<-5000 # number of simulations
Limits<-matrix(NA,nsims,2) # to hold results
beta.hats<-matrix(NA,nsims,1) # to hold estimates
for(i in 1:nsims){
  epsilon<-rnorm(n,0, sigma) # Errors
  y<-betas[1] + betas[2]*p.black + epsilon # Response
  lm.temp<-lm(y~p.black)

  # Beta.hat & Confidence limits
  beta.hats[i]<-coef(lm.temp)[2]
  Limits[i,]<-confint(lm.temp)[2,]
}
# True parameter in interval?
I.in<-betas[2]>=Limits[,1] & betas[2] <= Limits[,2]

# Coverage
sum(I.in)/nsims
```

```
[1] 0.9444
```

# First 100 simulations



```
summary(lm.nose)
```

Call:

```
lm(formula = age ~ proportion.black, data = LionNoses)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.5449	-1.1117	-0.5285	0.9635	4.3421

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.8790	0.5688	1.545	0.133
proportion.black	10.6471	1.5095	7.053	7.68e-08 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.669 on 30 degrees of freedom

Multiple R-squared: 0.6238, Adjusted R-squared: 0.6113

F-statistic: 49.75 on 1 and 30 DF, p-value: 7.677e-08

## P-values

If the null hypothesis,  $H_0 : \beta_1 = 0$ , is true, then:

$$t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$

## P-values

If the null hypothesis,  $H_0 : \beta_1 = 0$ , is true, then:

$$t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$

Is the value we get for  $t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} = 7.053$  consistent with  $H_0 : \beta_1 = 0$ ?

# P-values

If the null hypothesis,  $H_0 : \beta_1 = 0$ , is true, then:

$$t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$$


Is the value we get for  $t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} = 7.053$  consistent with  $H_0 : \beta_1 = 0$ ?

- Overlay  $t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} = 7.053$  on a  $t_{n-2}$  distribution
- Determine the probability of getting a t-statistic as or more extreme as the one we observed.



# Hypothesis test

- t-distribution tells about the sampling distribution of  $t = \frac{\hat{\beta}_1 - 0}{\widehat{SE}(\hat{\beta}_1)} \sim t_{n-2}$  when the null hypothesis is true
- our t-statistic falls in the tail of this distribution (so, the Null hypothesis is unlikely to be true!)



$R^2$ 

```
summary(lm.nose)
```

Call:

```
lm(formula = age ~ proportion.black, data = LionNoses)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.5449	-1.1117	-0.5285	0.9635	4.3421

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.8790	0.5688	1.545	0.133
proportion.black	10.6471	1.5095	7.053	7.68e-08 ***

---



Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.669 on 30 degrees of freedom



Multiple R-squared: 0.6238, Adjusted R-squared: 0.6113

F-statistic: 49.75 on 1 and 30 DF, p-value: 7.677e-08

# Sum of Squares





# Sum of Squares



$$\text{SST (Total sum of squares)} = \sum_i^n (Y_i - \bar{Y})^2$$



# Sum of Squares



$$\text{SST (Total sum of squares)} = \sum_i^n (Y_i - \bar{Y})^2$$

$$\text{SSE (Sum of Squares Error)} = \sum_i^n (Y_i - \hat{Y})^2$$

# Sum of Squares




$$\text{SST (Total sum of squares)} = \sum_i^n (Y_i - \bar{Y})^2$$

$$\text{SSE (Sum of Squares Error)} = \sum_i^n (Y_i - \hat{Y})^2$$

$$\text{SSR (sum of squares regression)} = \text{SST} - \text{SSE} = \sum_i^n (\hat{Y}_i - \bar{Y})^2$$

# Sum of Squares



$$SST \text{ (Total sum of squares)} = \sum_i^n (Y_i - \bar{Y})^2$$

$$SSE \text{ (Sum of Squares Error)} = \sum_i^n (Y_i - \hat{Y})^2$$

$$SSR \text{ (sum of squares regression)} = SST - SSE = \sum_i^n (\hat{Y}_i - \bar{Y})^2$$

$$R^2 = \frac{SST - SSE}{SST} = \frac{SSR}{SST} = \text{proportion of the variation explained by the linear model.}$$

Residual standard error =  $\hat{\sigma}$

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

- $\sigma$  describes the amount of variability about the regression line



Residual standard error =  $\hat{\sigma}$

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

- $\sigma$  describes the amount of variability about the regression line

$$\hat{\sigma} = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n-p}} = \sqrt{\frac{SSE}{n-p}} = \sqrt{MSE}$$

## Residual standard error = $\hat{\sigma}$

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

- $\sigma$  describes the amount of variability about the regression line

$$\hat{\sigma} = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n-p}} = \sqrt{\frac{SSE}{n-p}} = \sqrt{MSE}$$

- Listed as Residual Standard Error in R output from `summary` function

## Residual standard error = $\hat{\sigma}$

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, \sigma^2)$$

- $\sigma$  describes the amount of variability about the regression line

$$\hat{\sigma} = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n-p}} = \sqrt{\frac{SSE}{n-p}} = \sqrt{MSE}$$

- Listed as Residual Standard Error in R output from `summary` function
- $n - p$  since we lose one degree of freedom for each parameter we estimate

# Residual Standard Error

```
summary(lm.nose)
```

Call:

```
lm(formula = age ~ proportion.black, data = LionNoses)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.5449	-1.1117	-0.5285	0.9635	4.3421

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.8790	0.5688	1.545	0.133
proportion.black	10.6471	1.5095	7.053	7.68e-08 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.669 on 30 degrees of freedom

Multiple R-squared: 0.6238, Adjusted R-squared: 0.6113

F-statistic: 49.75 on 1 and 30 DF, p-value: 7.677e-08

# Residual Standard Error

```
summary(lm.nose)
```

Call:

```
lm(formula = age ~ proportion.black, data = LionNoses)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.5449	-1.1117	-0.5285	0.9635	4.3421

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	0.8790	0.5688	1.545	0.133
proportion.black	10.6471	1.5095	7.053	7.68e-08 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.669 on 30 degrees of freedom

Multiple R-squared: 0.6238, Adjusted R-squared: 0.6113

F-statistic: 49.75 on 1 and 30 DF, p-value: 7.677e-08

We expect 95% of the observations to fall within  $2 \times 1.669$  of the regression line.