

2007 AP® CALCULUS AB FREE-RESPONSE QUESTIONS**CALCULUS AB
SECTION II, Part B****Time—45 minutes****Number of problems—3****No calculator is allowed for these problems.**

-
4. A particle moves along the x -axis with position at time t given by $x(t) = e^{-t} \sin t$ for $0 \leq t \leq 2\pi$.
- (a) Find the time t at which the particle is farthest to the left. Justify your answer.
- (b) Find the value of the constant A for which $x(t)$ satisfies the equation $Ax''(t) + x'(t) + x(t) = 0$ for $0 < t < 2\pi$.
-

t (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

5. The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function r of time t , where t is measured in minutes. For $0 < t < 12$, the graph of r is concave down. The table above gives selected values of the rate of change, $r'(t)$, of the radius of the balloon over the time interval $0 \leq t \leq 12$. The radius of the balloon is 30 feet when $t = 5$. (Note: The volume of a sphere of radius r is given by $V = \frac{4}{3}\pi r^3$.)
- (a) Estimate the radius of the balloon when $t = 5.4$ using the tangent line approximation at $t = 5$. Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when $t = 5$. Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate $\int_0^{12} r'(t) dt$. Using correct units, explain the meaning of $\int_0^{12} r'(t) dt$ in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than $\int_0^{12} r'(t) dt$? Give a reason for your answer.
-

WRITE ALL WORK IN THE PINK EXAM BOOKLET.