

Mathematics

Part I: Calculus

Joanna Franaszek

spring 2019/2020

Warsaw School of Economics

Organisation

Important contacts

- joint course by Maria Ekes and me
- name: Joanna Franaszek
- mail: jfrana@sgh.waw.pl,
- webpage (slides, notes, announcements etc.):
<https://jfranaszek.github.io/>
- office hours: Monday 14:20, room TBA (*please e-mail me in advance*)

Grading

- 0-30 points: *average* of two written tests (mid-term and end-term, roughly 1/2 of the material)
- 0-30 points: final written exam (full material)
- 0-5 points: activity in exercise classes

score	grade
0-30	2
31-36	3
37-42	3.5
43-48	4
49-54	4.5
55+	5

Textbooks and other helpful resources

- official **e-book**
- WolframAlpha (desktop App or *<https://www.wolframalpha.com/>*
 - some **tutorials** are available
- Stewart James: Calculus Early Transcendentals, 2011, Brooks/Cole, Belmont CA,USA;
- Howard Anton, Chris Rorres: Elementary Linear Algebra with Supplemental Applications, 2010, Clarence Center Inc, Denver MA.

Sequences and limits

Key points

- definition of a sequence; arithmetic sequence; geometric sequence;
- bounded and monotone sequences;
- definition of a limit; simple arithmetic rules;
- squeeze theorem
- conditions for convergence;
- indeterminate forms

$$\frac{0}{0}, \frac{\infty}{\infty}, +\infty - \infty, 0 \cdot \infty, 1^{\infty}, 0^0, \infty^0$$

- the magical number **e**

Sequence

Definition (Sequence)

A sequence is a function $a : \mathbb{N} \rightarrow \mathbb{R}$, where \mathbb{N} is the set of natural numbers, and \mathbb{R} is the set of real numbers. The value $a(n) = a_n$ is called the n -th term of the sequence.

Notation: a_n is a single number, while $(a_n)_{n=1}^{\infty}$ or $(a_n)_{n=1}^{\infty}$ or simply $(a_n)_n$ or $\{a_n\}_n$ denote a sequence.

Special sequences

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall n \geq 1$ is called an **arithmetic sequence** with a common difference r .

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall_{n \geq 1}$ is called an **arithmetic sequence** with a common difference r .
- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n \cdot q \quad \forall_{n \geq 1}$ is called a **geometric sequence** with a common ratio r .

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall_{n \geq 1}$ is called an **arithmetic sequence** with a common difference r .
- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n \cdot q \quad \forall_{n \geq 1}$ is called a **geometric sequence** with a common ratio r .
- sequence is **(weakly) increasing** if $a_{n+1} \geq a_n \quad \forall_n$

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall n \geq 1$ is called an **arithmetic sequence** with a common difference r .
- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n \cdot q \quad \forall n \geq 1$ is called a **geometric sequence** with a common ratio r .
- sequence is **(weakly) increasing** if $a_{n+1} \geq a_n \quad \forall n$
- A sequence is **(weakly) decreasing** if $a_{n+1} \leq a_n \quad \forall n$

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall_{n \geq 1}$ is called an **arithmetic sequence** with a common difference r .
- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n \cdot q \quad \forall_{n \geq 1}$ is called a **geometric sequence** with a common ratio r .
- sequence is **(weakly) increasing** if $a_{n+1} \geq a_n \quad \forall_n$
- A sequence is **(weakly) decreasing** if $a_{n+1} \leq a_n \quad \forall_n$
- A sequence is **bounded from below** if $\exists m \quad \forall_n \quad a_n \geq m$

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall n \geq 1$ is called an **arithmetic sequence** with a common difference r .
- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n \cdot q \quad \forall n \geq 1$ is called a **geometric sequence** with a common ratio r .
- sequence is **(weakly) increasing** if $a_{n+1} \geq a_n \quad \forall n$
- A sequence is **(weakly) decreasing** if $a_{n+1} \leq a_n \quad \forall n$
- A sequence is **bounded from below** if $\exists m \quad \forall n \quad a_n \geq m$
- A sequence is **bounded from above** if $\exists M \quad \forall n \quad a_n \leq M$

Special sequences

- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n + r \quad \forall n \geq 1$ is called an **arithmetic sequence** with a common difference r .
- A sequence $(a_n)_n$ that satisfies: $a_{n+1} = a_n \cdot q \quad \forall n \geq 1$ is called a **geometric sequence** with a common ratio r .
- sequence is **(weakly) increasing** if $a_{n+1} \geq a_n \quad \forall n$
- A sequence is **(weakly) decreasing** if $a_{n+1} \leq a_n \quad \forall n$
- A sequence is **bounded from below** if $\exists m \quad \forall n \quad a_n \geq m$
- A sequence is **bounded from above** if $\exists M \quad \forall n \quad a_n \leq M$
- A sequence is **bounded** if it is bounded from above and below

Limit of a sequence

Definition (Limit)

A number $g \in \mathbb{R}$ is a limit of a sequence $(a_n)_n$ if:

$$\forall \epsilon > 0 \exists N \forall n > N |a_n - g| < \epsilon.$$

If such a number exists, we say the sequence **converges** to g .
Otherwise, the sequence is **divergent**.

Lemma

If $(a_n)_n$ has a limit, it is unique.

"Limits" in $+\infty$ or ∞

Definition (Improper limit)

A sequence $(a_n)_n$ has an improper limit in $+\infty$ ($-\infty$) if:

$$\forall M \exists N_M \forall n > N_M \quad a_n > M \quad (a_n < M).$$

We say the sequence **diverges to infinity (minus infinity)** and denote it by $\lim_{n \rightarrow \infty} a_n = +\infty$ ($\lim_{n \rightarrow \infty} a_n = -\infty$) or, shorter $a_n \rightarrow \pm\infty$

Basic properties of finite limits

Assume $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$ for $a, b, \in \mathbb{R}$:

Basic properties of finite limits

Assume $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$ for $a, b, \in \mathbb{R}$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = (\lim_{n \rightarrow \infty} a_n) \pm (\lim_{n \rightarrow \infty} b_n) = a \pm b$

Basic properties of finite limits

Assume $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$ for $a, b, \in \mathbb{R}$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = (\lim_{n \rightarrow \infty} a_n) \pm (\lim_{n \rightarrow \infty} b_n) = a \pm b$
- $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b$

Basic properties of finite limits

Assume $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$ for $a, b, \in \mathbb{R}$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = (\lim_{n \rightarrow \infty} a_n) \pm (\lim_{n \rightarrow \infty} b_n) = a \pm b$
- $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b$
- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$ if only both sides well-defined i.e. $b_n, b \neq 0$

Basic properties of finite limits

Assume $\lim_{n \rightarrow \infty} a_n = a$ and $\lim_{n \rightarrow \infty} b_n = b$ for $a, b, \in \mathbb{R}$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = (\lim_{n \rightarrow \infty} a_n) \pm (\lim_{n \rightarrow \infty} b_n) = a \pm b$
- $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = a \cdot b$
- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$ if only both sides well-defined i.e. $b_n, b \neq 0$
- $\lim_{n \rightarrow \infty} a_n^{b_n} = a^b$, if only both sides well-defined (note: 0^0 **not** well-defined)

Basic properties of limits with $\pm\infty$

Assume $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$ and $\lim_{n \rightarrow \infty} b_n = +\infty$:

Basic properties of limits with $\pm\infty$

Assume $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$ and $\lim_{n \rightarrow \infty} b_n = +\infty$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = \pm\infty$

Basic properties of limits with $\pm\infty$

Assume $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$ and $\lim_{n \rightarrow \infty} b_n = +\infty$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = \pm\infty$
- $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = \begin{cases} +\infty & \text{if } a > 0 \\ -\infty & \text{if } a < 0 \\ \text{indeterminate} & \text{if } a = 0 \end{cases}$

Basic properties of limits with $\pm\infty$

Assume $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$ and $\lim_{n \rightarrow \infty} b_n = +\infty$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = \pm\infty$
- $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = \begin{cases} +\infty & \text{if } a > 0 \\ -\infty & \text{if } a < 0 \\ \text{indeterminate} & \text{if } a = 0 \end{cases}$
- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$

Basic properties of limits with $\pm\infty$

Assume $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$ and $\lim_{n \rightarrow \infty} b_n = +\infty$:

$$\bullet \lim_{n \rightarrow \infty} (a_n \pm b_n) = \pm\infty$$

$$\bullet \lim_{n \rightarrow \infty} (a_n \cdot b_n) = \begin{cases} +\infty & \text{if } a > 0 \\ -\infty & \text{if } a < 0 \\ \text{indeterminate} & \text{if } a = 0 \end{cases}$$

$$\bullet \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$$

$$\bullet \lim_{n \rightarrow \infty} a_n^{b_n} = \begin{cases} +\infty & \text{if } a > 1 \\ 0 & \text{if } |a| < 1 \\ \text{does not exist} & \text{if } a < -1 \\ \text{indeterminate} & \text{if } a = 1 \end{cases}$$

Basic properties of limits with $\pm\infty$

Assume $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$ and $\lim_{n \rightarrow \infty} b_n = +\infty$:

- $\lim_{n \rightarrow \infty} (a_n \pm b_n) = \pm\infty$

- $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = \begin{cases} +\infty & \text{if } a > 0 \\ -\infty & \text{if } a < 0 \\ \text{indeterminate} & \text{if } a = 0 \end{cases}$

- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$

- $\lim_{n \rightarrow \infty} a_n^{b_n} = \begin{cases} +\infty & \text{if } a > 1 \\ 0 & \text{if } |a| < 1 \\ \text{does not exist} & \text{if } a < -1 \\ \text{indeterminate} & \text{if } a = 1 \end{cases}$

- Note: rules for $b_n \rightarrow -\infty$ can be derived using last two slides:

When simple rules do not work

- when trying to apply arithmetic rules, we can calculate well-defined limits...

When simple rules do not work

- when trying to apply arithmetic rules, we can calculate well-defined limits...
- ...but sometimes we encounter ill-defined statements:

$$\frac{0}{0}, \frac{\infty}{\infty}, +\infty - \infty, 0 \cdot \infty, 1^\infty, 0^0, \infty^0$$

When simple rules do not work

- when trying to apply arithmetic rules, we can calculate well-defined limits...
- ...but sometimes we encounter ill-defined statements:

$$\frac{0}{0}, \frac{\infty}{\infty}, +\infty - \infty, 0 \cdot \infty, 1^\infty, 0^0, \infty^0$$

- **indeterminate forms**

When simple rules do not work

- when trying to apply arithmetic rules, we can calculate well-defined limits...
- ...but sometimes we encounter ill-defined statements:

$$\frac{0}{0}, \frac{\infty}{\infty}, +\infty - \infty, 0 \cdot \infty, 1^\infty, 0^0, \infty^0$$

- **indeterminate forms**
- important note: encountering indeterminate form does **not** necessarily mean the limit does not exist; it means we have to work harder to find it!

When simple rules do not work

- when trying to apply arithmetic rules, we can calculate well-defined limits...
- ...but sometimes we encounter ill-defined statements:

$$\frac{0}{0}, \frac{\infty}{\infty}, +\infty - \infty, 0 \cdot \infty, 1^{\infty}, 0^0, \infty^0$$

- **indeterminate forms**
- important note: encountering indeterminate form does **not** necessarily mean the limit does not exist; it means we have to work harder to find it!
- important note 2: 'true' limits of statements in indeterminate form could be anything: a 'nice' number, zero, $-\infty$ etc.

"Classic" limits to remember

- exponential function diverges quicker than polynomial:

$$\lim_{n \rightarrow \infty} \frac{n^k}{2^n} = 0 \text{ for any } k > 0$$

(also works if we replace 2 with any number $a > 1$)

"Classic" limits to remember

- exponential function diverges quicker than polynomial:

$$\lim_{n \rightarrow \infty} \frac{n^k}{2^n} = 0 \text{ for any } k > 0$$

(also works if we replace 2 with any number $a > 1$)

- factorial converges quicker than polynomial and exponential:

$$\lim_{n \rightarrow \infty} \frac{n^k}{n!} = 0 \text{ and } \lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$$

"Classic" limits to remember

- exponential function diverges quicker than polynomial:

$$\lim_{n \rightarrow \infty} \frac{n^k}{2^n} = 0 \text{ for any } k > 0$$

(also works if we replace 2 with any number $a > 1$)

- factorial converges quicker than polynomial and exponential:

$$\lim_{n \rightarrow \infty} \frac{n^k}{n!} = 0 \text{ and } \lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$$

- for $a > 0$:

$$\lim_{n \rightarrow \infty} \sqrt[n]{a} = \lim_{n \rightarrow \infty} a^{\frac{1}{n}} = 1$$

"Classic" limits to remember

- exponential function diverges quicker than polynomial:

$$\lim_{n \rightarrow \infty} \frac{n^k}{2^n} = 0 \text{ for any } k > 0$$

(also works if we replace 2 with any number $a > 1$)

- factorial converges quicker than polynomial and exponential:

$$\lim_{n \rightarrow \infty} \frac{n^k}{n!} = 0 \text{ and } \lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$$

- for $a > 0$:

$$\lim_{n \rightarrow \infty} \sqrt[n]{a} = \lim_{n \rightarrow \infty} a^{\frac{1}{n}} = 1$$

- but also (which is less obvious):

$$\lim_{n \rightarrow \infty} \sqrt[n]{n} = \lim_{n \rightarrow \infty} n^{\frac{1}{n}} = 1$$

Squeeze theorem

Very, very useful

Theorem (Squeeze theorem)

Let sequences $(a_n)_n, (b_n)_n, (c_n)_n$, satisfy:

$$a_n \leq b_n \leq c_n \quad \forall_n \quad (\text{or at least } \exists_N \forall_{n>N})$$

Then if $a_n \rightarrow g$ and $c_n \rightarrow g$, it must be that $b_n \rightarrow g$.

Theorem (Divergence)

Let sequences $(a_n)_n, (b_n)_n$ satisfy:

$$a_n \leq b_n \quad \forall_n \quad (\text{or at least } \exists_N \forall_{n>N})$$

Then if $a_n \rightarrow \infty$, it must be that $b_n \rightarrow \infty$.

Theorem (Monotone convergence theorem)

Every monotone and bounded sequence converges (to a proper limit).

Example

Let $a_n = \left(1 + \frac{1}{n}\right)^n$. We will show that the sequence a_n is strictly increasing and bounded and therefore has a limit.

Euler's number

Definition

$$e := \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \approx 2.71828$$

Note: this is a *definition* of e (one of few alternatives).

Euler's number

Definition

$$e := \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \approx 2.71828$$

Note: this is a *definition* of e (one of few alternatives).

Lemma

Let $(a_n)_n$ be a sequence satisfying $a_n \rightarrow +\infty$ or $a_n \rightarrow -\infty$.

Then: $\left(1 + \frac{1}{a_n}\right)^{a_n} = e$

Euler's number

Definition

$$e := \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \approx 2.71828$$

Note: this is a *definition* of e (one of few alternatives).

Lemma

Let $(a_n)_n$ be a sequence satisfying $a_n \rightarrow +\infty$ or $a_n \rightarrow -\infty$.

Then: $\left(1 + \frac{1}{a_n}\right)^{a_n} = e$

Special case: $\left(1 + \frac{c}{n}\right)^n = e^c$.

Functions of one variable

Function. Domain and image.

Definition

A function $f: X \rightarrow Y$ is relation that associates with each element of X exactly one element of Y .

Function. Domain and image.

Definition

A function $f: X \rightarrow Y$ is a relation that associates with each element of X exactly one element of Y .

- X is the **domain** of f (also denoted D_f , especially if we need to determine it!)
- Y is the **co-domain**; but usually we are interested in:
- $f(X) \subset Y$ is an **image** (sometimes: range) of f

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)
- function is **"onto"** (surjective) if $\forall_{y \in Y} \exists_{x \in X} : f(x) = y$

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)
- function is **"onto"** (surjective) if $\forall_{y \in Y} \exists_{x \in X} : f(x) = y$
- function is **bijection** (\Rightarrow invertible!) if it is one-to-one and 'onto'

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)
- function is **"onto"** (surjective) if $\forall_{y \in Y} \exists_{x \in X} : f(x) = y$
- function is **bijjective** (\Rightarrow invertible!) if it is one-to-one and 'onto'
- note: it hinges crucially on X and Y ;
take $f(x) = x^2$:

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)
- function is **"onto"** (surjective) if $\forall_{y \in Y} \exists_{x \in X} : f(x) = y$
- function is **bijective** (\Rightarrow invertible!) if it is one-to-one and 'onto'
- note: it hinges crucially on X and Y ;
take $f(x) = x^2$:
 - if $f : \mathbb{R} \rightarrow \mathbb{R}$ it is neither 'onto' nor 1-1

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)
- function is **"onto"** (surjective) if $\forall_{y \in Y} \exists_{x \in X} : f(x) = y$
- function is **bijective** (\Rightarrow invertible!) if it is one-to-one and 'onto'
- note: it hinges crucially on X and Y ;
take $f(x) = x^2$:
 - if $f : \mathbb{R} \rightarrow \mathbb{R}$ it is neither 'onto' nor 1-1
 - if $f : \mathbb{R} \rightarrow [0, +\infty)$ it is 'onto'

Function one-to-one + "onto" = bijection

- function is **one-to-one** (injective) if $x \neq y \Rightarrow f(x) \neq f(y)$;
(it is usually easier to show equivalent statement $f(x) = f(y) \Rightarrow x = y$)
- function is **"onto"** (surjective) if $\forall_{y \in Y} \exists_{x \in X} : f(x) = y$
- function is **bijjective** (\Rightarrow invertible!) if it is one-to-one and 'onto'
- note: it hinges crucially on X and Y ;
take $f(x) = x^2$:
 - if $f : \mathbb{R} \rightarrow \mathbb{R}$ it is neither 'onto' nor 1-1
 - if $f : \mathbb{R} \rightarrow [0, +\infty)$ it is 'onto'
 - if $f : [0, +\infty) \rightarrow \mathbb{R}$ it is 'onto' and one-to-one
and has an inverse! $f^{-1}(y) = \sqrt{y}$

Image and preimage of a set

Let $f: X \rightarrow Y$ be a function

- image of $A \subset X$ is:

$$\{f(x) \in Y : x \in A\}$$

Notation: $f(A)$ or $f[A]$

Image and preimage of a set

Let $f : X \rightarrow Y$ be a function

- image of $A \subset X$ is:

$$\{f(x) \in Y : x \in A\}$$

Notation: $f(A)$ or $f[A]$

- preimage of $B \subset Y$ is:

$$\{x \in X : f(x) \in B\}$$

Notation: $f^{-1}(B)$ (not to be confused with an inverse function!) or $f^{-1}[B]$

Limit of a function

Definition (Limit points of a set)

A point $x \in X$ is a **limit point** of X if there exists a sequence $(x_n)_n$ such that $x_n \in X \setminus \{x\}$ and $x_n \rightarrow x$. Otherwise, we call x an isolated point.

Definition (Heine's limit)

Let x_0 be a limit point of X . A function $f(x) : X \rightarrow Y$ has a limit L in x_0 if for every sequence $(x_n)_n$ such that $x_n \in X \setminus \{x_0\}$ and $x_n \rightarrow x_0$ we have $f(x_n) \rightarrow L$.

Continuity

Definition

A function f is **continuous** at $x_0 \in X$ if for every sequence

$\lim_{n \rightarrow \infty} x_n \rightarrow x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.

Continuity

Definition

A function f is **continuous** at $x_0 \in X$ if for every sequence

$\lim_{n \rightarrow \infty} x_n \rightarrow x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.

- note: x_0 must be in domain of X to consider continuity!

Definition

A function f is **continuous on the domain** (or: on a set $A \subset X$) if it is continuous in every point of its domain (or: of $A \subset X$).

Continuity

Definition

A function f is **continuous** at $x_0 \in X$ if for every sequence $\lim_{n \rightarrow \infty} x_n \rightarrow x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.

- note: x_0 must be in domain of X to consider continuity!

Definition

A function f is **continuous on the domain** (or: on a set $A \subset X$) if it is continuous in every point of its domain (or: of $A \subset X$).

- sum, difference, product of continuous functions is continuous

Continuity

Definition

A function f is **continuous** at $x_0 \in X$ if for every sequence $\lim_{n \rightarrow \infty} x_n \rightarrow x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.

- note: x_0 must be in domain of X to consider continuity!

Definition

A function f is **continuous on the domain** (or: on a set $A \subset X$) if it is continuous in every point of its domain (or: of $A \subset X$).

- sum, difference, product of continuous functions is continuous
- quotient of continuous functions is continuous *if well-defined* (do not divide by 0!)

Continuity

Definition

A function f is **continuous** at $x_0 \in X$ if for every sequence $\lim_{n \rightarrow \infty} x_n \rightarrow x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.

- note: x_0 must be in domain of X to consider continuity!

Definition

A function f is **continuous on the domain** (or: on a set $A \subset X$) if it is continuous in every point of its domain (or: of $A \subset X$).

- sum, difference, product of continuous functions is continuous
- quotient of continuous functions is continuous *if well-defined* (do not divide by 0!)
- polynomial, exponential, rational, logarithmic, trigonometric functions are continuous on their respective

Composition

Definition (Composition)

Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. Then a composition $g \circ f =: h$ if a function $h : X \rightarrow Z$ defined by:

$$h(x) = g(f(x))$$

Composition

Definition (Composition)

Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. Then a composition $g \circ f =: h$ if a function $h : X \rightarrow Z$ defined by:

$$h(x) = g(f(x))$$

Lemma

A composition of continuous functions is a continuous function.

Asymptotes

Definition (Vertical asymptote)

A function $f(x)$ has a vertical asymptote $x = c$ if f has at least one-sided improper limit in c :

$$\lim_{x \rightarrow c^-} f(x) = \pm\infty \text{ or } \lim_{x \rightarrow c^+} f(x) = \pm\infty$$

Asymptotes

Definition (Vertical asymptote)

A function $f(x)$ has a vertical asymptote $x = c$ if f has at least one-sided improper limit in c :

$$\lim_{x \rightarrow c^-} f(x) = \pm\infty \text{ or } \lim_{x \rightarrow c^+} f(x) = \pm\infty$$

- we do not require (and typically do not have) that $c \in D_f$.

Asymptotes

Definition (Vertical asymptote)

A function $f(x)$ has a vertical asymptote $x = c$ if f has at least one-sided improper limit in c :

$$\lim_{x \rightarrow c^-} f(x) = \pm\infty \text{ or } \lim_{x \rightarrow c^+} f(x) = \pm\infty$$

- we do not require (and typically do not have) that $c \in D_f$.
- if f is continuous in x_0 , there can't be an asymptote in x_0 .

Asymptotes

Definition (Vertical asymptote)

A function $f(x)$ has a vertical asymptote $x = c$ if f has at least one-sided improper limit in c :

$$\lim_{x \rightarrow c^-} f(x) = \pm\infty \text{ or } \lim_{x \rightarrow c^+} f(x) = \pm\infty$$

- we do not require (and typically do not have) that $c \in D_f$.
- if f is continuous in x_0 , there can't be an asymptote in x_0 .
- \Rightarrow vertical asymptotes may exist only in the limit points outside of the domain or in discontinuity points.

Asymptotes

Definition (Horizontal asymptote)

A function $f(x)$ has a horizontal asymptote $y = b$ if:

$$\lim_{x \rightarrow +\infty} f(x) = b \text{ or } \lim_{x \rightarrow -\infty} f(x) = b$$

Definition (Oblique asymptote)

A function $f(x)$ has an oblique asymptote $y = ax + b$ if:

$$\lim_{x \rightarrow +\infty} (f(x) - ax - b) = 0 \text{ or } \lim_{x \rightarrow -\infty} (f(x) - ax - b) = 0$$

Useful rules for oblique asymptotes:

$$a := \lim_{x \rightarrow \pm\infty} \frac{f(x)}{x}, \quad b := \lim_{x \rightarrow \pm\infty} (f(x) - ax)$$

Three important limits

It is useful to remember those three rules:

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

Click [here](#) for a beautiful visual proof.

$$\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1$$

Here is a proof with e , but I'll also show another one.

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$$

Another visual proof.