

Mixed Element Methods for Stokes Equations

Will Frey

Virginia Tech

May 13, 2013

Overview

1 Theory

2 Numerical Results

Stokes with Homogeneous Boundary and Zero-Mean Pressure

$$\begin{aligned} -\operatorname{Re}^{-1} \nabla^2 u_1 + \frac{\partial}{\partial x} p &= f_1 \quad \text{in } \Omega, \\ -\operatorname{Re}^{-1} \nabla^2 u_2 + \frac{\partial}{\partial y} p &= f_2 \quad \text{in } \Omega, \\ \frac{\partial}{\partial x} u_1 + \frac{\partial}{\partial y} u_2 &= 0 \quad \text{in } \Omega, \\ u_1 &= 0 \quad \text{on } \partial\Omega, \\ u_2 &= 0 \quad \text{on } \partial\Omega, \\ \int p(x, y) d\Omega &= 0 \quad \text{on } \Omega. \end{aligned} \tag{1}$$

I take $\operatorname{Re} = 1$.

Framework

- ① $\mathbf{u} \in H_0^1(\Omega)^2$
- ② $H_0^1(\Omega)^2 = \{(u_1, u_2) : u_i \in H_0^1(\Omega)\}$
- ③ $p \in L_0^2(\Omega)$
- ④ $L_0^2(\Omega) = \{p \in L^2 : \int p \, d\Omega = 0\}$
- ⑤ Look for solution in $H_0^1(\Omega)^2 \times L_0^2(\Omega)$

We break up the function spaces for velocity \mathbf{u} and pressure p into two distinct Hilbert spaces. We then choose appropriate finite dimensional subspaces to find our finite element solution. I chose the **Taylor-Hood** $\mathbb{P}_2 - \mathbb{P}_1$ finite element.

Taylor-Hood Finite Element




Figure: The $\mathbb{P}_2 - \mathbb{P}_1$ Element

Variational Formulation

Find $\mathbf{u} \in H_0^1(\Omega)^2$ and $p \in L_0^2(\Omega)$ so that

$$\begin{aligned}(\nabla \mathbf{u}, \nabla \mathbf{v}) + (p, \nabla \cdot \mathbf{v}) &= (\mathbf{f}, \mathbf{v}) \quad \text{for all } \mathbf{v} \in H_0^1(\Omega)^2, \\ -(\nabla \cdot \mathbf{u}, q) &= 0 \quad \text{for all } q \in L_0^2(\Omega).\end{aligned}\tag{2}$$

Variational Weak Form Cont'd

Find $(u_{1,h}, u_{2,h}) \in V_h \subset H_0^1(\Omega)^2$ and $p_h \in Q_h \subset L_0^2(\Omega)$ such that

$$\begin{aligned} \int \nabla u_{1,h} \cdot \nabla v_{1,h} d\Omega - \int p_h \frac{\partial v_{1,h}}{\partial x} d\Omega &= \int f_1 v_{1,h} d\Omega \quad \forall v_{1,h} \in V_h, \\ \int \nabla u_{2,h} \cdot \nabla v_{2,h} d\Omega - \int p_h \frac{\partial v_{2,h}}{\partial y} d\Omega &= \int f_2 v_{2,h} d\Omega \quad \forall v_{2,h} \in V_h \text{ and } , \\ \int \left(\frac{\partial u_{1,h}}{\partial x} + \frac{\partial u_{2,h}}{\partial y} \right) q_h d\Omega &= 0 \quad \forall q_h \in Q_h. \end{aligned} \tag{3}$$

Conditions for Well-Posedness

We get a system of the form
$$\begin{bmatrix} A & B \\ B^T & 0 \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} = \begin{bmatrix} f \\ 0 \end{bmatrix}.$$

Ladyzenskaya-Babuska-Breezi Condition (LBB)

A solution exists if and only if

$$\inf_{v \in V} \sup_{u \in U} \frac{a(u, v)}{\|u\| \|v\|} > \alpha_E > 0$$

and the solution is unique if and only if

$$\inf_{u \in U} \sup_{v \in V} \frac{(a(u, v))}{\|u\| \|v\|} > \alpha_U > 0$$

for a generic bilinear form $a(\cdot, \cdot) : U \times V \rightarrow \mathbb{R}$ defined on the Cartesian product of Hilbert spaces $U \times V$.

Stokes Error Estimate

Let $V_h \times Q_h$ satisfy the LBB conditions with polynomials of order max order k in V_h and l in Q_h . Let $u \in H^{k+1}(\Omega)^2$ and $p \in L^2(\Omega)$ be the solution of the Stokes equation. Then

$$\|\nabla(v - v_h)\| + \|p - p_h\| \leq Ch^{\min\{k, l+1\}} (\|\nabla^{k+1} v\| + \|\nabla^l p\|). \quad (4)$$

We see that the error is optimal for both spaces when $k = l + 1$. This is why $\mathbb{P}_2 - \mathbb{P}_1$ works.

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: p_h for $h = 0.1$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: $|u_h|$ for $h = 0.1$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: $|v_h|$ for $h = 0.1$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: Quiver Plot for $h = 0.1$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: p_h for $h = 0.025$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: $|u_h|$ for $h = 0.025$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: $|v_h|$ for $h = 0.025$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: Quiver Plot for $h = 0.025$

$$f_1(x, y) = x^2 y^3 \cos(\pi y), f_2(x, y) = 1e - 4 \cdot |y - 0.5|$$




Figure: p_h for $h = 0.025$

$$f_1(x, y) = x^2 y^3 \cos(\pi y), f_2(x, y) = 1e - 4 \cdot |y - 0.5|$$




Figure: $|u_h|$ for $h = 0.025$

$$f_1(x, y) = x^2 y^3 \cos(\pi y), f_2(x, y) = 1e - 4 \cdot |y - 0.5|$$




Figure: $|v_h|$ for $h = 0.025$

$$f_1(x, y) = y, f_2(x, y) = -x$$




Figure: Quiver Plot for $h = 0.025$

References



Dietrich Braess,

Finite elements: Theory, fast solvers, and applications in solid mechanics,
Cambridge University Press, Third Edition, 2006.



Vivette Girault and Pierre-Arnaud Raviart,

Finite Element Methods for Navier-Stokes Equations: Theory and Algorithms,
Springer-Verlag, Germany, 1986.



Claes Johnson,

Numerical Solution to Partial Differential Equations by the Finite Element Method,
Dover, Mineola, New York, 2009.



Sang Dong Kim, Yong Hun Lee, and Byeong Chun Shin,

Newton's Method for the Navier-Stokes Equations with Finite-Element Initial Guess of Stokes Equation,
Computers and Mathematics with Applications 51 (2006), pp. 805–816.



J.E. Dennis, Jr. and Robert B. Schnabel,

Numerical Methods for Unconstrained Optimization and Nonlinear Equations,
Prentice-Hall, Inc, Englewood Cliffs, New Jersey, 1983.



Philippe G. Ciarlet,

The Finite Element Method for Elliptic Problems,
North Holland Publishing Company, Volume 4, 1978.

The End