

Math 324 C - Summer 2016  
Midterm exam 2  
Wednesday, July 29th, 2016

Name: \_\_\_\_\_

Problem 1	10	
Problem 2	10	
Problem 3	10	
Problem 4	10	
Problem 5	10	
Total	50	


- There are 5 questions on this exam. Make sure you have all five.
- You must show your work on all problems. The correct answer with no supporting work may result in no credit. **Put a box around your FINAL ANSWER for each problem and cross out any work that you don't want to be graded.**
- Give exact answers, and simplify as much as possible. For example,  $\frac{\pi}{\sqrt{2}}$  is acceptable, but  $3\sqrt{3} + \frac{1}{\sqrt{3}}$  should be reduced to  $\frac{10\sqrt{3}}{3}$ .
- If you need more room, use the backs of the pages and indicate to the grader that you have done so.
- Raise your hand if you have a question.
- Any student found engaging in academic misconduct will receive a score of 0 on this exam.
- You have 60 minutes to complete the exam. Budget your time wisely!

GOOD LUCK!

1. (10 pts) Consider the function  $f(x, y, z) = 3x \sin(y) - xz$ .
  - a. Find  $\frac{\partial f}{\partial x}$ ,  $\frac{\partial f}{\partial y}$  and  $\frac{\partial f}{\partial z}$ .
  - b. Let  $v$  be the unit vector with tail at the origin and tip at the point  $(1, \frac{\pi}{3}, \frac{\pi}{2})$  in spherical coordinates  $(\rho, \theta, \phi)$ . Find the directional derivative  $D_v f$  at the point  $(3, -\pi, 1)$ .
  - c. Give the unit vector  $u$  that maximizes  $D_u f(3, -\pi, 1)$ .

- 2a. (3 pts) Give a definition in words of the tangent plane to a surface  $F(x, y, z) = k$  at a point  $p$  in terms of the gradient of  $F$ . (Don't just write down the equation.)
- 2b. (7 pts) Compute the tangent plane to the implicitly defined surface  $x^2z + 3y^3 - z^3 - 3z = 0$  at the point  $(x, y, z) = (1, 1, 1)$ .

- 3a. (3 pts) Let  $f(x, y) = \frac{1}{2}(x^2 + y^2)$ . Write down a formula for  $\nabla f(x, y)$ , and draw the vector  $\nabla f(x, y)$  at each of the three points  $(1, 0)$ ,  $(-2, 1)$  and  $(2, -2)$ .



- 3b. (3 pts) Define what it means for a vector field  $F$  defined on an *arbitrary domain*  $D$  in the plane to be conservative. (You may give any definition equivalent to the one we used in class.)
- 3c. (4 pts) Consider the vector field  $F(x, y) = \langle y^3 \cos(x), -3y^2 \sin(x) \rangle$ . Is  $F$  conservative? If so, give a potential function; if not, explain how you know it isn't conservative.

- 4a. (5 pts) State the fundamental theorem for line integrals, and use it to compute


$$\int_C \sin(y)e^{x \sin(y)} dx + x \cos(y)e^{x \sin(y)} dy,$$

where  $C$  is the line segment from  $(1, \pi)$  to  $(2, \pi)$  followed by the line segment from  $(2, \pi)$  to  $(2, 2\pi)$ .

- 4b. (5 pts) Let  $F$  denote the force field  $F(x, y) = \langle 3 \ln(2xy - 1), x^2 - y^2 \rangle$  defined on  $\{(x, y) : 2xy > 1\}$ . Find the work done by  $F$  on a particle traveling along the path  $r(t) = (t, t^{-1}), 1 \leq t \leq 2$ .

5. Consider the curve  $C$  given by the equation  $xy^2 = 1$  for  $0 < x < \infty$ . Let  $D$  be the (infinite) region above  $C$ , so  $D = \{(x, y) : x > 0, y > \frac{1}{\sqrt{x}}\}$ . Also, define the vector field  $F$  by  $F(x, y) = \langle \frac{1}{x^2+y^2}, \frac{1}{1+y^2} \rangle$ .
- 5a. (5 pts) Let  $S_R$  be the quarter circle of radius  $R$  centered at  $(0, 0)$  in the first quadrant parameterized counter-clockwise. By evaluating the line integral directly, show that

$$\lim_{R \rightarrow \infty} \int_{S_R} F \cdot dr = \frac{\pi}{2}.$$



- 5b. (5 pts) Use the result from part a and Green's theorem to relate

$$\int_C F \cdot dr$$

to a double integral over the region  $D$ . (Hint: use Green's theorem on a closed curve consisting partly of  $C$  and partly of  $S_R$ , and then let  $R \rightarrow \infty$ . You may assume that  $C$  and  $S_R$  meet at the points  $(R, 0)$  and  $(0, R)$ : the error in doing so is negligible.)

5b. (cont.)

5c. (Extra credit) Evaluate your integral from part b: you may use the fact that

$$\int \frac{x}{x^3+1} dx = \frac{1}{3} \log \frac{\sqrt{x^2-x+1}}{x+1} + \frac{1}{\sqrt{3}} \tan^{-1} \frac{2x-1}{\sqrt{3}} + C.$$