

Math 302  
Midterm exam  
Friday, May 28, 5pm

## Instructions

- There are 4 questions on this exam.
- You have 60 minutes to complete the exam, then an additional 20 minutes to upload pictures/scans of your solutions to Canvas.
- Write your name on the top of each page of work that you submit.
- **You must show your work on all problems.** The correct answer with no supporting work may result in no credit. Put a box around your FINAL ANSWER for each problem and cross out any work that you don't want to be graded.
- Give exact answers.
- Any student found engaging in academic misconduct will receive a score of 0 on this exam.

GOOD LUCK!

1. (12 points; 3 each) Deal a hand of 4 cards from a standard 52 card deck – that is, sample 4 cards from the deck without replacement.

- (a) What is the probability of getting two pairs, but not four of a kind? (e.g. 4H, 4S, JC, JS)

A: Count the number of hands that have two pairs but not four of a kind. We have to choose the two values – there are  $\binom{13}{2}$  possibilities – then the suits for each pair – there are  $\binom{4}{2}$  suit combinations for each pair. So the desired probability is  $\frac{\binom{13}{2}\binom{4}{2}^2}{\binom{52}{4}}$ .

- (b) What is the probability that no two of the cards have the same suit?

A: The only way this can happen is if we have one card from each suit. So for each of the four suits we pick a value. Thus the desired probability is  $\frac{13^4}{\binom{52}{4}}$ .

- (c) What is the probability that two of the cards are the ace and king of spades?

A: The number of such hands is  $\binom{50}{2}$ , since we just have to pick the remaining two cards. So the desired probability is  $\frac{\binom{50}{2}}{\binom{52}{4}}$ .

- (d) What is the probability of getting four twos, conditioned on the event that the hand has at least two twos?

A: The probability of having exactly  $j$  twos is  $p_j = \frac{\binom{4}{j}\binom{48}{4-j}}{\binom{52}{4}}$ . The desired probability is  $\frac{p_4}{p_2+p_3+p_4}$ .

2. (12 points; 4 each) Let  $(\Omega, P)$  be a probability space, and let  $A, B$  be events in  $\Omega$ . Assume that  $\mathbb{P}(A \cap B) = 1/2$  and  $\mathbb{P}(A) = 2/3$ . Find  $\mathbb{P}(B)$  in each of the following cases:

(a)  $A, B$  are independent

A: If  $A, B$  are independent, then  $1/2 = \mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B) = 2/3 \mathbb{P}(B)$ , so  $\mathbb{P}(B) = 3/4$  in this case.

(b)  $\mathbb{P}(A \cup B) = 1$

A: By the inclusion-exclusion formula,  $\mathbb{P}(B) = \mathbb{P}(A \cup B) - \mathbb{P}(A) + \mathbb{P}(A \cap B) = 1 - 2/3 + 1/2 = 5/6$ .

(c)  $\mathbb{P}(A|B) = 3/4$ .

A: By the conditional probability formula,  $3/4 = \mathbb{P}(A|B) = \mathbb{P}(A \cap B) / \mathbb{P}(B) = \frac{1}{2\mathbb{P}(B)}$ , so  $\mathbb{P}(B) = 2/3$ .

3. (14 points) There are  $2n$  seats at a circular dinner table, labeled clockwise from 1 to  $2n$ . Suppose that of the  $2n$  guests, exactly  $n$  are vegetarians, and they choose random seats in such a way that all  $\binom{2n}{n}$  arrangements of vegetarians are equally likely. For  $i \in \{1, 2, \dots, 2n\}$ , let  $X_i$  be the indicator random variable of the event that there is a vegetarian in both seats  $i$  and  $i + 1$ .

(a) (3pts) Give the probability mass function of  $X_i$  for any  $i \in \{1, 2, \dots, 2n\}$ .

A:  $X_i$  takes values 0 or 1. For  $X_i = 1$ , seats  $i$  and  $i + 1$  must be occupied. This has probability  $\frac{\binom{2n-2}{n-2}}{\binom{2n}{n}} = \frac{n(n-1)}{2n(2n-1)} = \frac{n-1}{4n-2}$ . Thus  $\mathbb{P}(X_i = 0) = 1 - \mathbb{P}(X_i = 1) = \frac{3n-1}{4n-2}$ .

(b) (3pts) Compute  $\mathbb{E}X_i$ .

A: From the calculation in part (a),  $\mathbb{E}[X_i] = \frac{n-1}{4n-2}$ .

(c) (3pts) Compute  $\mathbb{E}[X_1X_2]$ .

A:  $X_1X_2$  is the indicator of the event that seats 1, 2, and 3 are all occupied. This has probability  $\frac{\binom{2n-3}{n-3}}{\binom{2n}{n}}$ .

(d) (3pts) Let  $X$  denote the total number of pairs of vegetarians sitting next to each other (each pair counts once). Give a formula for  $X$  using the  $X_i$ 's.

A:  $X = \sum_{i=1}^{2n} X_i$ .

(e) (2pts) Compute  $\mathbb{E}[X]$ .

A: By linearity,  $\mathbb{E}[X] = 2n\mathbb{E}[X_i] = \frac{n^2-n}{2n-1}$ .

4. (12 points; 4 each) Consider the random variable  $Y$  with PDF given by

$$f_Y(y) = \frac{e}{2}ye^{-y}, y \geq 1.$$

- (a) Let  $Z = e^Y$ . Find the PDF of  $Z$ .

A: Note that the function  $e^y$  is increasing. Thus,  $\mathbb{P}(Z \leq z) = \mathbb{P}(Y \leq \log z)$ . By the chain rule, we obtain

$$f_Z(z) = \frac{d}{dz}\mathbb{P}(Y \leq \log z) = f_Y(\log z) \cdot \frac{d}{dz}(\log z) = \frac{e}{2}(\log z)e^{-\log z} \cdot \frac{1}{z} = \frac{e}{2} \frac{\log z}{z^2}.$$

- (b) Use your PDF from part (a) to give an integral expression for  $\mathbb{E}[Z]$ , and show that the integral diverges (i.e. that it is  $+\infty$ ).

A: Using the PDF from (a) and the law of the unconscious statistician,  $\mathbb{E}[Z] = \int_e^\infty \frac{e}{2} \frac{\log z}{z} dz \geq \frac{e}{2} \int_e^\infty \frac{1}{z} dz = \infty$ , where we have used the fact that the integrand is non-negative, and  $\log z \geq 1$  for  $z \geq e$ . That the final integral is infinite follows from the 'p-test.'

- (c) Let  $W = \log Y$ . Compute  $\mathbb{P}(1 \leq W \leq 2)$ . You can leave your answer as an integral.

A: Since the function  $e^y$  is increasing in  $y$ ,

$$\mathbb{P}(1 \leq W \leq 2) = \mathbb{P}(e \leq Y \leq e^2) = \int_e^{e^2} f_Y(y) dy.$$