

Digging Holes
in the Desert

Matthew
Arran

Observation
and relevance

Mechanisms

Differential
aeolian transport
Granular
segregation

Investigation

Preliminary
results

Further work

Digging Holes in the Desert: A field-based investigation of sand-dune structure

Matthew Arran
work done with Nathalie Vriend, Michel Louge

GKB Lab Lunch

March 13, 2015

Observation
and relevance

Mechanisms

Differential
aeolian transport
Granular
segregation

Investigation

Preliminary
results

Further work

Introduction

Structure: Observation and relevance

Possible mechanisms for layering

Field Investigation

Preliminary Results

Further Work

Digging Holes in the Desert

Matthew
Arran

Observation
and relevance

Mechanisms

Differential
aeolian transport
Granular
segregation

Investigation

Preliminary
results

Further work

Observations

Digging Holes in the Desert

Matthew
Arran

Observation
and relevance


Mechanisms

Differential
aeolian transport
Granular
segregation

Investigation

Preliminary
results

Further work



Credit: Prof. M. Louge, Jan 2015

Observations

Digging Holes in the Desert

Matthew
Arran

Observation
and relevance

Mechanisms


Differential
aeolian transport
Granular
segregation

Investigation


Preliminary
results

Further work

Observations



Credit: Prof. M. Louge, Jan 2015



Digging Holes in the Desert

Matthew
Arran

Observation
and relevance


Mechanisms

Differential
aeolian transport
Granular
segregation



Investigation

Preliminary
results

Further work



Credit: Prof. M. Louge, Jan 2015



Observations

Relevance

- Oil and gas recovery
- Desert hydrology
- Microbiological dune stabilisation
- Interpretation of sandstones
 - ① Environment of deposition
 - ② Climatic conditions

Dune Migration

Observation
and relevance

Mechanisms


Differential
aeolian transport
Granular
segregation

Investigation

Preliminary
results

Further work

- Migration is via two mechanisms:
 - ① Aeolian transport
 - ② Slip-face avalanches
- Segregation is associated with both processes:
 - ① Differential transport under variable wind conditions
 - ② Kinetic sieving and squeeze expulsion



Threshold velocity

- Consider aeolian transport in inviscid regime

$$Re_p = \frac{u_* d}{\nu} \gg 1 \quad \text{for } \begin{cases} \text{shear velocity } u_* = \sqrt{\tau / \rho_a} \\ \text{particle diameter } d \end{cases}$$

- Ratio of destabilising to stabilising forces on a particle the Shields' number

$$\theta = \frac{\rho_a u_*^2}{(\rho_s - \rho_a)gd} \quad \text{for } \begin{cases} \text{air density } \rho_a \\ \text{sand density } \rho_d \end{cases}$$

- Therefore threshold shear velocity $u_{*t} \propto \sqrt{d}$

Sand flux

- Above threshold velocity, in steady conditions, time-averaged sand flux

$$Q \approx \frac{C_b \rho_a}{g} \sqrt{\frac{\bar{d}}{d_{ref}}} u_*^3 \quad \text{for const. } C_b, d_{ref}$$


- Fraction of flux consisting of particles of diameter d , making up mass fraction M_d/M of source,

$$\frac{Q_d}{Q} \approx \gamma_1 \left(\frac{d}{\bar{d}} \right)^{-\gamma_2} \frac{M_d}{M} \quad \text{for const. } \gamma_1, \gamma_2$$

- But wind conditions fundamentally turbulent, unsteady
- Complex dependence of transported particle size distribution on source distribution, wind conditions

Kinetic sieving and squeeze expulsion

- Consider particle-level motion in avalanche
- Kinetic sieving:
 - ① Smaller particles may fall through smaller voids
 - ② Systematic downwards flux of smaller particles
- Squeeze expulsion:
 - ① More frequent collisions at higher local particle density
 - ② Size-unbiased, diffusive, upwards flux of particles



Distinguishing features

- Magnitude of each effect hard to calculate directly
- Timescales also difficult to quantify
- Layer thickness
 - ① Set by wind variation timescale, dune migration rate
 - ② Set by fundamental avalanche processes
- Grading in layers
 - ① Varies depending on wind variation
 - ② Consistent inverse grading

Approach


- Fieldwork conducted between Doha and Messaieed, Qatar
- Two dunes in dune field studied
- Theodolite and ground penetrating radar surveys
- Samples taken with 'comb' and 'aquarium'



Satellite images courtesy of NASA, CNES/Astrium, Google

'Comb' samples

- Sand wetted for cohesion and pit dug
- Seven adjacent 7mm wide samples extracted
- Size analysis by laser diffraction particle sizer



'Aquarium' samples



- Cuboidal 'aquarium' inserted into undisturbed sand
- Pit dug and 'aquarium' extracted
- Front plate removed
- Low viscosity acrylic resin added and allowed to set
- Sample removed and cut into sections
- Images to be taken with reflected light microscope
- Images to be analysed for particle size distributions



Photos courtesy of Prof M. Louge


Single sample size distribution

Particle size distribution in each sample lognormal, with clear max-likelihood location and shape parameters.



Comparison of location parameters

- Variation between layers greater than in each layer
- No distinct inverse grading observed in grainflow layers
- Grading in aeolian layer similar to that in grainflow layers



Questions raised

- Variation along a single layer?
- Visual distinctiveness of layers?
- Systemic error?

Further work

- 'Aquarium' sample analysis
- GPR survey comparison
- Comparison with site data
 - ① Weather station data
 - ② Sand trap data
- Comparison with lab experiments
 - ① Avalanche depth distribution
 - ② Extent of segregation
- Theoretical modelling

Acknowledgements

- NERC, QNRF
- WCMC-Q, Department of Earth Sciences, Department of Geography
- Prof. Michel Louge, Dr. Ali Sultan, Prof. Anthony Hay, Prof. Anil Netravali
- Paul Mitton, David Page-Croft, Martin Walker, Chris Rolfe

Digging Holes in the Desert

Matthew
Arran

Observation
and relevance

Mechanisms

Differential
aeolian transport
Granular
segregation

Investigation

Preliminary
results

Further work

