

Stat 610 Homework 8

Thursday, November 17, 11:59pm

Assignment

1. The beta distribution $\text{Beta}(\alpha, \beta)$ is a continuous distribution that takes values in $[0, 1]$. The probability density function for $\text{Beta}(\alpha, \beta)$ is

$$\frac{x^{\alpha-1}(1-x)^{\beta-1}}{\Gamma(\alpha)\Gamma(\beta)/\Gamma(\alpha+\beta)}.$$

Implement an accept-reject algorithm where the proposal distribution is uniform on $[0, 1]$ and the target distribution is (a) $\text{Beta}(2, 2)$ or (b) $\text{Beta}(10, 10)$.

In your version of the accept-reject algorithm, include code that allows you to monitor how many times you had to propose a value before one was accepted.

For each type of beta distribution, plot a histogram of the accepted values you obtained from the algorithm and report the average number of proposals per accepted sample. Write a small description of what accounts for the difference in the number of accepted proposals between (a) and (b).

Some notes that might be helpful:

- Notice that when $\alpha = \beta$, the pdf is symmetric.
 - If they are both also greater than 1, the pdf takes its maximum value at $1/2$.
 - The Γ function is a generalization of the factorial function, and is available in R as `gamma`.
 - You can use the `hist` function to plot a histogram.
2. (a) Make a Monte Carlo estimate of the integral $\int_0^1 \cos(\pi x/2) dx$, noticing that the integral should be equivalent to $E[\cos(\pi U/2)]$, where U is a random variable that is uniform on the interval $[0, 1]$.
(b) Make an importance sampling estimate using the importance density $h(x) = 3(1 - x^2)/2$.
(c) Estimate the variances of each estimate and compare. What accounts for the difference?

Submission parameters

Submit two files:

- A pdf writeup containing your plots and answers to the questions.
- A file containing the code you used.