

E115_BirdWatching App Wireframe

March 13, 2025

Bird Watching App

[About](#)[Donate](#)[FAQ](#)

Welcome...! We Help You Learn About Birds in Yanachaga Chemillen National Park

Record Bird Acoustics
To Identify the Bird Species

Button

If the identified bird is one of the 500 species found in the national park, but not one of the 9 listed in the dropdown menu on the right, the app will display *only the species name* and return to this page when clicked again.

Otherwise, it will proceed to the next page.

Would you like to learn more about birds in Yanachaga Chemillén?

Drop down menu:

- Andigena hypoglauca
- Aulacorhynchus coeruleicinctis
- Doliornis sclateri
- Gallinago jamesoni
- Hapalopsittaca melanotis
- Pionus tumultuosus
- Pipile cumanensis
- Rupicola peruvianus
- Tinamus osgoodi

If the identified bird based on its call is one of the nine species in the dropdown menu, or if the user manually selects a bird from the dropdown menu...

Bird Watching App

[About](#) [Donate](#) [FAQ](#)

Do you have any question
about Rupicola Peruvianus?



Enter your question here

Chatbot Response

Your bird is a Rupicola Peruvianus

Familia: Cotingidae
Genus: Rupicola
Species: ***Rupicola peruvianus***
Subspecies: *R. p. aequatorialis* –
R. p. peruvianus – *R. p. sanguinolentus*
– *R. p. saturatus*

Name

Rupicola peruvianus (Latham, 1790)
• Type locality: Chanchamayo, Junín, Peru.

Synonyms


- *Pipra peruviana* (protonym)
- *Rupicola peruviana* (Latham, 1790)

Distribution



One of the iconic birds of the Andes. Fairly large, significantly larger and chunkier than a thrush. Found in cloud forest at middle elevations, most often at a dawn display site where males make piglike squeals and jump and dance for females. Sometimes seen elsewhere in the forest, especially at fruiting trees. Males can be either brilliant crimson or orange, always with black wings and broad silver tertials, orange legs, and puffy rounded crest. Female duller brown with smaller crest; note staring pale eye. Nests on large rocks or cliff faces. Only similar species is Guianan Cock-of-the-rock, but no range overlap.

Previous Rupicola Peruvianus locations within Yanachaga Chemillén



[Click here](#) to learn more about the bird's habitat
or [click here](#) to explore other locations inside the
national park where the bird is likely to be found
based on model predictions.

Coming in next Milestone


If the user selects the option to learn more about the bird's habitat or explore other locations where it is likely to be found...

Bird Watching App


[About](#) [Donate](#) [FAQ](#)

Learn about Habitat Conditions of *Rupicola Peruvianus*: **Deforestation** (drop-down menu with different habitat indicators)

Deforestation



- Previous locations of *Rupicola Peruvianus* inside Yanachaga Chemillen
- Deforested areas inside Yanachaga Chemillen



Deforestation within the national park has been decreasing over the years, although with a peak in 2017. However, it remains a significant threat to birds, as it reduces their food sources. As expected, *Rupicola Peruvianus* has not been previously observed in deforested areas within the park.

Do you have any question about *Rupicola Peruvianus*?


Enter your question here

Chatbot Response

Bird Watching App

[About](#)[Donate](#)[FAQ](#)

Other locations where *Rupicola Peruvianus* may be found inside Yanachaga Chemillen



- Likely to be found
- Less likely to be found

Beta Version – Based on **AI model predictions** derived from similar habitat characteristics where the bird was previously observed. Please consult your local guide or forest ranger before visiting different areas of the national park.

Do you have any question
about *Rupicola Peruvianus*?

Enter your question here

Chatbot Response