E115_BirdWatching App Wireframe

March 13, 2025

ACOUSTICS MONITORING

About

Donate

FAQ

Welcome to Acoustics Monitoring Where We Help You Learn About Birds

Record Bird Acoustics
To Identify the Bird Species



If the identified bird is one of the 500 species found in the national park, but not one of the 9 listed in the dropdown menu on the right, the app will display only the species name and return to this page when clicked again. Otherwise, it will proceed to the next page.

Would you like to learn more about the birds in Yanachaga Chemillén?

Drop down menu:

- Andigena hypoglauca
- Aulacorhynchus coeruleicinctis
- Doliornis sclateri
- Gallinago jamesoni
- Hapalopsittaca melanotis
- Pionus tumultuosus
- Pipile cumanensis
- Rupicola peruvianus
- Tinamus osgoodi

The user pressed the record button to capture a bird's call, which corresponds to one of the nine species in the dropdown menu, or selected a species directly from the dropdown menu...

ACOUSTICS MONITORING

About Donate FAQ

Do you have any question about Rupicola Peruvianus?



Enter your question here

Chatbot Response

Your bird is a Rupicola Peruvianus

Familia: <u>Cotingidae</u> Genus: <u>Rupicola</u>

Species: *Rupicola peruvianus*Subspecies: *R. p. aequatorialis* – *R. p. peruvianus* – *R. p. sanguinolentus*

-R. p. saturatus

Name

Rupicola peruvianus (Latham, 1790)

• Type locality: Chanchamayo, Junín, Peru.

Synonyms


- Pipra peruviana (protonym)
- Rupicola peruviana (Latham, 1790)

Migratory Path



One of the iconic birds of the Andes. Fairly large, significantly larger and chunkier than a thrush. Found in cloud forest at middle elevations, most often at a dawn display site where males make piglike squeals and jump and dance for females. Sometimes seen elsewhere in the forest, especially at fruiting trees. Males can be either brilliant crimson or orange, always with black wings and broad silver tertials, orange legs, and puffy rounded crest. Female duller brown with smaller crest; note staring pale eye. Nests on large rocks or cliff faces. Only similar species is Guianan Cock-of-the-rock, but no range overlap.

Previous Rupicola Peruvianus locations within Yanachaga Chemillen



Click here to view other locations where the birds are likely to be found, based on model predictions (available in upcoming milestones)