

MATH4301 Lecture Notes

Lectures by Ole Warnaar
Notes taken by Jackson Gatenby

MATH4301 Advanced Algebra
The University of Queensland

Version 0.4.0

September 17, 2013

Contents

0.1	Meta	1
1	Notes	2
1.1	Presentations	2
1.2	Coxeter Groups	3
1.3	Dihedral Groups	4
1.4	...?	5
1.5	Lecture 5	6
1.5.1	Semi-direct products	6
1.5.2	Symmetric Groups	7
1.6	The General Theory of Finite Reflection Groups	7
A	Version History	10

0.1 Meta

It is the author's intention that these notes, and the L^AT_EX source code, be publicly released and made available for future reference, so this section records notes about these notes.

At time of writing, this document can be found at <http://jgat.github.io/math4301-notes/lectures.pdf>, and its source code can be found at <https://github.com/jgat/math4301-notes>.

The 2013 offering of MATH4301 Advanced Algebra was divided into two halves, the first half on Galois Theory, taught by Victor Scharaschkin, and the second half on Coxeter Groups, taught by Ole Warnaar. Each half of the course was accompanied by an assignment worth 20% of the grade, and an exam worth 30% of the grade.

Victor has provided typed notes to supplement the first half of the course, not included here. This document contains detailed lecture notes taken during the second half of the course. The notes correspond closely to what was written and said in lectures.

Each section within this document corresponds to notes taken in a single lecture.

Chapter 1

Notes

1.1 Presentations

Let A be an alphabet, the free group $F(A)$ consists of all words over $A \cup A^{-1}$ in which the pairs aa^{-1} and $a^{-1}a$ are forbidden (i.e. $aa^{-1} = a^{-1}a = 1$). The group multiplication corresponds to concatenation of words and removal of forbidden pairs.

Example: if $A = \{a\}$, $F(A) = \{a^k \mid k \in \mathbb{Z}\} \cong (\mathbb{Z}, +)$. If $w_1 = a^4$, $w_2 = a^{-2}$, then $w_1 w_2 = aaaa^{-1}a^{-1} = a^2$.

To make life more interesting we need relations. For example, $A = \{a, b\}$ with relation $b = 1$ gives $(\mathbb{Z}, +)$.

A *presentation* (of a group) $\langle A \mid R \rangle$ consists of a set A of *generators* and a set of relations R between the generators (and their inverses). Elements of the group are again words in A , but two words represent the same element in the group if they can be transformed into each other by the use of R . More formally, $G \cong F(A)/N$ where N is the normal subgroup generated by R .

Example: $\langle a \mid a^k = 1 \rangle \cong \mathbb{Z}/k\mathbb{Z} = \mathbb{Z}_k$ (for $k = 1, 2, \dots$). Formally, $\langle a \mid a^k = 1 \rangle \cong F(a)/\langle a^k \rangle$.

Example: $\langle a, b \mid a^2 = b^2 = (ab)^2 = 1 \rangle$ contains elements $1, a, b, ab, ba, \dots$, however note that $ba = (ab)^{-1} = ab$. Simply guessing which words are distinct is not going to work. The multiplication table of the group is (Exercise: Show that this is all of the elements in the group):

G	1	a	b	ab
1	1	a	b	ab
a	a	1	ab	b
b	b	ab	1	a
ab	ab	b	a	1

Note that $bab = a^{-1}abab = a^{-1} = a$. This is the Klein 4-group $\cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. Geometrically it is the symmetry group of the (non-square) rectangle and a rhombus, where a and b are reflections and ab is rotation by π .

The *word problem* is to decide if two *distinct* words in the generators represent the same/different elements in the group. In 1955, Novikov showed that the word problem is undecidable. This is not the case for Coxeter groups.

1.2 Coxeter Groups

References:

- Bjorner & Brenti: Combinatorics of Coxeter groups (Springer GTM231, '05)
- Bourbaki: Lie groups & Lie algebras (Chap 4-6)
- Cohen: Coxeter groups
- Humphreys: Reflection Groups and Coxeter Groups
- Davis: The Geometry and Topology of Coxeter Groups

Let M be an $r \times r$ symmetric matrix with entries m_{ij} in $\{1, 2, 3, \dots\} \cup \{\infty\}$ with $m_{ii} = 1$ and $m_{ij} = m_{ji} > 1$ for $i \neq j$. Such a matrix is called a *Coxeter matrix*. For example,

$$\begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}$$

Coxeter matrices are often represented as a graph with r labelled vertices $(1, 2, \dots, r)$, and if $m_{ij} \geq 3$, an edge between i and j with a labelling of the edge by m_{ij} . It is standard to drop edge labels which are 3. Hence the above example can be expressed as



Given a Coxeter matrix M (or graph), a *Coxeter system* (W, S) of type M is a set $S = \{s_1, \dots, s_r\}$ and a group

$$W = \langle S \mid (s_i s_j)^{m_{ij}} = 1, \quad 1 \leq i, j \leq r, m_{ij} \neq \infty \rangle.$$

(That is, whenever $m_{ij} \neq \infty$, impose a relation $(s_i s_j)^{m_{ij}} = 1$). The group W is called a *Coxeter group* (of type M). The number r is known as the *rank* of W . Note that $s_i^2 = 1$ for all $1 \leq i \leq r$.

Example: For rank 1, there is only one Coxeter group, $M = (1)$, with the trivial graph:



and the corresponding Coxeter group $W = \langle s \mid s^2 = 1 \rangle \cong \mathbb{Z}_2 = \mathbb{Z}/2\mathbb{Z}$.

For rank 2, we have first,

$$M = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix},$$

with corresponding graph



and corresponding group

$$W = \langle s, t \mid s^2 = t^2 = (st)^2 = 1 \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2$$

(Note that $(s_i s_j)^{m_{ij}} = 1$ implies that $(s_j s_i)^{m_{ij}} = 1$. Why: $(s_j s_i)^{m_{ij}} = (s_j s_i)^{m_{ij}} s_j^2 = s_j (s_i s_j)^{m_{ij}} s_j = s_j^2 = 1$) We will later show that if a Coxeter system has a disconnected graph, then the Coxeter

group will be the direct product of the corresponding groups for each component; hence we will focus on connected graphs. We also have

$$M = \begin{pmatrix} 1 & m \\ m & 1 \end{pmatrix}, \quad m \geq 3,$$

and

$$W = \langle s, t \mid s^2 = t^2 = (st)^m = 1 \rangle.$$

This is known as the *dihedral* group of order $2m$ ($D_m / D_{2m} / I_2(m)$). The dihedral group is the symmetry group of the regular m -gon. For example, $I_2(3)$ is the symmetry group of the triangle, where $s, t, sts = tst$ are reflections and st, ts are rotations. $I_2(4)$ has reflections $s, t, sts = s(ts)^2 = (st)^2s, tst = t(st)^2 = (ts)^2t$.

Note that a word of odd length corresponds to a reflection, and a word of even length corresponds to a rotation; also note that the relation $(st)^m = 1$ embodies the “rotate m times to get the identity” property of the m -gon.

1.3 Dihedral Groups

Recall that a Coxeter group is of the form

$$W = \langle S \mid (s_i s_j)^{m_{ij}} = 1, 1 \leq i \leq j \leq r, m_{ij} \neq \infty \rangle,$$

with the associated matrix $M = (m_{ij})$ where $m_{ii} = 1$ and $M^T = M$. A special case is, for $m > 2$,

$$I_2(m) = \langle s, t \mid s^2 = t^2 = (st)^m = 1 \rangle,$$

a Coxeter group of rank 2. We saw that $I_2(3)$ is the group of symmetries of the equilateral triangle, and $I_2(4)$ is the group of symmetries of the square. We claim that $I_2(m)$ is a group of order $2m$ consisting of m reflections and m rotations of the regular m -gon.

First, for a vector $\alpha \in \mathbb{R}^n$, let H_α denote the hyperplane with normal α , and denote reflection in H_α by r_α . Now, for any vector λ ,

$$r_\alpha(\lambda) = \lambda - \frac{2(\alpha, \lambda)\alpha}{(\alpha, \alpha)}, \quad (1.1)$$

where (a, b) denotes the inner product (vector dot product). Note that $r_\alpha(\lambda) = \lambda$ for every $\lambda \in H_\alpha$, and $r_\alpha(\alpha) = \alpha - \frac{2(\alpha, \alpha)\alpha}{(\alpha, \alpha)} = -\alpha$ as expected. Since we have verified this for a hyperplane of codimension 1 and for a vector normal to the hyperplane, the result is true for all vectors (by Linear Algebra).

Proof. Let s and t be reflections, where the axes of symmetry have an angle of $\theta = \frac{\pi}{m}$, i.e. $s := r_{(1,0)}$ and $t := r_{(\cos \theta, -\sin \theta)}$. Then, $s(1, 0) = (-1, 0)$ and $s(0, 1) = (0, 1)$, so

$$\hat{s} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$

is a matrix representation of s , and

$$\hat{t} = \begin{pmatrix} -\cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix}$$

is a matrix representation of t (Exercise).

If we can show that st is a rotation over $\frac{2\pi}{m}$, then $(st)^k$ will be a rotation over $\frac{2\pi k}{m}$, which will give m distinct rotations. Now,

$$\hat{st} = \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix},$$

which is a rotation matrix for rotation over 2θ (Exercise). Then $(\hat{st})^k$ is a rotation by $2k\theta$ (easy to see geometrically, or show inductively that $(\hat{st})^k$ is a rotation matrix).

It remains to show that there are m distinct words of odd length, and they are all reflections. WLOG, we can say that all words of odd length are of the form $t(st)^{k-1}$ for some $k = 1, 2, \dots, m$, hence there are m distinct words of odd length. Now,


$$\hat{t}(\hat{st})^k = \begin{pmatrix} -\cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix} \begin{pmatrix} \cos 2\theta(k-1) & -\sin 2\theta(k-1) \\ \sin 2\theta(k-1) & \cos 2\theta(k-1) \end{pmatrix} = \begin{pmatrix} -\cos(2k\theta) & \sin 2k\theta \\ \sin 2k\theta & \cos 2k\theta \end{pmatrix}$$

(Exercise), so $t(st)^{k-1} := r_{(\cos k\theta, -\sin k\theta)}$. Hence the m words of odd length are all reflections. \square

Finally, we consider the remaining Coxeter group of rank 2, given by

$$M = \begin{pmatrix} 1 & \infty \\ \infty & 1 \end{pmatrix},$$

with corresponding graph



and corresponding group

$$I_2(\infty) = \langle s, t \mid s^2 = t^2 = 1 \rangle.$$

Elements of this group, known as the ∞ -dihedral group are the words of the form: $1, s, t, st, ts, sts, tst, \dots$. Again this group has a geometric interpretation in terms of reflections (etc.). Before we can describe this, we need some more notation.

Let $V = \mathbb{R}^n$ and for $\alpha \in V$ let H_α denote the hyperplane perpendicular to α . Algebraically, $H_\alpha = \{\lambda \in V : (\lambda, \alpha) = 0\}$. As we have seen, the reflection r_α in H_α is given by its action on $\lambda \in V$ as

$$r_\alpha(\lambda) = \lambda - \frac{2(\alpha, \lambda)}{(\alpha, \alpha)}\alpha = \lambda - \frac{2(\lambda, \alpha)}{\|\alpha\|^2}\alpha = \lambda - (\lambda, \alpha^V)\alpha,$$

where $\alpha^V = \frac{2\alpha}{\|\alpha\|^2}$ (a covector).

1.4 ...?

I was absent from this lecture (Thursday 12 September) (because I slept in) :(. Anyone have notes I can copy?

1.5 Lecture 5

Let $s = r_0$, $t = r_1$, in the group $W = \langle s, t \mid s^2 = t^2 = 1 \rangle$. From last time, we had the following claims:

Claim 1: $r_k = s(st)^k$, $t_{2k} = (ts)^k$, $k \in \mathbb{Z}$.

Claim 2: $\tilde{A}_1 = A_1 \rtimes T$, $A_1 = \{1, r_0\}$, $T = \{t_{2k} : k \in \mathbb{Z}\}$.

Proof. Recall that

$$r_{a,k}(\lambda) = \lambda - \{(\lambda, \alpha) - k\}\alpha^V,$$

if $\alpha = 1$, then $\alpha^V = 2$ (since the dot product must be 2), so

$$r_k = \lambda - 2\{\lambda - k\} = 2k - \lambda.$$

Then,

$$ts(\lambda) = r_1 r_0(\lambda) = r_1(-\lambda) = 2 + \lambda = t_2(\lambda),$$

thus $ts = t_2$, so $(ts)^k = t_{2k}$. Furthermore,

$$s(st)^k(\lambda) = r_0 t_{-2k}(\lambda) = r_0(\lambda - 2k) = -\lambda + 2k = r_k(\lambda),$$

so $s(st)^k = r_k$ □

Note that

$$r_{\alpha,\kappa}(\lambda + \mu) \neq r_{\alpha,\kappa}(\lambda) + r_{\alpha,\kappa}(\mu),$$

instead,

$$\begin{aligned} r_{\alpha,\kappa}(\lambda + \mu) &= \lambda + \mu - \{(\lambda + \mu, \alpha)\alpha^V - \kappa\} \\ &= \lambda + \mu - \{(\lambda, \alpha)\alpha^V + (\mu, \alpha)\alpha^V - \kappa\} \\ &= r_{\alpha,\kappa}(\lambda) + r_{\alpha,\kappa}(\mu) - \kappa\alpha^V \end{aligned}$$

Now,

$$\begin{aligned} r_{\alpha,\kappa} t_\lambda r_{\alpha,\kappa}(\mu) &= r_{\alpha,\kappa}(\lambda + r_{\alpha,\kappa}(\mu)) \\ &= r_{\alpha,\kappa}(\lambda) + \mu - \kappa\alpha^V \\ &= r_{\alpha,0}(\lambda) + \mu, \end{aligned}$$

so $r_{\alpha,\kappa} t_\lambda r_{\alpha,\kappa} = t_{r_\alpha(\lambda)}$.

1.5.1 Semi-direct products

Recall that, for a group G , if $K \leq G$, $N \triangleleft G$, $K \cap N = \{1\}$, and $G = NK$, then we say $G = K \rtimes N$ or $G = N \rtimes K$. Now, consider claim 2:

$$\tilde{A}_1 = \{t_{2k}, r_k \mid k \in \mathbb{Z}\},$$

and consider the subgroups

$$N = T = \{t_{2k} : k \in \mathbb{Z}\}, K = \{1, r_0\} = A_1.$$

Note that $r_k = r_0 t_{-2k}$, so $\tilde{A}_1 = A_1 T$. Thus, claim 2 holds.

1.5.2 Symmetric Groups

Before moving to the general theory, we will discuss one more important example of a finite reflection group (Coxeter group), the symmetric group S_n , also denoted A_{n-1} (not to be confused with the alternating group!) Let us define S_n to be

$$S_n = \langle s_1, \dots, s_{n-1} \mid s_i^2 = 1, (s_i s_{i+1})^3 = 1, (s_i s_j)^2 = 1, |i - j| > 1 \rangle.$$

Clearly the associated Coxeter matrix is

$$M = \begin{pmatrix} 1 & 3 & 2 & \cdots & 2 \\ 3 & 1 & 3 & \cdots & 2 \\ 2 & 3 & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 3 \\ 2 & 2 & \cdots & 3 & 1 \end{pmatrix}$$

with Coxeter graph

The most common description of S_n is as the group of permutations on n letters with s_i acting as “adjacent” transpositions, interchanging the letters in positions i and $i + 1$ (i.e. the cycle $(i \ i + 1)$). For example, $s_2(2, 3, 6, 4, 5, 1) = (2, 6, 3, 4, 5, 1)$. The relation $(s_i s_j)^2 = 1$ for $|i - j| > 1$ may be recast as the commutation relation $s_i s_j = s_j s_i$. Finally, $(s_i s_{i+1})^3 = 1$ can be restated as $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, known as Artin’s braid relation.

Of course, S_n can also be interpreted as a reflection group. Let $V = \mathbb{R}^n$ and s_i the reflection in the hyperplane $H_{\varepsilon_i - \varepsilon_{i+1}} = H_{\alpha_i}$, where ε_i is the i^{th} basis vector in the standard basis. Then, it holds that

$$s_i(\varepsilon_k) = \begin{cases} \varepsilon_k & k \neq i, i + 1 \\ \varepsilon_{i+1} & k = i \\ \varepsilon_i & k = i + 1 \end{cases}$$

so that indeed $s_i^2 = 1$, $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, and $s_i s_j = s_j s_i$ for $|i - j| > 1$. (Exercise: show this).

Note that $\varepsilon_1 + \cdots + \varepsilon_n$ is fixed by the action of $A_{n-1}(= S_n)$, and no other vector linearly independent to this is fixed, hence A_{n-1} only acts on an $(n - 1)$ -dimensional subspace of \mathbb{R}^n , namely $\{\lambda \in \mathbb{R}^n \mid (\lambda, \varepsilon_1 + \cdots + \varepsilon_n) = 0\}$. For example in the case of $S_3 = A_2$,

Now, A_{n-1} is the symmetry group of the $(n - 1)$ -simplex, where for instance the 2-simplex is the triangle, the 3-simplex is the tetrahedron, and so on.

1.6 The General Theory of Finite Reflection Groups

Let V be an Euclidean space, that is, a vector space over \mathbb{R} with a positive definite symmetric bilinear form $(\cdot, \cdot) : V \times V \rightarrow \mathbb{R}$, for example the dot product $(a, b) = a \cdot b$ in \mathbb{R}^n . (Here it suffices to think of \mathbb{R}^n when using Euclidean spaces).

A *reflection* in V is a linear map s_α ($\alpha \in V$) such that

$$s_\alpha(\lambda) = \lambda - (\lambda, \alpha^V) \alpha,$$

where $\alpha^V := \frac{2\alpha}{(\alpha, \alpha)}$ (cf. Equation (1.1)). Note that s_α is an *involution* as well as an *orthogonal transformation*:

Proof.

$$\begin{aligned}
s_\alpha^2(\lambda) &= s_\alpha(\lambda - (\lambda, \alpha^V)\alpha) \\
&= s_\alpha(\lambda) - (\lambda, \alpha^V)s_\alpha(\alpha) \\
&= \lambda - (\lambda, \alpha^V)\alpha + (\lambda, \alpha^V)\alpha \\
&= \lambda,
\end{aligned}$$

so s_α is an involution. Also,

$$\begin{aligned}
(s_\alpha(\lambda), s_\alpha(\mu)) &= (\lambda - (\lambda, \alpha^V)\alpha, \mu - (\mu, \alpha^V)\alpha) \\
&= (\lambda, \mu) - (\mu, \alpha^V)(\lambda, \alpha) - (\lambda, \alpha^V)(\mu, \alpha) + (\lambda, \alpha^V)(\mu, \alpha^V)(\alpha, \alpha) \\
&= (\lambda, \mu) - 2(\mu, \alpha)(\lambda, \alpha)\frac{\alpha}{(\alpha, \alpha)} + 4(\lambda, \alpha)(\mu, \alpha)\frac{(\alpha, \alpha)}{(\alpha, \alpha)^2} \\
&= (\lambda, \mu),
\end{aligned}$$

so s_α is an orthogonal transformation. (Exercise: convince yourself of these results.) \square

A *finite reflection group* is a finite subgroup of the group of orthogonal transformations on V generated by reflections. As we shall see (I hope), all finite reflection groups are Coxeter groups. The converse also holds, however we will not show this.

Lemma 1. *Let $O(V)$ be the group of orthogonal transformations on V , and $W < O(V)$ be a finite reflection group. If $s_\alpha \in W$ is a reflection and $g \in O(V)$ then $gs_\alpha g^{-1} = s_{g(\alpha)}$.*

(Note that there may be elements in W which are not reflections. Hence this does not say that W is normal in $O(V)$, since this lemma does not necessarily hold for all elements of W .)

Proof. Let $\beta = g(\alpha) \in V$. Then first,

$$gs_\alpha g^{-1}(\beta) = gs_\alpha g^{-1}g(\alpha) = gs_\alpha(\alpha) = g(-\alpha) = -g(\alpha) = -\beta.$$

If we can show that $gs_\alpha g^{-1}(H_\beta) = H_\beta$ pointwise, then we are done, because $gs_\alpha g^{-1}$ must then be the reflection s_β .

Let $\lambda \in H_\alpha$. Note that $\lambda \in H_\alpha \iff g(\lambda) \in H_\beta$, since $0 = (\lambda, \alpha) = (g(\lambda), \beta)$. Now,

$$gs_\alpha g^{-1}(g(\lambda)) = gs_\alpha(\lambda) = g(\lambda),$$

so indeed $g(\lambda)$ is fixed by $gs_\alpha g^{-1}$. \square

Corollary 1. *If $s_\alpha, w \in W$ then $s_{w(\alpha)} \in W$. i.e. if H_α is a reflection hyperplane, so is $H_{w(\alpha)}$.*

Proof. Set $g = w$ in Lemma 1. \square

We conclude that reflecting hyperplanes are permuted by the action of w . For example, take $W = A_2 (= S_3)$

$s_\alpha, s_\beta, s_\alpha s_\beta, s_\alpha s_\beta s_\alpha$. Now,

$$\begin{aligned}
H_\alpha &= s_\alpha(H_\alpha) = s_\beta(H_{\alpha+\beta}) = s_\alpha s_\beta(H_{\alpha+\beta}) = s_\beta s_\alpha(H_\beta) = s_\alpha s_\beta s_\alpha(H_\beta) \\
H_\beta &= s_\alpha(H_{\alpha+\beta}) = s_\beta(H_\beta) = s_\alpha s_\beta(H_\alpha) = s_\beta s_\alpha(H_{\alpha+\beta}) = s_\alpha s_\beta s_\alpha(H_\alpha) \\
H_{\alpha+\beta} &= s_\alpha(H_\beta) = s_\beta(H_\alpha) = s_\alpha s_\beta(H_\beta) = s_\beta s_\alpha(H_\alpha) = s_\alpha s_\beta s_\alpha(H_{\alpha+\beta})
\end{aligned}$$

We see that each hyperplane occurs twice as a permutation of each other hyperplane. What we don't see here is that the group elements will also permute the normals of the hyperplane.

To better understand the structure of W , we introduce the notion of a *root system*.

Def: Let Φ be a finite subset of V . Φ is called a root system if, for all $\alpha \in \Phi$,

1. The only multiples of α in Φ are $\alpha, -\alpha$ (so that for each normal vector α , we only have it and its negative).
2. $s_\alpha(\Phi) = \Phi$.

Several remarks are in order:

- Sometimes, condition 1 is dropped from the definition, allowing for *non-reduced* root systems.
- Sometimes a third condition is assumed, that $(\alpha, \beta^\vee) \in \mathbb{Z}$ for $\alpha, \beta \in \Phi$. (We will not assume this.) This leads to *crystallographic* root systems, important in Lie theory.

Lemma 2. *The classification of finite reflection groups boils down to the classification of root systems.*

Proof. Homework. □

Appendix A

Version History

Below we describe briefly the version history of this document, based on versions which have been published at <http://jgat.github.io/math4301-notes/lectures.pdf>. Version numbers will roughly follow the format $v0.n.m$, where n indicates the number of lectures which notes have been written for, and m indicates the number of minor revisions. Once notes have been taken for all lectures, the version number will increment to $v1.0$.

v0.3.0: Initial publication with lectures 1 to 3, §1.1-1.3.

v0.4.0: Lecture 4 placeholder.