

Computing skew-stickiness

Jim Gatheral



Bloomberg Quant Seminar
November 28, 2023

Outline of this talk

- The skew-stickiness ratio (SSR)
- The diamond product
- Trees and forests
- The \mathbb{F} -expansion and stochastic volatility
 - The Bergomi-Guyon smile expansion to all orders
- The SSR in terms of diamonds for very short times
- The time series of SSR from Vola Dynamics
- Properties of the SSR

Implied volatility

According to the definition of implied volatility $\sigma_{BS}(k, T)$, the market price of an option is given by

$$C(S, K, T) = C_{BS}(S, K, T, \sigma_{BS}(k, T))$$

where C_{BS} denotes the Black-Scholes formula and $k = \log K/S$ is the log-strike.

Hedging European options

To hedge options using the Black-Scholes formula, market makers need to hedge two effects:

- The explicit spot effect

$$\frac{\partial C}{\partial S} \delta S$$

and

- The change in implied volatility conditional on a change in the spot

$$\frac{\partial C}{\partial \sigma} \mathbb{E} [\delta \sigma | \delta S].$$

Estimating $\mathbb{E}[\delta\sigma(T)|\delta X]$

- ATM implied volatilities $\sigma(T) = \sigma_{BS}(0, T)$ and stock prices are both observable.
- Market makers can estimate the second component using a simple regression:

$$\delta\sigma(T) = \beta(T) \frac{\delta S}{S} + \text{noise} =: \beta(T) \delta X + \text{noise}. \quad (1)$$

- Then

$$\beta(T) = \frac{\mathbb{E}[\delta\sigma(T)|\delta X]}{\delta X}.$$

.

The skew-stickiness ratio

- For a given time to expiration T , we define the ATM volatility skew

$$\psi(T) = \left. \frac{\partial}{\partial k} \sigma_{BS}(k, T) \right|_{k=0}.$$

- Bergomi [[Ber09](#), [Ber16](#)] calls

$$\mathcal{R}(T) = \frac{\beta(T)}{\psi(T)}$$

the *skew-stickiness ratio* or *SSR*.

Sticky delta and sticky strike

In the old days, traders would typically make one of two assumptions:

- *Sticky delta* where the ATM volatility is fixed.
 - In this case, when S increases to $S + \delta S$, $\delta\sigma(T) = 0$ so $\mathcal{R}(T) = 0$.

or

- *Sticky strike* where the implied volatility is fixed for a given strike independent of the stock price.
 - In this case, when S increases to $S + \delta S$,

$$\delta\sigma(T) = \sigma_{BS}(S + \delta S, T) - \sigma_{BS}(S, T) \approx \psi(T) \delta S$$

so $\beta(T) = \psi(T)$ and $\mathcal{R} = 1$.

- Listed options were thought of as sticky strike and OTC options as sticky delta.

Empirical SSR

- Let's check some skew-stickiness ratios over the period June 1, 2010 to June 1, 2011, reproducing figures from an article I wrote with Mike Kamal [KG09] in the Encyclopedia of Quantitative Finance.

1-month SSR




Figure 1: The 1-month skew-stickiness ratio (SSR). The "sticky strike" green line with slope 1 clearly doesn't fit.

3-month SSR




Figure 2: The 3-month skew-stickiness ratio (SSR). The "sticky strike" green line with slope 1 clearly doesn't fit.

6-month SSR




Figure 3: The 6-month skew-stickiness ratio (SSR). The "sticky strike" green line with slope 1 clearly doesn't fit.

12-month SSR




Figure 4: The 12-month skew-stickiness ratio (SSR). The "sticky strike" green line with slope 1 clearly doesn't fit.

The diamond product

Definition

Given two continuous semimartingales A, B with integrable covariation process $\langle A, B \rangle$, the diamond product of A and B is another continuous semimartingale given by

$$(A \diamond B)_t(T) := \mathbb{E}[\langle A, B \rangle_{t,T} | \mathcal{F}_t] = \mathbb{E}[\langle A, B \rangle_T | \mathcal{F}_t] - \langle A, B \rangle_t,$$

where $\langle A, B \rangle_{t,T} = \langle A, B \rangle_T - \langle A, B \rangle_t$.

Properties of the diamond product

- Commutative: $A \diamond B = B \diamond A$.
- Non-associative: $(A \diamond B) \diamond C \neq A \diamond (B \diamond C)$.
- $A \diamond B$ depends only on the respective martingale parts of A and B .
- $A \diamond B$ is in general not a martingale.

The \mathbb{G} -forest expansion

Theorem 1 (Theorem 1.1 of [FGR22])

Let Y_T be a real-valued, \mathcal{F}_T -measurable random variable with associated martingale $Y_t = \mathbb{E}_t[Y_T]$. Under natural integrability conditions, with a, b small enough, there is a.s. convergence of

$$\log \mathbb{E} \left[e^{aY_T + b\langle Y \rangle_T} \middle| \mathcal{F}_t \right] = aY_t + b\langle Y \rangle_t + \sum_{k \geq 2} \mathbb{G}_t^k(T), \quad (2)$$

where

$$\begin{aligned} \mathbb{G}^2 &= \left(\frac{1}{2}a^2 + b \right) (Y \diamond Y)_t(T), \\ \mathbb{G}^k &= \frac{1}{2} \sum_{j=2}^{k-2} \mathbb{G}^{k-j} \diamond \mathbb{G}^j + (a Y \diamond \mathbb{G}^{k-1}) \text{ for } k > 2. \end{aligned} \quad (3)$$

Trees and forests

- The general term $\mathbb{G}_t^n(T)$ in (3) is naturally written as a linear combination of binary diamond trees¹.
- Hence the terminology \mathbb{G} -forest expansion for (2).
- Specifically, writing \bullet as a short-hand for Y , interpreted as single leaf, we have



$$\begin{aligned}
 \mathbb{G}^2 &= \left(\frac{1}{2}a^2 + b\right) \bullet \text{---} \bullet \\
 \mathbb{G}^3 &= a\left(\frac{1}{2}a^2 + b\right) \bullet \text{---} \bullet \text{---} \bullet \\
 \mathbb{G}^4 &= \frac{1}{2}\left(\frac{1}{2}a^2 + b\right)^2 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet + a^2\left(\frac{1}{2}a^2 + b\right) \bullet \text{---} \bullet \text{---} \bullet \\
 \mathbb{G}^5 &= a\left(\frac{1}{2}a^2 + b\right)^2 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet + \frac{1}{2}a\left(\frac{1}{2}a^2 + b\right)^2 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\
 &\quad + a^3\left(\frac{1}{2}a^2 + b\right) \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \quad (4)
 \end{aligned}$$

¹Trees stolen from [Hai13]!

Forward variance models

- Let S be a strictly positive continuous martingale.
 - Then $X := \log S$ is a semimartingale with quadratic variation process $\langle X \rangle$.
- Defining $V_t dt := d\langle X \rangle_t$, forward variances are given by $\xi_t(u) := \mathbb{E}[V_u | \mathcal{F}_t]$, $u > t$.
 - Forward variances are tradable assets (unlike spot variance).
 - We get a family of martingales indexed by their individual time horizons u .
- Following [BG12], it is natural to specify the dynamics of $\xi_t(u)$ for each $u > t$.
 - all conventional finite-dimensional Markovian stochastic volatility models may be cast as forward variance models.

Trees with colored leaves

- Denote $X \equiv \bullet$.
- We could define $(X \diamond X) = M$, or  $\equiv \bullet$, resulting in trees with leaves of two colors.
 - In a forward variance model, X_t represents the log-stock price and $M_t(T)$, the expected total variance $\int_t^T \xi_t(u) du$.
- In general, we can always identify subtrees in this way and assign them a new variable name (and leaf color).
 - For example, we could define  $\equiv \bullet$ to get



and so on.

The $\tilde{\mathbb{F}}$ -forest expansion

A corollary of the \mathbb{G} -expansion is the $\tilde{\mathbb{F}}$ -expansion of [AGR2020]:

Corollary

The cumulant generating function (CGF) is given by

$$\psi_t(T; a) = \log \mathbb{E}_t \left[e^{i a X_T} \right] = i a X_t - \frac{1}{2} a (a + i) M_t(T) + \sum_{\ell=1}^{\infty} \tilde{\mathbb{F}}_{\ell}(a). \quad (5)$$

where the $\tilde{\mathbb{F}}_{\ell}$ satisfy the recursion

$$\tilde{\mathbb{F}}_0 = -\frac{1}{2} a (a + i) M_t = -\frac{1}{2} a (a + i) \bullet \text{ and for } k > 0,$$

$$\tilde{\mathbb{F}}_{\ell} = \frac{1}{2} \sum_{j=0}^{\ell-2} \left(\tilde{\mathbb{F}}_{\ell-2-j} \diamond \tilde{\mathbb{F}}_j \right) + i a \left(X \diamond \tilde{\mathbb{F}}_{\ell-1} \right). \quad (6)$$

Applying the recursion (6), the first few $\tilde{\mathbb{F}}$ forests are given by

$$\tilde{\mathbb{F}}_0 = -\frac{1}{2}a(a+i) \bullet$$

$$\tilde{\mathbb{F}}_1 = -\frac{i}{2}a^2(a+i) \bullet \text{---} \bullet$$

$$\tilde{\mathbb{F}}_2 = \frac{1}{2^3}a^2(a+i)^2 \bullet \text{---} \bullet + \frac{1}{2}a^3(a+i) \bullet \text{---} \bullet \text{---} \bullet$$

$$\tilde{\mathbb{F}}_3 = (\tilde{\mathbb{F}}_0 \diamond \tilde{\mathbb{F}}_1) + ia \bullet \diamond \tilde{\mathbb{F}}_2$$


$$= \frac{i}{2^2}a^3(a+i)^2 \bullet \text{---} \bullet \text{---} \bullet + \frac{i}{2^3}a^3(a+i)^2 \bullet \text{---} \bullet \text{---} \bullet + \frac{i}{2}a^4(a+i) \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet$$

- Note that the total probability and martingale constraints are satisfied for each tree.
 - That is $\psi_t^T(0) = \psi_t^T(-i) = 0$.

The Bergomi-Guyon smile expansion

- The Bergomi-Guyon (BG) smile expansion (Equation (14) of [BG12]) reads

$$\sigma_{BS}(k, T) = \hat{\sigma}_T + \mathcal{S}_T k + \mathcal{C}_T k^2 + \mathcal{O}(\epsilon^3)$$

where the coefficients $\hat{\sigma}_T$, \mathcal{S}_T and \mathcal{C}_T are complicated combinations of trees such as .

- The beauty of the BG expansion is that in some sense, it yields direct relationships between the smile and autocovariance functionals.

A formal expansion

- Regarding the forest expansion (5) as a formal power series in ϵ whose power counts the forest index ℓ , the characteristic function of the log stock price may be written in the form

$$\varphi_t(T; a) = \exp \left\{ i a X_t - \frac{1}{2} a (a + i) M_t(T) + \sum_{\ell=1}^{\infty} \epsilon^{\ell} \tilde{\mathbb{F}}_{\ell}(a) \right\}.$$

- On the other hand, from for example equation (5.7) of [Gat06], with $X_t = 0$,

$$\int_0^{\infty} \frac{du}{u^2 + \frac{1}{4}} \operatorname{Re} \left[e^{-iuk} \left(\varphi_t^T(u - i/2) - e^{-\frac{1}{2}(u^2 + \frac{1}{4})\Sigma(k)} \right) \right] = 0 \quad (7)$$

where, fixing T , $\Sigma(k) = \sigma_{\text{BS}}^2(k, T)$ T is the implied total variance smile.

- Formally expand $\Sigma(k)$ as

$$\Sigma(k) = \sum_{\ell=0}^{\infty} \epsilon^{\ell} a_{\ell}(k).$$

- The power of ϵ counts the order of the forest expansion.
- We set $\epsilon = 1$ at the end of the computation.
- Equation (7) may then be rewritten in the form

$$\begin{aligned} & \int_0^{\infty} \frac{du}{u^2 + \frac{1}{4}} \operatorname{Re} \left[e^{-iuk} \exp \left\{ -\frac{1}{2} \left(u^2 + \frac{1}{4} \right) \sum_{\ell=0}^{\infty} \epsilon^{\ell} a_{\ell}(k) \right\} \right] \\ &= \int_0^{\infty} \frac{du}{u^2 + \frac{1}{4}} \operatorname{Re} \left[e^{-iuk} e^{-\frac{1}{2} \left(u^2 + \frac{1}{4} \right) M_t(T)} \exp \left\{ \sum_{\ell=1}^{\infty} \epsilon^{\ell} \tilde{\mathbb{F}}_{\ell}(u - i/2) \right\} \right]. \end{aligned} \quad (8)$$

- Matching powers of ϵ on each side of (8) gives the coefficients $a_\ell(k)$ in terms of diamond trees, for any $\ell \in \mathbb{Z}^+$.

$$\begin{aligned}
a_0(k) &= M_t(T) = \bullet \\
a_1(k) &= \left(\frac{k}{M} + \frac{1}{2} \right) \bullet \text{---} \bullet \\
a_2(k) &= \frac{1}{4} (\bullet \text{---} \bullet)^2 \left\{ -\frac{5k^2}{M^3} - \frac{2k}{M^2} + \frac{3}{M^2} + \frac{1}{4M} \right\} \\
&\quad + \frac{1}{4} (\bullet \text{---} \bullet \text{---} \bullet) \left\{ \frac{k^2}{M^2} - \frac{1}{M} - \frac{1}{4} \right\} \\
&\quad + (\bullet \text{---} \bullet \text{---} \bullet) \left\{ \frac{k^2}{M^2} + \frac{k}{M} - \frac{1}{M} + \frac{1}{4} \right\}. \tag{9}
\end{aligned}$$

- It is straightforward to verify that the resulting expansion coincides with that of Bergomi and Guyon up to second order in ϵ .

Computation of the ATM skew $\psi(T)$

- To first order in the forest expansion,

$$\Sigma(k) = M + \epsilon \left(\frac{k}{M} + \frac{1}{2} \right) \text{ (skew stickiness icon) } + \text{ higher order.}$$

- Wlog, set $t = 0$ and drop the subscript.
- To lowest order in the forest expansion

$$\Sigma(0) = \text{ (skew stickiness icon) } = M(T) = \int_0^T \xi(u) du + \text{ higher order.}$$

and, by definition of ψ ,

$$2\sqrt{M}\sqrt{T}\psi(T) = \Sigma'(0) = \epsilon \frac{1}{M} \text{ (skew stickiness icon) } + \text{ higher order.} \quad (10)$$

Computation of the regression coefficient $\beta(T)$

- We rewrite the regression (1) defining $\beta(T)$ as

$$\delta\sigma(0) = \frac{1}{2\sqrt{M}\sqrt{T}} \delta M_t(T) = \alpha + \beta(T) \frac{\delta S_t}{S_t} + \text{noise}.$$

- Thus, for fixed T , (reintroducing t),

$$\beta_t(T) = \frac{\mathbb{E}[d\langle M, X \rangle_t]}{2\sqrt{M}\sqrt{\tau}\mathbb{E}[d\langle X \rangle_t]} = \frac{-\frac{d}{dt}(X \diamond M)_t(T)}{2\sqrt{M}\sqrt{\tau}V_t} + \mathcal{O}(\epsilon), \quad (11)$$

where $X = \log S$, $\tau = T - t$.

- Recall that $(X \diamond M)_t(T) = \mathbb{E}_t \left[\int_t^T d\langle X, M(T) \rangle_s \right]$.

Computation of the SSR for very short times

- Putting (10) and (11) together, we get

$$\mathcal{R}_t(T) = \frac{\beta_t(T)}{\psi_t(T)} = -\frac{M_t(T) \frac{d}{dt}(X \diamond M)_t(T)}{V_t(X \diamond M)_t(T)} + \mathcal{O}(\epsilon^2).$$

- For very short times, with $\tau = T - t$, $M_t(T) \approx V_t \tau$, so

$$\mathcal{R}_t(T) \approx -\tau \frac{d}{dt} \log(X \diamond M)_t(T).$$

To first order in the forest expansion, for τ very small:

The SSR \mathcal{R} is given by the time derivative of the spot-volatility correlation functional $X \diamond M$.

A wild guess

- Suppose that something like

$$\mathcal{R}_t(T) \approx -\tau \frac{d}{dt} \log(X \diamond M)_t(T).$$

holds for all expirations, as implicitly argued in [Ber16].

- Then we have the following examples:

Some SSR examples

- The SABR model
 - $(X \diamond M)_t(T) \propto (T - t)^2$ so $\mathcal{R}_t(T) \approx 2$.
- The Heston model (with $V_t = \bar{V}$),

$$(X \diamond M)_t(T) = \rho \eta \bar{V} \int_0^T dt \int_t^T e^{-\kappa(u-t)} du$$

- For $\tau \ll 1/\kappa$, $(X \diamond M)_t(T) \sim \tau^2$ and $\mathcal{R}_t(T) \approx 2$.
- For $\tau \gg 1/\kappa$, $(X \diamond M)(\tau) \sim \tau$ and $\mathcal{R}_t(T) \approx 1$.

The n -factor Bergomi model

- Let κ_1 be the shortest timescale (largest) mean reversion coefficient and κ_n be the longest timescale (smallest) mean reversion coefficient.
 - For $\tau \ll 1/\kappa_1$, $(X \diamond M)_t(T) \sim \tau^2$ and $\mathcal{R}_t(T) \approx 2$.
 - For $\tau \gg 1/\kappa_n$, $(X \diamond M)_t(T) \sim \tau$ and $\mathcal{R}_t(T) \approx 1$.
- For classical stochastic volatility models in general, $\mathcal{R}_t(T) \approx 2$ for τ small and $\mathcal{R}_t(T) \approx 1$ for τ large.

The skew-stickiness ratio under rough volatility

- Under rough Heston

$$(X \diamond M)_t(T) = \frac{\rho \nu}{\Gamma(1 + \alpha)} \int_t^T \xi_t(s) (T-s)^\alpha ds \approx \frac{\rho \nu}{\Gamma(2 + \alpha)} V_t \tau^{\alpha+1}.$$

- Then, for very short times,

$$\mathcal{R}_t(T) \approx \tau \frac{d}{d\tau} \log(X \diamond M)(\tau) \approx \alpha + 1.$$

- Rough Heston is thus an explicit example of a model in which

$$\mathcal{R}_t(T) \approx 1 + \alpha = H + \frac{3}{2}.$$

- Fukasawa [Fuk21] derives the above result formally for generic rough volatility models in the limit $\tau \rightarrow 0$.

The time series of SSR from Vola Dynamics




Figure 5: The 30 day, 90 day, and 1 year skew-stickiness ratios (SSR), with a trailing window of 60 days from Vola Dynamics.

The time series of SSR from Vola Dynamics




Figure 6: The 30 day, 90 day, and 1 year skew-stickiness ratios (SSR), with a trailing window of 120 days from Vola Dynamics.

Is rough volatility consistent with the SSR time series?

- To be consistent with rough volatility, we would need $\mathcal{R}(T) > \frac{3}{2}$.
 - We see that, empirically, $0.9 < \mathcal{R}(T) < 1.7$.
- Maybe volatility is super-rough?
- Moreover the previous analysis only applies in the limit $T - t = \tau \rightarrow 0$
- and, we haven't taken account of the shape of the forward variance curve.

More carefully: The SSR in AFV models

- In affine forward variance models [GKR19],

$$\frac{dS_t}{S_t} = \sqrt{V_t} dZ_t; \quad d\xi_t(u) = \kappa(u-t) \sqrt{V_t} dW_t.$$

- $M_t(T) = \int_t^T \xi_t(u) du$ so

$$dM_t(T) = \left(\int_t^T \kappa(u-t) du \right) \sqrt{V_t} dW_t =: \tilde{\kappa}(T-t) \sqrt{V_t} dW_t,$$

and

$$d\langle X \rangle_t = V_t dt; \quad d\langle X, M \rangle_t = \rho \tilde{\kappa}(T-t) V_t dt.$$

- Thus

$$(X \diamond M)_t(T) = \rho \int_t^T \xi_t(s) \tilde{\kappa}(T-s) ds.$$

Dependence of the SSR on ξ

- Recall that

$$\mathcal{R}_t(T) = \frac{\beta_t(T)}{\psi_t(T)} = \frac{M_t(T) d \langle X, M \rangle_t}{d \langle X \rangle_t (X \diamond M)_t(T)} + \mathcal{O}(\epsilon^2).$$

- Then, in AFV models,

$$\mathcal{R}_t(T) = \frac{\left(\int_t^T \xi_t(s) ds \right) \tilde{\kappa}(T-t)}{\int_t^T \xi_t(s) \tilde{\kappa}(T-s) ds}. \quad (12)$$

- $\mathcal{R}_t(T)$ depends on the shape of $\xi_t(u)$!
 - A monotonic increasing ξ causes the SSR to increase, and vice versa.

Dependence of the SSR on ξ




Figure 7: The SSR depends on the shape of the forward variance curve.

Path-dependence of the SSR

- With $dW_t = \rho dZ_t + \sqrt{1 - \rho^2} dZ_t^\perp$,

$$\begin{aligned}\xi_t(u) &= \bar{\xi} + \int_{-\infty}^t \kappa(u - r) \sqrt{V_r} dW_r \\ &= \bar{\xi} + \rho \int_{-\infty}^t \kappa(u - r) \frac{dS_r}{S_r} + \text{independent noise.}\end{aligned}$$

- The forward variance curve depends on a weighted average of historical stock returns.
- We deduce that $\mathcal{R}_t(T)$ depends on weighted average historical stock returns.
 - Historical negative returns should cause the SSR to increase, and vice versa.
 - From Figure 7, the SSR seems pretty sensitive to the shape of the forward variance curve.

How good is the skew approximation?

- Under rough Heston, with reasonable parameters:




Figure 8: The rough Heston skew is in red and the first order approximation in blue.

- The first order approximation gets the shape correct but not the level.

How short is short?




Figure 9: The ratio of the rough Heston skew to the BG first order approximation.

- The approximate formula overestimates the skew.
 - The ratio hits 0.95 for expirations later than 15 seconds!

Cancelation of errors?

- Maybe errors in the skew $\psi(T)$ are balanced to some extent by similar errors in $\beta(T)$.
- Figure 10 confirms this guess.
 - However, the skew stickiness ratio remains far above $H + \frac{3}{2}$ in the rough Heston model, for any expiration of practical interest.
- The short term limiting expression for the SSR in terms of $X \diamond M$ has no practical relevance!

SSR under rough Heston ($\lambda = 0$) for various H




Figure 10: Theoretical SSR under rough Heston with $\xi = 0.025$ and the Padé (4,4) approximation [GR19] of the characteristic function.

Summary

- Empirically, implied volatility moves proportionally to the implied volatility skew.
 - The constant of proportionality, the SSR, is historically between 1 and 1.5.
- Under stochastic volatility, the SSR for very short expirations is given by the time derivative of $\log X \diamond M$.
 - $\mathcal{R}_t(T) \approx H + \frac{3}{2}$ where H is the Hurst exponent of V_t .
 - The SSR depends on the recent history of the underlying stock return process.
- For longer expirations, under rough Heston at least, it seems that $\mathcal{R}_t(T) \gtrapprox 2$.
 - We need to check if this remains true under other modeling assumptions.
- For the moment, the empirical behavior of the SSR seems to be inconsistent with (classical or rough) stochastic volatility!

References



Elisa Alòs, Jim Gatheral, and Radoš Radoičić.

Exponentiation of conditional expectations under stochastic volatility.

Quantitative Finance, 20(1):13–27, 2020.



Lorenzo Bergomi

Smile dynamics IV.

Risk December, pages 94–100, 2009.



Lorenzo Bergomi and Julien Guyon.

Stochastic volatility's orderly smiles.

Risk May, pages 60–66, 2012.



Lorenzo Bergomi.

Stochastic Volatility Modeling.

Chapman and Hall/CRC, 2016.



Peter K Friz, Jim Gatheral and Radoš Radoičić.

Forests, cumulants, martingales.

The Annals of Probability, 50(4):1418–1445, 2022.



Volatility has to be rough.

Quantitative Finance, 21(01):1–8, 2021.



The volatility surface: A practitioner's guide.

John Wiley & Sons, 2006.



Affine forward variance models.

Finance and Stochastics, 23(3):501–533, 2019.



Rational approximation of the rough Heston solution.

International Journal of Theoretical and Applied Finance, 22(3):1950010–19, 2019.



Solving the KPZ equation,

Annals of Mathematics, 178:559–664, 2013.



Implied Volatility Surface.

Encyclopedia of Quantitative Finance, Wiley, 2009.