

a
intact forest
ghost forest
prop roots
168 ± 42
GPP
2,962 ± 263
downed wood
151 ± 21
stems +
branches
486 ± 58
soil
233 ± 21
leaves
710 ± 28
water
191 ± 47
CH4
1.1 ± 0.6
wood
+131
(65–197)
CH4 from
15% water
54% soil
28% roots
litterfall
389
(345–456)
mortality
150
(100–300)
sediment
root
production
289
lateral export 346 (235–938): DIC 145, DOC 56, POC 145
unexplained 420 (-185 to 594)
burial
123 (69–157)
to estuary
dead stems +
prop roots
19 ± 2.4
branches
165 ± 40
downed wood
66 ± 52
no GPP
soil
89 ± 52
water
155 ± 35
CH4
8.9 ± 3.6
CH4 from
87% water
12% soil
0% roots
sediment
lateral export: not measured
burial or peat loss: not measured
to estuary
photosynthesis
CO2 emission
CH4 emission
litter / wood / roots
lateral export
burial
unexplained
g C m⁻² yr⁻¹; arrow width ∝ √flux. Solid: measured here (tower GPP; chambers × scanned surface). Translucent, dashed: literature or closure residual.
b
CH4 (g CH4 m−2 yr−1)
15
10
5
0
lateral
(dissolved)
intact
ghost
water
soil
prop root
stem + branch
downed wood
airborne, mean of 4 deployments
bottom-up