
Ghost floor without standing water (Mar 2023)
BL60 dieback soil, Mar 2023; ≤2 cm counted as exposed
BL60 dieback soil, Mar 2023
FLM30 soil, Mar 2022 (same site); ≤2 cm counted as exposed
Pooled ghost soil (FLM30 2022 + MI); ≤2 cm counted as exposed
Pooled ghost soil (FLM30 2022 + MI)
Marco Island ghost soil, Mar 2022-23; ≤2 cm counted as exposed
Marco Island ghost soil, Mar 2022-23
Floor fully inundated
Leaf respiration
Rd25 1.28, LAI 2.3
Rd25 1.62, LAI 5.55
Flooding of the intact floor
all-or-nothing, campaign months
area-weighted, campaign months (high)
area-weighted, campaign months (low)
all-or-nothing, long-term record
high and low tide weighted equally
area-weighted, long-term record
Downed wood volume
negligible (0.01 m³ ha⁻¹)
Krauss low (13 m³ ha⁻¹)
Krauss eyewall (132 m³ ha⁻¹)
Krauss high (181 m³ ha⁻¹)
Q10, day to 24 h (chamber CO2)
stem chambers (4.33)
literature (2.00)
none (1.00)
Tidal phase, intact water CH4
x 0.6
x 2.0
CH4, day to 24 h
x 1.30 (all CH4)
Carbon-balance framing
lateral export, alkalinity retained
lateral export, all returned to air
storage only (burial + wood)
4000
5000
6000
7000
8000
Intact-to-ghost switch (g CO2-eq m−2 yr−1, GWP20)