

Chapter 7

GREEN'S FUNCTIONS



George Green

(1793-1841)

Math/Physics

"Self-Taught" Education

Green's Functions

Math for EM Waves

Lecture 23

7.1 Green's Function : Superposition of Impulse Responses

7.1 Green's Function : Superposition of Impulse Responses

In physics and electrical engineering, the Green's function is one of the most important mathematical tools. There are many different mathematical approaches to define the Green's function. However, here we take a **physics approach** which is very simple and clear:

ex) $L = (\nabla^2 + k^2)$

$$Lf(x) = s(x) \quad (7.1)$$

where L is a **linear** differential operator, $f(x)$ is a classical or quantum mechanical field, and $s(x)$ is a source function.

For a linear system, we can use the **superposition principle** along with the **sifting property** of the Dirac delta function:

$$s(x) = \int dx' s(x') \delta(x - x') \quad (7.2)$$


This means that we can first solve the **impulse response** or the **Green's Function**

$$Lg(x, x') = -\delta(x - x')$$

and then we can find the solution*,

$$f(x) = \int dx' g(x') \delta(x - x')$$

field point.
 $g(x, x')$,
 source point.



(7.4)

This is just a **particular solution**. For the complete solution, we have to need to add a **homogeneous solution**.