Per-atom and per-bond strain across models · fixed cells at a = 2.46 Å

4 × 4 cell 5 × 5 cell reconstructed vacancy

Areal strain | Maximum shear strain | Bond strain

MACE-MP-0 small
MACE-MP-0 medium
MACE-MP-0 large
MACE-MP-0b2 small
MACE-MP-0b2 medium
MACE-MP-0b2 large
MACE-MP-0b3 medium
MACE-MPA-0 medium
MACE-OMAT-0 small
MACE-OMAT-0 medium
MACE-MATPES-PBE-0
MACE-MATPES-r2SCAN-0
MACE-MH-0 omat_pbe
MACE-MH-0 matpes_r2scan
MACE-MH-1 omat_pbe
MACE-MH-1 matpes_r2scan
MatterSim v1.0.0-1M
MatterSim v1.0.0-5M
MACE-OFF23 medium*

−4 −2 0 2 4 6 8
$\varepsilon_A = \det F - 1$ (%)

0 1 2 3 4 5 6
$\gamma = (E_1 - E_2)/2$ (%)

−5.0 −2.5 0.0 2.5 5.0 7.5 10.0 12.5
$\varepsilon_b = \ell/\ell_0 - 1$ (%)

One dot per atom (areal, shear) or per C–C bond (bond strain) · * out of training domain