Host-Microbe Symbiosis





Olhão, Algarve Portugal














SymbNET International Conference on Host-Microbe Symbioses 10-13 June 2024, Olhão, Portugal

ABOUT

Hosts and microbes form long-term symbiotic interactions that shape their biology at multiple levels, from physiology to ecology and evolution. This is a growing field of research which impacts many other areas, from environmental sciences to human health.

The "SymbNET International Conference on Host-Microbe Symbiosis" brings together researchers working on a diverse range of questions, approaches, and model systems. Our meeting, with 6 plenary sessions, will cover different host systems such as plants, animals and humans and different molecular mechanisms, functional understanding and ecological models of the interactions. The purpose is to highlight the most recent advances in the field, common principles between systems, and future directions to explore.

This meeting has 164 participants from 26 countries and it is organized in the context of the EU twinning grant <u>SymbNET</u>, with specific sessions organized by <u>NCCR Microbiomes</u> and <u>CRC Metaorganisms</u>.

P51- Study of gut microbiota associated with gestational diabetes mellitus

<u>Tizziani Benítez-Guerrero</u>¹, Juan Manuel Vélez-Ixta¹, Carmen Josefina Juárez-Castelán¹, Karina Corona-Cervantes¹, Alberto Piña-Escobedo¹, Helga Martínez-Corona¹, Amapola De Sales-Millán¹, Yair Cruz-Narváez Carlos², Yamel Gómez-Cruz², Tito Ramírez-Lozada³, Gustavo Acosta-Altamirano⁴, Mónica Sierra-Martínez⁵, Paola Berenice Zárate-Segura⁶ and Jaime García-Mena¹

- 1 Departamento de Genética y Biología Molecular, Cinvestav, Av. Instituto Politécnico Nacional 2508, Ciudad de México 07360. Mexico
- 2 Laboratorio de Posgrado de Operaciones Unitarias, Escuela Superior de Ingeniería Química e Industrias Extractivas, Instituto Politécnico Nacional, Ciudad de México 07738, Mexico
- 3 Unidad de Ginecología y Obstetricia, Hospital Regional de Alta Especialidad de Ixtapaluca, Carretera Federal México-Puebla Km. 34.5, Col. Zoquiapan, Ixtapaluca 56530, Mexico
- 4 Dirección de Planeación, Enseñanza e Investigación, Hospital Regional de Alta Especialidad de Ixtapaluca, Carretera Federal México-Puebla Km. 34.5, Col. Zoquiapan, Ixtapaluca 56530, Mexico
- 5 Unidad de Investigación en Salud, Hospital Regional de Alta Especialidad de Ixtapaluca, Carretera Federal México-Puebla Km. 34.5, Col. Zoquiapan, Ixtapaluca 56530, Mexico
- 6 Laboratorio de Medicina Traslacional, Escuela Superior de Medicina, Instituto Politécnico Nacional, Ciudad de México 11340, Mexico

Worldwide, changes have been found in the composition of the gut microbiota of pregnant women with gestational diabetes compared to normoglycemic pregnant women. Our objective was to determine the relationship of gut microbiota with clinical, anthropometric, metabolic, and dietary parameters in Mexican pregnant women diagnosed with GD. Stool samples from 54 pregnant women divided into 4 groups were studied: 30 healthy pregnant women (control group), 11 pregnant women diagnosed with GD, 8 women diagnosed with preeclampsia, 5 women with gestational prediabetes. Taxonomic characterization of the gut microbiota was performed. In relation to what was found in the gut microbiota, differences were observed in the relative abundances at the level of phylum and genus between the 4 study groups. Differences in beta diversity were found between the gestational diabetes and control groups. There are differential taxa in each study group evaluated by DESeq2. There are differences in metabolic pathways that are associated with each study group. In our study we observed a greater abundance of genera such as Blautia, Colinsella, Bosea, which have already been associated with gestational diabetes, however, we also found other genera such as Rothia and Enterobacter that could be interesting to explore their role in the pathophysiology of the disease. When the mycobiota was evaluated, a higher abundance of Saccharomyces was found in the case groups and a higher abundance of Fusarium in the control group. Work supported by CONACyT 163235 INFR-2011-01 and CONACYT FORDECYT-PRONACES/6669/2020_Programa Presupuestario F003-Ciencia de Frontera 2019.

P52- Mechanism of metabolic regulation in grain pest beetle-endosymbiont interaction

Soumi Bhattacharyya¹, Aurélien Vigneron², Jean Keller³, Tobias Engl¹, Martin Kaltenpoth¹

- 1 Department of Insect Symbiosis, Max-Planck Institute for Chemical Ecology, 07745 Jena, Germany
- 2 Université Claude Bernard Lyon 1 Faculté des Sciences et Technologies, 69100 Villeurbanne, Lyon, France
- 3 Centre national de la recherche scientifique, 31400 Toulouse, France

Symbiotic associations between insects and microorganisms are crucial for insect evolutionary success. Intracellular symbiotic bacteria, known as endosymbionts, provide essential metabolic components to the insects that are absent in the diet. Tyrosine is necessary for synthesizing the hard, melanized cuticular exoskeleton — a primary defense in beetles. Since beetles cannot synthesize tyrosine themselves, endosymbionts in specialized organs called bacteriomes supplement tyrosine precursors to the host, particularly in nutritionally unbalanced environments. However, maintaining optimal symbiont growth imposes metabolic costs on the host, necessitating a balance between costs and benefits aligned with physiological demands. However, the molecular mechanisms regulating this interaction remain elusive. *Oryzaephilus surinamensis*, a grain pest beetle, houses highly specialized bacterial endosymbionts *Shikimatogenerans silvanidophilus*, that provide tyrosine and L-DOPA precursors crucial for cuticle melanization and sclerotization to the host. However, exposure to



Host-Microbe Symbiosis

10-13th June 2024

GULBENKIAN



This certificate presented to:

Tizziani Benítez Guerrero

Olhão, Portugal, during June 10th-13th, 2024. for attendance and poster presentation International Conference on Host-Microbe Symbiosis, held in at the SymbNET

Luís Teixeira (Project Coordinator)







