

Characterization of Vertically transmitted miRNAs during breastfeeding

##En preparación##

1 Juan Manuel Vélez-Ixta¹, Carmen Josefina Juárez-Castelán¹, Noemí del Socorro Lázaro-
2 Pérez¹, J. J. Castro-Arellano², Silvia Romero-Maldonado³, Enrique Rico-Arzate⁴, Carlos Hoyo-
3 Vadillo⁵, Aparna Krishnakumar¹, Alberto Piña-Escobedo¹, Maria Luisa Pizano-Zarate³, Yair
4 Cruz-Narváez⁴, Jaime García-Mena^{1*}

5 ¹Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Departamento
6 de Genética y Biología Molecular, Ciudad de México, CDMX, México.

7 ²Hospital General “Dr. José María Rodríguez”, Ecatepec de Morelos, Estado de México, México.

8 ³Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Departamento
9 de Farmacología, Ciudad de México, CDMX, México.

10 ⁴Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Gronostajowa 7,
11 Kraków, Poland.

12 ⁵Unidad de Biología Integrativa, The Institute for Obesity Research, Tecnológico de Monterrey, N. L.
13 México.

14 * Correspondence:

15 Jaime García Mena

16 jgmena@cinvestav.mx

17 **Keywords:** Breastfeeding, Human Milk, miRNAs, Neonatal gut, Vertical transmission.

18 Abstract

19 Archaea have been identified as early colonizers of the human intestine, appearing from the first days
20 of life. It is hypothesized that the origin of many of these archaea is through vertical transmission
21 during breastfeeding. In this study, we aimed to characterize the archaeal composition in samples of
22 mother-neonate pairs to observe the potential vertical transmission. We performed a cross-sectional
23 study characterizing the archaeal diversity of 40 human colostrum-neonatal stool samples by next-
24 generation sequencing of V5-V6 16S rDNA libraries. Intra- and inter-sample analyses were carried
25 out to characterize the Archaeal diversity in each sample type. Human colostrum and neonatal stool
26 presented similar core microbiota, mainly composed of the methanogens *Methanoculleus* and
27 *Methanosarcina*. Beta diversity and metabolic prediction results suggest homogeneity between
28 sample types. Further, the co-occurrence network analysis showed associations between Archaea and
29 Bacteria, which might be relevant for these organisms' presence in the human milk and neonatal
30 stool ecosystems. According to relative abundance proportions, beta diversity and co-occurrence
31 analyses, the similarities found imply there is vertical transmission of archaea through breastfeeding.
32 Nonetheless, differential abundances between the sample types suggest other relevant sources for the
33 colonization of archaea to the neonatal gut.

34 1 Introduction

35 Breastfeeding is considered the normative standard for infant feeding and nutrition

36 [Eidelman2012], since this practice has been associated with multiple infant health and immune

development benefits, namely, less incidence of gastrointestinal disease, lower risk to develop respiratory and intestinal tract infections, improved cognitive development and lower mortality rates [@Lyons2020; @Koh2017, @Westerfield2018]. Also, several advantages for the lactating mother have been reported, like less incidence of breast and ovary cancer, metabolic syndrome and postpartum depression [@Sattari2019; @Sayres2018]; also breastfeeding helps to reduce postpartum stress produced by preterm labor [@Black2012].

Due to the wide variety of benefits that breastfeeding provides, multiple health care associations, recomend exclusive breastfeeding for at least the first six months and to continue this practice up to two years [@CDC2012; @WHO2021; @Eidelman2012].

It's worth noting that despite the known benefits of breastfeeding, some mothers may face challenges or barriers to breastfeeding, such as difficulty with latching, inadequate milk supply, or returning to work or school. In these cases, alternative feeding methods such as formula may be necessary to ensure adequate infant nutrition. However, healthcare professionals can provide support and resources to help mothers overcome these challenges and continue breastfeeding if they choose to do so [@Academy].

Human milk (HM) is an unique high nutritional value biofluid, resulted of millions of years of an evolutionary process, that composes the optimum feeding regime both for the mother and the neonate/infant [@Christian2021; @Lyons2020]. The composition of HM widely varies along lactation period, nursing process and circadian cycle [@Ballard2013; @Andreas2015; @Italianer2020], but broadly speaking, it consists on macro and micronutrients, responsible for energy and nutritional HM intake, also biological components esential to the correct development of the infant [@Andreas2015].

As mentioned before, HM can be differentiated in three major stages according to lactation period, these are colostrum, transitional milk and mature milk [@Ballard2013].

In addition to macro and micronutrients, human milk also contains various bioactive molecules, including growth factors, hormones, enzymes, immune modulators, and other compounds that have important roles in infant development and protection [@Ballard2013; @Andreas2015]. For example, human milk contains immunoglobulins, lactoferrin, lysozyme, and other proteins that provide passive immunity to the infant and protect against infections [@Hassiotou2013]. HM also contains oligosaccharides that support the growth of beneficial gut bacteria and reduce the risk of infections and other diseases [@Bode2012]. Additionally, human milk is rich in long-chain polyunsaturated fatty acids, such as docosahexaenoic acid (DHA) and arachidonic acid (ARA), which are important for the development of the brain and nervous system [@Andreas2015].

Additionally, HM contains a complex array of bioactive compounds that confer numerous health benefits to the infant, including antimicrobial, anti-inflammatory, and immunomodulatory effects. These bioactive components include lactoferrin, lysozyme, secretory immunoglobulin A (IgA), oligosaccharides, cytokines, growth factors, and nucleotides, among others [@Ballard2013; @Andreas2015]. The specific composition and concentration of these bioactive components also varies throughout lactation, suggesting that HM provides tailored immunological support to the infant during different stages of development.

MicroRNAs (miRNAs), are highly conserved small Ribonucleic Acid (RNA) molecules from 17-25 nucleotides lenght, miRNAs play important roles in biological processes like human development, metabolism, cellular cycle, proliferation and apoptosis, at postranscriptional level by translational

80 inhibition or mRNA degradation [@Sousa2019; @Tafrihi2018]. MiRNAs can perform their actions
81 either by binding with 3'UTR mRNA transcripts or also by forming complexes with proteins and
82 other miRNAs [@Sousa2019; @Tafrihi2018; @Tome-Carneiro2018].

83 MiRNAs are first transcribed in the cell nucleus from genomic Desoxyribonucleic Acid (gDNA) by
84 the action of RNA polymerase II, in an initial structure called pri-miRNA, thereafter sliced by a
85 complex formed between Drosha and its cofactor DGCR8 in order to produce a pre-miRNA. This
86 new miRNA precursor is later exported to the cytoplasm by exportin 5 and its cofactor Ran. Finally,
87 exported pre-miRNAs are processed by Dicer and argonaut proteins to produce mature miRNAs
88 [@He2016].

89 MicroRNAs (miRNAs) are important regulators of gene expression, and they play crucial roles in
90 many biological processes, including development, differentiation, metabolism, and disease. They
91 are small non-coding RNA molecules that are typically 17-25 nucleotides in length, and they act
92 post-transcriptionally to regulate gene expression by binding to complementary sequences in target
93 messenger RNA (mRNA) transcripts, resulting in translational inhibition or mRNA degradation
94 [@Sousa2019; @Tafrihi2018].

95 MiRNAs can interact with mRNA transcripts by binding to their 3' untranslated region (UTR), but
96 they can also form complexes with proteins and other miRNAs to regulate gene expression
97 [@Sousa2019; @Tafrihi2018; @Tome-Carneiro2018].

98 MiRNAs are initially transcribed from genomic DNA by RNA polymerase II into a primary
99 transcript (pri-miRNA), which is processed in the nucleus by the Drosha-DGCR8 complex to
100 generate a precursor miRNA (pre-miRNA). The pre-miRNA is then transported to the cytoplasm by
101 Exportin 5 and its cofactor Ran, where it is further processed by Dicer and Argonaut proteins to
102 generate mature miRNAs that can then bind to their target mRNA transcripts [@He2016].

103 miRBase is the primary public repository and online resource for microRNA sequences and
104 annotation, is responsible for microRNA gene nomenclature and provides a wide-range of
105 information on published microRNAs, including their sequences, their biogenesis precursors,
106 genome coordinates and context, literature references, deep sequencing expression data and
107 community-driven annotation [@Kozomara2019].

108 miRBase is a database that collects, curates and provides comprehensive information on microRNAs.
109 It was established in 2002 and has since become the primary source for microRNA nomenclature,
110 sequences and functional annotation. The database is managed and maintained by a team of curators,
111 who collect and organize data on new and existing microRNAs from published literature and
112 experimental datasets.

113 In addition to providing information on microRNA sequences and annotation, miRBase also includes
114 tools for sequence search, target prediction, and expression analysis. Users can search for
115 microRNAs by name, sequence, or genome location, and can also browse and download datasets of
116 microRNA expression profiles from a variety of tissues and conditions.

117 Despite previous works had studied and documented miRNAs in human milk, yet little is known
118 about miRNome at population level. Moreover, investigation regarding human milk components,
119 both in mother and neonate intestine, could help to clarify the influence of human milk in the infant
120 gut miRNome and the possible similarities with the mother.

2 Materials and Methods

Study design and selection of subjects

The cross-sectional descriptive study consisted of 40 mother-neonatal pairs of patients from the "Dr José María Rodríguez" General Hospital, located in Ecatepec de Morelos, State of Mexico (19°36'35'' N, 99° 3'36'' W). The samples were obtained from healthy lactating women and exclusively breastfed newborns. Colostrum and neonatal stool samples were collected from each mother-infant pair 1 to 6 days after delivery, from November 2017 to January 2018. The inclusion criteria were as follows: (1) Mexican origin with at least two generations of ancestry, (2) gestational age between 37 and 41 weeks, (3) birth weight between 2500 - 4500g, (4) Apgar score greater than 7 at 5 min after birth. Exclusion criteria: (1) Probiotic and alcohol consumption, (2) smoking, (3) diabetes before or during pregnancy, (4) antibiotic use during the last trimester of pregnancy and before sampling. The participants were given a survey where sociodemographic and clinical data was recorded (maternal age, gestational age at delivery, and type of delivery; newborn sex, and age). Written informed consent was obtained from all participants before the study, following the 2013 Declaration of Helsinki. The protocol was approved by the Ethics Committee of the General Hospital "Dr José María Rodríguez" (Project identification code: 217B560002018006).

Sample collection

he 60 dyad samples will be collected from mothers who gave their informed consent according to the Declaration of Helsinki, 2013, the protocol was approved by the "Instituto Nacional de Perinatología" Bioethics Committee. After collection, samples will be stored at -20°C until processing at "Laboratorio de Referencia y apoyo para la Caracterización de Genomas, Transcriptomas y Microbiomas" Genetics and Molecular Biology, Cinvestav Zacatenco. Inclusion criteria for participants are samples from healthy women, residing in Mexico, full-term delivery, at first breastfeeding week (0 to 7 postpartum days) and who had previously breastfed their infant. Neonate stool sample will correspond to mother offspring which human milk sample was collected and had breastfed. In addition, exclusion criteria will be alcohol, cigarettes or other drugs consumption during pregnancy or sample collection, mothers who do not breastfed and mothers suffering metabolic syndrome, chronic or neurodegenerative diseases.

RNA extraction

One (1) mL of colostrum will be separated in Neofuge 12R centrifuge (HealForce®) using fresh 1.5 mL polypropylene tubes at 13 800 rpm, 5 min, 4° C, repeat this process until 3 distinct phases are clearly distinguishable. Discard intermediate (proteic) phase with 100 µL filter pipette tips and corresponding micropipette; keep lipidic and cellular fraction. Wash two times with 1 mL cold PBS, ensuring to disrupt the pellet in vortex, centrifuge under the same conditions after each wash. Spin 10 seconds. For the stool samples, weigh 200 mg and wash with cold PBS in a 0.4 µm filter, collect in 15 mL Falcon tube. Transfer 1 mL to a fresh centrifuge tube. For total RNA extraction, use mirVana™ miRNA Isolation Kit (Invitrogen™, Cat. AM1560) following manufacturer's instructions. Check RNA integrity in 2% low melting point agarose gel. Concentration was assessed in Qubit with Qubit™ RNA High Sensitivity kit (Cat. Q32852).

Preparation of the 16S rDNA library and next generation sequencing

The small RNA (sRNA) Library with TruSeq® Small RNA Library Prep (Cat., RS-200-0012) were prepared following manufacturer's instructions as follows. Starting total RNA should be at least at

200 ng and sRNA 10-50 ng/ μ L. Use DNase treatment after extraction. Positive control is Human Brain Total RNA (Cat., # AM7962). For 3' adapter ligation, add 1 μ L RA3 and 1 μ L RNA in 5 μ L water, mix and incubate 70 °C for 2 minutes. In a fresh PCR tube, mix the following reagents for each sample: 2 μ L HML, 1 μ L RNase inhibitor and 1 μ L T4 RNA ligase, Deletion Mutant. Add 4 μ L to each sample, incubate at 28° C for 1 hour, add 1 μ L STP, mix and incubate for 15 min more, then place on ice. For 5' adapter ligation, take a fresh PCR tube and for each sample: add 1.1 μ L RA5, incubate 70°C, 2 min, add 1.1 μ L ATP 10 mM, mix and add 1.1 μ L T4 RNA ligase. Add 3 μ L to each 3' adapter ligated sample. Incubate at 28° C for 1 hour, then place on ice. From 6 μ L ligated RNA, add 1 μ L RNA RT Primer, mix and incubate 70° C, 2 min, then place on ice. In a fresh PCR tube, mix for each sample: 2 μ L First Strand Buffer, 0.5 dNTP Mix 12.5 mM, 1 μ L DDT 100 mM, 1 μ L RNase inhibitor and 1 μ L SuperScript II Reverse Transcriptase. Add 5.5 to each library and incubate at 50° C for 1 hour. For each index, mix in a fresh PCR tube: 8.5 μ L ultrapure water, 25 μ L PML, 2 μ L RP1 and 2 μ L RPIx. Add 37.5 μ L to each library, incubate on thermal cycler with the following program: 98° C, 30 s; 11 cycle of 98° C for 10 s, 60° C for 30 s and 72° C for 15 s; then 72° C, 10 min, and hold at 4° C. Place each library in High Sensitivity DNA chip (Agilent, Cat. 5067-4626) to verify the presence of clear bands corresponding to sRNA region. This is a safe stop point, cDNA can be stored at -25° C up to 7 days. cDNA construct purification will be conducted with 7% polyacrylamide gel (Novex TBE Gels, Cat., EC6265BOX). Load 1 μ L CRL with 1 μ L DNA loading dye in wells 1 and 2; 1 μ L HRL with 1 μ L DNA loading dye in well 3; and 25 μ L cDNA library with 25 μ L DNA loading dye in wells 4 and 5. Run the gel for 60 min, 145 V. Stain the gel with ethidium bromide for 2-3 min, cut the bands between 147-157 nt, corresponding to sRNA of 22-30 nt, respectively. Disrupt the bands in 0.5 mL gel breaker tube, centrifuge 20 000 x g for 2 min, add 200 μ L ultrapure water and mix for two hours to elute DNA. Place in 5 μ m filter and centrifuge at 600 x g for 10 seconds.

187 **Data analysis**

188 All bioinformatic analyses were conducted using R 4.3.3 in RStudio, version 2024.04.2.0, along with
 189 additional packages including ComplexHeatmap version 2.18.0, DESeq2 version 1.42.1 (Love et al.,
 190 2014), ALDEx2 version 1.34 (Fernandes et al., 2013), and edgeR version 4.0.16 (Robinson et al.,
 191 2010). A heatmap, a statistical visualization technique used to reveal similar patterns shared by
 192 subsets of rows and columns in matrix-type data (Gu, 2022), was generated using the
 193 ComplexHeatmap package. For this, we included 100% of the miRNAs with at least 50% abundance.
 194 Additionally, a chart was created to highlight the most representative miRNAs found in the two
 195 groups of milk samples used in this study by calculating the mean and standard deviation for miRNA
 196 abundance in the Before and After milk samples.

197 For differential analysis, three methods available in R were applied. The first, DESeq2 version
 198 1.42.1, employs shrinkage estimation for dispersions and fold changes to improve the stability and
 199 interpretability of the estimates (Love et al., 2014). This method generated a volcano plot, a scatter
 200 plot displaying the statistical significance (P-value) against the magnitude of change (fold change),
 201 allowing for the visual identification of genes with significantly large fold changes. In the volcano
 202 plot, the most statistically significant genes are located at the top; upregulated genes are to the right,
 203 while downregulated genes appear on the left (Doyle). For the volcano plot, we used
 204 log2FoldChange and P-value variables, and for DESeq2 bar plots, we selected genes with P-value <
 205 0.05 and log2FoldChange > 1.5. The second method, ALDEx2, takes as input the sequencing count
 206 per gene for each replicate and a list indicating which replicate belongs to each condition. It produces
 207 a table for each gene that includes individual mean expression per gene for each sample, mean
 208 expression per gene for each condition, overall mean expression, average absolute fold difference,

mean effect size, median within-condition difference, and zero quantile (Fernandes et al., 2013). A volcano plot was also created to determine variable cutoffs, utilizing mean difference values between the log-ratio (clr) of the groups (diff.btw) and the expected P-value from the Wilcoxon rank-sum test (wi.ep). For ALDEx bar plots, diff.btw and miRNA data were included.

Finally, edgeR version 4.0.16 (Robinson et al., 2010) was employed to estimate gene-specific dispersions using conditional maximum likelihood on total counts for each gene. An empirical Bayes procedure shrinks these dispersions toward a consensus value, and differential expression for each gene is then evaluated using a test for overdispersed data. Here, a volcano plot was also generated, using Log Fold Change (logFC) and P-value variables. For ALDEx bar plots, diff.btw and the miRNAs found in the samples were included.

3 Results

EN PREPARACIÓN

4 Discussion

EN PREPARACIÓN

5 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6 Author Contributions

Salas-López M: Conceptualization, Investigation, Methodology, Visualization, Writing-original draft, reviewing & editing. Velez-Ixta J-M: Data curation, Formal analysis, Visualization, Software. Rojas-Guerrero D-L: Investigation, Methodology, Visualization. Piña-Escobedo A: Methodology, Supervision. Hernández-Hernández J-M: Validation, Writing- reviewing & editing. Rangel-Calvillo M-N: Conceptualization, Methodology, Funding acquisition. Pérez-Cruz C: Conceptualization, Validation, Writing- reviewing & editing. Corona-Cervantes K: Data curation, Methodology, Writing – reviewing & editing, Investigation. Juárez-Castelán C-J: Methodology, Investigation, Supervision, García-Mena J: Writing – original draft, reviewing & editing, Resources, Project administration and Funding acquisition.

7 Funding

This work was supported by Consejo Nacional de Ciencia y Tecnología (CONACYT-163235) INFR-2011-01, and by CONACyT FORDECYT-PRONACES/6669/2020_Programa Presupuestario F003-Ciencia de Frontera 2019. The funding body was not involved in study design; collection, management, analysis, and interpretation of data; or the decision to submit for publication.

8 Acknowledgments

We are grateful to all families who agreed to participate in the study, to Rodrigo García-Gutiérrez for technical support in the laboratory, and Viridiana Rosas-Ocegueda for administrative assistance. CP-C (47399), JMH-H (225832), and JG-M (19815) are Fellows from the Sistema Nacional de Investigadores, Mexico. We thank for CONAHCyT Doctoral Fellowships 997152 to JMV-I, 777953 to KC-C; Master Fellowships 1140881 to MS-L, 1140600 to DLR-G, and Estancias-Posdoctorales-por-México Fellowship 321600 to CJJ-C.

248 9 Reference styles

- 249 Auguie, B. (2017). *gridExtra: Miscellaneous Functions for “Grid” Graphics*. Available at:
 250 <https://CRAN.R-project.org/package=gridExtra>
- 251 Ballard, O., and Morrow, A. L. (2013). Human Milk Composition: Nutrients and Bioactive Factors.
 252 *Pediatric Clinics* 60, 49–74. doi: 10.1016/j.pcl.2012.10.002
- 253 Bang, C., Weidenbach, K., Gutschmann, T., Heine, H., and Schmitz, R. A. (2014). The Intestinal
 254 Archaea *Methanosphaera stadtmanae* and *Methanobrevibacter smithii* Activate Human
 255 Dendritic Cells. *PLOS ONE* 9, e99411. doi: 10.1371/journal.pone.0099411
- 256 Bisanz, J. E. (2018). qiime2R: Importing QIIME2 artifacts and associated data into R sessions.
 257 Available at: <https://github.com/jbisanz/qiime2R>
- 258 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al.
 259 (2019). Reproducible, interactive, scalable and extensible microbiome data science using
 260 QIIME 2. *Nat Biotechnol* 37, 852–857. doi: 10.1038/s41587-019-0209-9
- 261 Brugère, J.-F., Borrel, G., Gaci, N., Tottey, W., O’Toole, P. W., and Malpuech-Brugère, C. (2014).
 262 Archaeobiotics. *Gut Microbes* 5, 5–10. doi: 10.4161/gmic.26749
- 263 Camara, A., Konate, S., Tidjani Alou, M., Kodio, A., Togo, A. H., Cortaredona, S., et al. (2021).
 264 Clinical evidence of the role of *Methanobrevibacter smithii* in severe acute malnutrition. *Sci*
 265 *Rep* 11, 5426. doi: 10.1038/s41598-021-84641-8
- 266 Chaudhary, P. P., Conway, P. L., and Schlundt, J. (2018). Methanogens in humans: potentially
 267 beneficial or harmful for health. *Appl Microbiol Biotechnol* 102, 3095–3104. doi:
 268 10.1007/s00253-018-8871-2
- 269 Corona-Cervantes, K., García-González, I., Villalobos-Flores, L. E., Hernández-Quiroz, F., Piña-
 270 Escobedo, A., Hoyo-Vadillo, C., et al. (2020). Human milk microbiota associated with early
 271 colonization of the neonatal gut in Mexican newborns. *PeerJ* 8, e9205. doi:
 272 10.7717/peerj.9205
- 273 David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., et al.
 274 (2014). Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 505, 559–
 275 563. doi: 10.1038/nature12820
- 276 Deng, Z.-L., Szafranski, S. P., Jarek, M., Bhujju, S., and Wagner-Döbler, I. (2017). Dysbiosis in
 277 chronic periodontitis: Key microbial players and interactions with the human host. *Sci Rep* 7,
 278 3703. doi: 10.1038/s41598-017-03804-8
- 279 Di Simone, N., Santamaria Ortiz, A., Specchia, M., Tersigni, C., Villa, P., Gasbarrini, A., et al.
 280 (2020). Recent Insights on the Maternal Microbiota: Impact on Pregnancy Outcomes.
 281 *Frontiers in Immunology* 11. Available at:
 282 <https://www.frontiersin.org/articles/10.3389/fimmu.2020.528202> (Accessed October 2,
 283 2023).
- 284 Djemai, K., Drancourt, M., and Tidjani Alou, M. (2022). Bacteria and Methanogens in the Human
 285 Microbiome: a Review of Syntrophic Interactions. *Microb Ecol* 83, 536–554. doi:
 286 10.1007/s00248-021-01796-7
- 287 Douglas, G. M., Maffei, V. J., Zaneveld, J. R., Yurgel, S. N., Brown, J. R., Taylor, C. M., et al.
 288 (2020). PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol* 38, 685–688. doi:
 289 10.1038/s41587-020-0548-6
- 290 Drancourt, M., Djemai, K., Gouriet, F., Grine, G., Loukil, A., Bedotto, M., et al. (2021).
 291 *Methanobrevibacter smithii* Archaeemia in Febrile Patients With Bacteremia, Including Those
 292 With Endocarditis. *Clinical Infectious Diseases* 73, e2571–e2579. doi: 10.1093/cid/ciaa998
- 293 Drevland, R. M., Waheed, A., and Graham, D. E. (2007). Enzymology and evolution of the pyruvate
 294 pathway to 2-oxobutyrate in *Methanocaldococcus jannaschii*. *J Bacteriol* 189, 4391–4400.
 295 doi: 10.1128/JB.00166-07

296 Dridi, B., Henry, M., Khéchine, A. E., Raoult, D., and Drancourt, M. (2009). High Prevalence of
 297 Methanobrevibacter smithii and Methanosphaera stadtmanae Detected in the Human Gut
 298 Using an Improved DNA Detection Protocol. *PLOS ONE* 4, e7063. doi:
 299 10.1371/journal.pone.0007063

300 Eikmanns, B., Jaenchen, R., and Thauer, R. K. (1983). Propionate assimilation by methanogenic
 301 bacteria. *Arch. Microbiol.* 136, 106–110. doi: 10.1007/BF00404782

302 Ekiel, I., Sprott, G. D., and Patel, G. B. (1985). Acetate and CO₂ assimilation by Methanothrix
 303 concilii. *J Bacteriol* 162, 905–908. doi: 10.1128/jb.162.3.905-908.1985

304 Fernández, L., Langa, S., Martín, V., Maldonado, A., Jiménez, E., Martín, R., et al. (2013). The
 305 human milk microbiota: Origin and potential roles in health and disease. *Pharmacological*
 306 *Research* 69, 1–10. doi: 10.1016/j.phrs.2012.09.001

307 Fitzstevens, J. L., Smith, K. C., Hagadorn, J. I., Caimano, M. J., Matson, A. P., and Brownell, E. A.
 308 (2017). Systematic Review of the Human Milk Microbiota. *Nutrition in Clinical Practice* 32,
 309 354–364. doi: 10.1177/0884533616670150

310 Gaci, N., Borrel, G., Tottey, W., O'Toole, P. W., and Brugère, J.-F. (2014). Archaea and the human
 311 gut: New beginning of an old story. *World J Gastroenterol* 20, 16062–16078. doi:
 312 10.3748/wjg.v20.i43.16062

313 Gállego-Bravo, A. K., García-Mena, J., Piña-Escobedo, A., López-Jiménez, G., Gutiérrez-Castillo,
 314 M. E., and Tovar-Gálvez, L. R. (2023). Monitoring of a microbial community during
 315 bioaugmentation with hydrogenotrophic methanogens to improve methane yield of an
 316 anaerobic digestion process. *Biotechnol Lett* 45, 1339–1353. doi: 10.1007/s10529-023-03414-
 317 7

318 Ghavami, S. B., Rostami, E., Sephay, A. A., Shahrokh, S., Balaii, H., Aghdaei, H. A., et al. (2018).
 319 Alterations of the human gut Methanobrevibacter smithii as a biomarker for inflammatory
 320 bowel diseases. *Microbial Pathogenesis* 117, 285–289. doi: 10.1016/j.micpath.2018.01.029

321 Gila-Diaz, A., Arribas, S. M., Algara, A., Martín-Cabrejas, M. A., López de Pablo, Á. L., Sáenz de
 322 Pipaón, M., et al. (2019). A Review of Bioactive Factors in Human Breastmilk: A Focus on
 323 Prematurity. *Nutrients* 11, 1307. doi: 10.3390/nu11061307

324 Gomathi, V., Ramasamy, K., Ramalakshmi, A., and Ramanathan, A. (2009). Methan Emission by
 325 Gut Symbionts of Termites.

326 Goodchild, A., Raftery, M., Saunders, N. F. W., Guilhaus, M., and Cavicchioli, R. (2004). Biology of
 327 the cold adapted archaeon, Methanococcoides burtonii determined by proteomics using liquid
 328 chromatography-tandem mass spectrometry. *J Proteome Res* 3, 1164–1176. doi:
 329 10.1021/pr0498988

330 Grine, G., Boualam, M. A., and Drancourt, M. (2017). Methanobrevibacter smithii, a methanogen
 331 consistently colonising the newborn stomach. *Eur J Clin Microbiol Infect Dis* 36, 2449–2455.
 332 doi: 10.1007/s10096-017-3084-7

333 Gu, Z. (2022). Complex Heatmap Visualization. *iMeta*. doi: 10.1002/imt2.43

334 Hasan, N., and Yang, H. (2019). Factors affecting the composition of the gut microbiota, and its
 335 modulation. *PeerJ* 7, e7502. doi: 10.7717/peerj.7502

336 Hugon, P., Lagier, J.-C., Colson, P., Bittar, F., and Raoult, D. (2017). Repertoire of human gut
 337 microbes. *Microbial Pathogenesis* 106, 103–112. doi: 10.1016/j.micpath.2016.06.020

338 Huynh, H. T. T., Pignoly, M., Nkamga, V. D., Drancourt, M., and Aboudharam, G. (2015). The
 339 Repertoire of Archaea Cultivated from Severe Periodontitis. *PLOS ONE* 10, e0121565. doi:
 340 10.1371/journal.pone.0121565

341 Ignacio, A., Fernandes, M. R., Rodrigues, V. A. A., Groppo, F. C., Cardoso, A. L., Avila-Campos,
 342 M. J., et al. (2016). Correlation between body mass index and faecal microbiota from
 343 children. *Clinical Microbiology and Infection* 22, 258.e1-258.e8. doi:
 344 10.1016/j.cmi.2015.10.031

Jiménez, E., de Andrés, J., Manrique, M., Pareja-Tobes, P., Tobes, R., Martínez-Blanch, J. F., et al. (2015). Metagenomic Analysis of Milk of Healthy and Mastitis-Suffering Women. *J Hum Lact* 31, 406–415. doi: 10.1177/0890334415585078

Kers, J. G., and Saccenti, E. (2022). The Power of Microbiome Studies: Some Considerations on Which Alpha and Beta Metrics to Use and How to Report Results. *Frontiers in Microbiology* 12. Available at: <https://www.frontiersin.org/articles/10.3389/fmicb.2021.796025> (Accessed July 26, 2023).

Koskinen, K., Pausan, M. R., Perras, A. K., Beck, M., Bang, C., Mora, M., et al. (2017). First Insights into the Diverse Human Archaeome: Specific Detection of Archaea in the Gastrointestinal Tract, Lung, and Nose and on Skin. *mBio* 8, e00824-17. doi: 10.1128/mBio.00824-17

Le Doare, K., Holder, B., Bassett, A., and Pannaraj, P. S. (2018). Mother's Milk: A Purposeful Contribution to the Development of the Infant Microbiota and Immunity. *Frontiers in Immunology* 9. Available at: <https://www.frontiersin.org/articles/10.3389/fimmu.2018.00361> (Accessed April 28, 2023).

Lecours, P. B., Marsolais, D., Cormier, Y., Berberi, M., Haché, C., Bourdages, R., et al. (2014). Increased Prevalence of *Methanosphaera stadtmanae* in Inflammatory Bowel Diseases. *PLOS ONE* 9, e87734. doi: 10.1371/journal.pone.0087734

Liu, C., Cui, Y., Li, X., and Yao, M. (2021). microeco: an R package for data mining in microbial community ecology. *FEMS Microbiology Ecology* 97, fiae255. doi: 10.1093/femsec/fiae255

Liu, C., Li, C., Jiang, Y., Zeng, R. J., Yao, M., and Li, X. (2023). A guide for comparing microbial co-occurrence networks. *iMeta* 2, e71. doi: 10.1002/imt2.71

Love, M., Ahlmann-Eltze, C., Forbes, K., Anders, S., Huber, W., FP7, R. E., et al. (2023). DESeq2: Differential gene expression analysis based on the negative binomial distribution. doi: 10.18129/B9.bioc.DESeq2

Lurie-Weinberger, M. N., and Gophna, U. (2015). Archaea in and on the Human Body: Health Implications and Future Directions. *PLOS Pathogens* 11, e1004833. doi: 10.1371/journal.ppat.1004833

Lyu, Z., Shao, N., Akinyemi, T., and Whitman, W. B. (2018). Methanogenesis. *Curr Biol* 28, R727–R732. doi: 10.1016/j.cub.2018.05.021

Matarazzo, F., Ribeiro, A. C., Feres, M., Faveri, M., and Mayer, M. P. A. (2011). Diversity and quantitative analysis of Archaea in aggressive periodontitis and periodontally healthy subjects. *Journal of Clinical Periodontology* 38, 621–627. doi: 10.1111/j.1600-051X.2011.01734.x

Maya-Lucas, O., Murugesan, S., Nirmalkar, K., Alcaraz, L. D., Hoyo-Vadillo, C., Pizano-Zárate, M. L., et al. (2019). The gut microbiome of Mexican children affected by obesity. *Anaerobe* 55, 11–23. doi: 10.1016/j.anaerobe.2018.10.009

McMurdie, P. J., and Holmes, S. (2013). phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* 8, e61217.

Mihajlovski, A., Alric, M., and Brügère, J.-F. (2008). A putative new order of methanogenic Archaea inhabiting the human gut, as revealed by molecular analyses of the *mcrA* gene. *Research in Microbiology* 159, 516–521. doi: 10.1016/j.resmic.2008.06.007

Mihajlovski, A., Doré, J., Levenez, F., Alric, M., and Brügère, J.-F. (2010). Molecular evaluation of the human gut methanogenic archaeal microbiota reveals an age-associated increase of the diversity. *Environmental Microbiology Reports* 2, 272–280. doi: 10.1111/j.1758-2229.2009.00116.x

Moossavi, S., and Azad, M. B. (2020). Origins of human milk microbiota: new evidence and arising questions. *Gut Microbes* 12, 1667722. doi: 10.1080/19490976.2019.1667722

393 Moubareck, C. A. (2021). Human Milk Microbiota and Oligosaccharides: A Glimpse into Benefits,
394 Diversity, and Correlations. *Nutrients* 13, 1123. doi: 10.3390/nu13041123

395 Mukhopadhyay, B., Purwantini, E., Conway de Macario, E., and Daniels, L. (1991). Characterization
396 of aMethanosarcina strain isolated from goat feces, and that grows on H₂-CO₂ only after
397 adaptation. *Current Microbiology* 23, 165–173. doi: 10.1007/BF02091977

398 Nava, G. M., Carbonero, F., Croix, J. A., Greenberg, E., and Gaskins, H. R. (2012). Abundance and
399 diversity of mucosa-associated hydrogenotrophic microbes in the healthy human colon. *ISME*
400 *J* 6, 57–70. doi: 10.1038/ismej.2011.90

401 Nguyen-Hieu, T., Khelaifia, S., Aboudharam, G., and Drancourt, M. (2013). Methanogenic archaea
402 in subgingival sites: a review. *APMIS* 121, 467–477. doi: 10.1111/apm.12015

403 Nkanga, V. D., Henrissat, B., and Drancourt, M. (2017). Archaea: Essential inhabitants of the human
404 digestive microbiota. *Human Microbiome Journal* 3, 1–8. doi: 10.1016/j.humic.2016.11.005

405 Notarbartolo, V., Giuffrè, M., Montante, C., Corsello, G., and Carta, M. (2022). Composition of
406 Human Breast Milk Microbiota and Its Role in Children’s Health. *Pediatr Gastroenterol*
407 *Hepatol Nutr* 25, 194–210. doi: 10.5223/pghn.2022.25.3.194

408 Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., et al. (2022).
409 *vegan: Community Ecology Package*. Available at: [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
410 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan)

411 Pace, R. M., Williams, J. E., Robertson, B., Lackey, K. A., Meehan, C. L., Price, W. J., et al. (2021).
412 Variation in Human Milk Composition Is Related to Differences in Milk and Infant Fecal
413 Microbial Communities. *Microorganisms* 9, 1153. doi: 10.3390/microorganisms9061153

414 Palmer, C., Bik, E. M., Giulio, D. D., Realman, D. A., and Brown, P. O. (2007). Development of the
415 Human Infant Intestinal Microbiota | PLOS Biology. Available at:
416 <https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.0050177> (Accessed July
417 11, 2022).

418 Probst, A. J., Auerbach, A. K., and Moissl-Eichinger, C. (2013). Archaea on Human Skin. *PLOS*
419 *ONE* 8, e65388. doi: 10.1371/journal.pone.0065388

420 R Core Team (2021). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
421 Foundation for Statistical Computing. Available at: <https://www.R-project.org/>

422 Ramezani, A., Nolin, T. D., Barrows, I. R., Serrano, M. G., Buck, G. A., Regunathan-Shenk, R., et al.
423 (2018). Gut Colonization with Methanogenic Archaea Lowers Plasma Trimethylamine N-
424 oxide Concentrations in Apolipoprotein e^{-/-} Mice. *Sci Rep* 8, 14752. doi: 10.1038/s41598-
425 018-33018-5

426 Rescigno, M., Urbano, M., Valzasina, B., Francolini, M., Rotta, G., Bonasio, R., et al. (2001).
427 Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to
428 sample bacteria. *Nature immunology* 2. doi: 10.1038/86373

429 Robeson, M. S., O’Rourke, D. R., Kaehler, B. D., Ziemski, M., Dillon, M. R., Foster, J. T., et al.
430 (2020). RESCRIPt: Reproducible sequence taxonomy reference database management for the
431 masses. 2020.10.05.326504. doi: 10.1101/2020.10.05.326504

432 Robichaux, M., Howell, M., and Boopathy, R. (2003). Methanogenic Activity in Human Periodontal
433 Pocket. *Curr Microbiol* 46, 0053–0058. doi: 10.1007/s00284-002-3807-5

434 Rodríguez, J. M. (2014). The Origin of Human Milk Bacteria: Is There a Bacterial Entero-Mammary
435 Pathway during Late Pregnancy and Lactation? *Advances in Nutrition* 5, 779–784. doi:
436 10.3945/an.114.007229

437 RStudio Team (2022). *RStudio: Integrated Development Environment for R*. Boston, MA: RStudio,
438 PBC. Available at: <http://www.rstudio.com/>

439 Ruiz, L., García-Carral, C., and Rodríguez, J. M. (2019). Unfolding the Human Milk Microbiome
440 Landscape in the Omics Era. *Frontiers in Microbiology* 10. Available at:

- <https://www.frontiersin.org/articles/10.3389/fmicb.2019.01378> (Accessed September 29, 2023).
- Saghehdu, V., Patrone, V., Miragoli, F., and Morelli, L. (2017). Abundance and Diversity of Hydrogenotrophic Microorganisms in the Infant Gut before the Weaning Period Assessed by Denaturing Gradient Gel Electrophoresis and Quantitative PCR. *Frontiers in Nutrition* 4. Available at: <https://www.frontiersin.org/articles/10.3389/fnut.2017.00029> (Accessed September 30, 2023).
- Samuel, B. S., and Gordon, J. I. (2006). A humanized gnotobiotic mouse model of host–archaeal–bacterial mutualism. *Proceedings of the National Academy of Sciences* 103, 10011–10016. doi: 10.1073/pnas.0602187103
- Scanlan, P. D., Shanahan, F., and Marchesi, J. R. (2008). Human methanogen diversity and incidence in healthy and diseased colonic groups using mcrA gene analysis. *BMC Microbiology* 8, 79. doi: 10.1186/1471-2180-8-79
- Selma-Royo, M., Calvo-Lerma, J., Bäuerl, C., Esteban-Torres, M., Cabrera-Rubio, R., and Collado, M. C. (2022). Human milk microbiota: what did we learn in the last 20 years? *Microbiome Research Reports* 1, 19. doi: 10.20517/mrr.2022.05
- Sereme, Y., Mezouar, S., Grine, G., Mege, J. L., Drancourt, M., Corbeau, P., et al. (2019). Methanogenic Archaea: Emerging Partners in the Field of Allergic Diseases. *Clinic Rev Allerg Immunol* 57, 456–466. doi: 10.1007/s12016-019-08766-5
- Shima, S., Warkentin, E., Thauer, R. K., and Ermler, U. (2002). Structure and function of enzymes involved in the methanogenic pathway utilizing carbon dioxide and molecular hydrogen. *Journal of Bioscience and Bioengineering* 93, 519–530. doi: 10.1016/S1389-1723(02)80232-8
- Stinson, L. F., Sindi, A. S. M., Cheema, A. S., Lai, C. T., Mühlhäusler, B. S., Wlodek, M. E., et al. (2021). The human milk microbiome: who, what, when, where, why, and how? *Nutrition Reviews* 79, 529–543. doi: 10.1093/nutrit/nuaa029
- Thapa, B. R. (2005). Health factors in colostrum. *Indian J Pediatr* 72, 579–581. doi: 10.1007/BF02724182
- Togo, A. H., Grine, G., Khelaifia, S., des Robert, C., Brevaut, V., Caputo, A., et al. (2019). Culture of Methanogenic Archaea from Human Colostrum and Milk. *Sci Rep* 9, 18653. doi: 10.1038/s41598-019-54759-x
- Uruakpa, F. O., Ismond, M. A. H., and Akobundu, E. N. T. (2002). Colostrum and its benefits: a review. *Nutrition Research* 22, 755–767. doi: 10.1016/S0271-5317(02)00373-1
- van de Pol, J. A. A., van Best, N., Mbakwa, C. A., Thijs, C., Savelkoul, P. H., Arts, I. C. W., et al. (2017). Gut Colonization by Methanogenic Archaea Is Associated with Organic Dairy Consumption in Children. *Front Microbiol* 8, 355. doi: 10.3389/fmicb.2017.00355
- Wampach, L., Heintz-Buschart, A., Hogan, A., Muller, E. E. L., Narayanasamy, S., Laczny, C. C., et al. (2017). Colonization and Succession within the Human Gut Microbiome by Archaea, Bacteria, and Microeukaryotes during the First Year of Life. *Frontiers in Microbiology* 8. Available at: <https://www.frontiersin.org/articles/10.3389/fmicb.2017.00738> (Accessed July 11, 2022).
- Weaver, G. A., Krause, J. A., Miller, T. L., and Wolin, M. J. (1986). Incidence of methanogenic bacteria in a sigmoidoscopy population: an association of methanogenic bacteria and diverticulosis. *Gut* 27, 698–704. doi: 10.1136/gut.27.6.698
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., et al. (2019). Welcome to the tidyverse. *Journal of Open Source Software* 4, 1686. doi: 10.21105/joss.01686

- Willis, A. D. (2019). Rarefaction, Alpha Diversity, and Statistics. *Frontiers in Microbiology* 10. Available at: <https://www.frontiersin.org/articles/10.3389/fmicb.2019.02407> (Accessed July 26, 2023).
- Xiong, X., Rao, Y., Tu, X., Wang, Z., Gong, J., Yang, Y., et al. (2022). Gut archaea associated with bacteria colonization and succession during piglet weaning transitions. *BMC Vet Res* 18, 243. doi: 10.1186/s12917-022-03330-4
- Yazici, C., Arslan, D. C., Abraham, R., Cushing, K., Keshavarzian, A., and Mutlu, E. A. (2016). Breath Methane Levels Are Increased Among Patients with Diverticulosis. *Dig Dis Sci* 61, 2648–2654. doi: 10.1007/s10620-016-4174-6
- Yu, Y., Lee, C., Kim, J., and Hwang, S. (2005). Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnology and Bioengineering* 89, 670–679. doi: 10.1002/bit.20347
- Zhang, H., DiBaise, J. K., Zuccolo, A., Kudrna, D., Braidotti, M., Yu, Y., et al. (2009). Human gut microbiota in obesity and after gastric bypass. *Proceedings of the National Academy of Sciences* 106, 2365–2370. doi: 10.1073/pnas.0812600106

11 Supplementary Material

Supplementary Material can be found at

Data Availability Statement

The datasets generated for this study can be found in the the NCBI BioProject ID XXXX Link <https://www.ncbi.nlm.nih.gov/bioproject/XXXX>.