

# Paradise Lost? Estimating the Economic Impact of Hawaii’s Mandatory Quarantine using Synthetic Historical Controls

JARED GREATHOUSE<sup>†</sup>

<sup>†</sup>*Department of Policy and Management, Georgia State University*  
E-mail: [jgreathouse3@student.gsu.edu](mailto:jgreathouse3@student.gsu.edu)

**Summary** This paper estimates the causal impact of Hawaii’s March 2020 mandatory quarantine on its tourism industry and labor market using the Synthetic Historical Control method. Analyzing visitor days, hotel occupancy, leisure/hospitality employment, and unemployment claims (all in annualized year-over-year growth rates), the study reveals substantial negative impacts from the quarantine. Observed outcomes fall significantly outside 90% conformal out-of-sample prediction intervals, indicating a clear treatment effect. Placebo tests further validate the model’s internal credibility. This research demonstrates the method’s utility and informs the broader debate on economic impacts of anti-COVID policy.

**Keywords:** *Synthetic Controls, Policy Evaluation, COVID-19*

## 1. INTRODUCTION

Many of the most pressing policy questions in economics concern one-time, systemic shocks: What would the United States’ economy have looked like absent the revolution? What would have happened to unemployment in Great Britain without Brexit? In these cases, credible untreated units often do not exist, making standard causal inference tools, like difference-in-differences (DiD) and synthetic control (SCM), difficult or impossible to apply.

This paper takes up one such case: Hawaii’s near-total quarantine in March 2020. While most regions experienced economic contraction from COVID-19 through voluntary distancing, fear, or uneven mandates, Hawaii implemented one of the few deliberate shut-downs in the U.S.: a policy aimed not merely at mitigation of the virus by agreement, but by actively suppressing travel and commerce. On March 26, 2020, the state imposed a 14-day quarantine on all arrivals, effectively sealing its borders. Hawaii’s economic and tourism collapse was thus, in part, an intended consequence. Within a month, a third of the state’s labor force was unemployed [Ivanova \(2020\)](#).

Quantifying economic impacts of such an intervention poses serious challenges. As noted by [Callaway and Li \(2023\)](#) and [Bilgel \(2022\)](#), the assumptions underlying DiD and SCM are often violated in COVID-19 settings.<sup>1</sup> In a global pandemic, there are no unaffected

<sup>1</sup>Of course, this is not universal. Researchers have productively applied SCM and related panel methods to study the effects of non-pharmaceutical interventions during the pandemic, especially when some entities implemented policies like mask mandates or lockdowns while others did not [Yang \(2021\)](#); [Friedson et al. \(2021\)](#); [Gibson and Sun \(2023\)](#); [Breidenbach and Mitze \(2021\)](#); [Bulle et al. \(2022\)](#); [Ke and Hsiao \(2022, 2023\)](#). These comparative case study designs work well when variation in treatment exists across geography.

units. Hawaii is geographically and economically distinct, and every plausible control — whether U.S. state or island nation — faced concurrent shocks. In such a setting, there is no valid cross-sectional counterfactual.

To address this, I apply the Synthetic Historical Control (SHC) method introduced by [Chen et al. \(2024\)](#), which constructs a counterfactual using only the treated unit’s own pre-intervention history. SHC represents a fundamental shift in the SCM paradigm: rather than borrowing strength across space, it borrows strength across time. This allows us to recover counterfactual outcomes when the treated unit is one-of-a-kind and no credible donors exist, exactly the situation in Hawaii.

This paper contributes to the literature on the broader economic consequences of lockdown-style policies—beyond their direct effects on infection or mortality. Understanding the economic costs of aggressive containment measures is vital for future health policy, particularly in tourism-dependent regions where travel restrictions can have outsized effects. Hawaii’s case offers a uniquely compelling setting: its geographic isolation, centralized governance, and heavy reliance on tourism allowed it to pursue a form of near-total border control that was effectively impossible in mainland states. As such, Hawaii provides an empirical glimpse into a “zero-COVID”-like strategy implemented within the U.S. federal system—what might be called Great Isolation with Aloha characteristics.

Methodologically, this is the first real-world policy application of the SHC approach outside of its original proposal. More broadly, this paper illustrates how SHC opens a path forward for causal inference in settings where standard panel methods are unusable — including many of the policy shocks economists most want to study.

## 2. METHODOLOGY

We know what happened given that Hawaii mandated two-week quarantines. We do not know what would have happened had they not done this. Given its novelty and radical deviation from SCM, this section summarizes the SHC method by [Chen et al. \(2024\)](#). For a comprehensive treatment of the method, including the theoretical proofs and asymptotic properties, readers are referred to [Chen et al. \(2024\)](#).

**Notation** Let  $\mathbb{R}$  denote the set of real numbers, and let  $\|\cdot\|_1$  denote the usual  $\ell_1$  vector norm. Throughout, I denote sets using calligraphic letters (e.g.,  $\mathcal{T}, \mathcal{S}, \mathcal{N}$ ) for clarity. Scalars are represented using plain lowercase letters (e.g.,  $h, t, n, m$ ), and matrices are denoted by bold uppercase letters (e.g.,  $\mathbf{X}, \mathbf{W}, \mathbf{Y}$ ). Given a vector  $\mathbf{x} = (x_1, x_2, \dots, x_T)^\top \in \mathbb{R}^T$  and an index set  $\mathcal{I} \subseteq \{1, \dots, T\}$ , I write  $\mathbf{x}_{\mathcal{I}} := (x_t)_{t \in \mathcal{I}} \in \mathbb{R}^{|\mathcal{I}|}$  to denote the subvector of  $\mathbf{x}$  corresponding to indices in  $\mathcal{I}$ , preserving their original order.

Let  $\mathcal{T} = \{1, 2, \dots, T\}$  index time at a monthly frequency. Define the pre-treatment period as  $\mathcal{T}_1 := \{t \in \mathcal{T} : t \leq T_0\}$  and the post-treatment period as  $\mathcal{T}_2 := \{t \in \mathcal{T} : t > T_0\}$ . The number of post-treatment periods is  $n = T - T_0$ . Let  $m \in \mathbb{N}$  denote the evaluation window length, i.e., the number of periods used to construct the SHC match. Define the evaluation period as the final  $m$  months of the pre-treatment period  $\mathcal{T}_{\text{eval}} := \{T_0 - m + 1, \dots, T_0\} \subset \mathcal{T}_1$ .

Let  $\mathbf{y} = (y_1, y_2, \dots, y_T)^\top \in \mathbb{R}^T$  denote the observed outcome vector for the treated unit.

Define the pre-treatment outcome vector  $\mathbf{y}_{\text{pre}} := \mathbf{y}_{\mathcal{T}_1} \in \mathbb{R}^{T_0}$ , the evaluation subvector  $\mathbf{y}_{\text{eval}} := \mathbf{y}_{\mathcal{T}_{\text{eval}}} \in \mathbb{R}^m$ , and the post-treatment outcome vector  $\mathbf{y}_{\text{post}} := \mathbf{y}_{\mathcal{T}_2} \in \mathbb{R}^n$ .

The evaluation subvector  $\mathbf{y}_{\text{eval}}$  is the target pattern to be reconstructed using convex combinations of earlier segments from the pre-treatment period. To reconstruct the evaluation window, we compare it against earlier segments of the treated unit's own pre-treatment trajectory.

Define the donor matrix  $\tilde{\mathbf{Y}}_{\text{pre}} \in \mathbb{R}^{m \times N}$  as a collection of  $N$  overlapping length- $m$  subvectors extracted from the smoothed pre-treatment series. Each column is a contiguous subvector of the form  $\tilde{\mathbf{y}}_{[i, i+m-1]}$ , with  $i$  ranging over eligible start points in  $\mathcal{T}_1$  that precede the evaluation window. The precise construction is given in [Section 2.2](#).

### 2.1. Synthetic Historical Control Method

**Potential Outcomes** Following the Rubin potential outcomes framework, let  $y_t^1$  and  $y_t^0$  denote the potential outcomes for the treated unit at time  $t \in \mathcal{T}$ , corresponding to the outcomes under treatment and no treatment, respectively. The observed outcome at time  $t$  is  $y_t = d_t y_t^1 + (1 - d_t) y_t^0$ , where  $d_t = \mathbf{1}\{t > T_0\}$ .

The fundamental problem of causal inference is that for  $t > T_0$ , we observe only the treated potential outcome  $y_t^1 = y_t$ , while the untreated potential outcome  $y_t^0$  remains unobserved. Let the SHC estimator for  $y_t^0$  be denoted  $\hat{y}_t^0 := y_t^{\text{SHC}}$  for all  $t \in \mathcal{T}_2$ . Define the observed post-treatment outcome vector as  $\mathbf{y}_{\text{post}} := (y_t^1)_{t \in \mathcal{T}_2}$  and the estimated counterfactual vector as  $\mathbf{y}_{\text{post}}^0 := (\hat{y}_t^0)_{t \in \mathcal{T}_2}$ . The out-of-sample treatment effect is

$$\alpha_t := y_t^1 - y_t^0, \quad \text{for } t \in \mathcal{T}_2, \quad (2.1)$$

with estimator  $\hat{\alpha}_t := y_t - \hat{y}_t^0$ . Stacking over  $t \in \mathcal{T}_2$  yields:

$$\hat{\boldsymbol{\alpha}} := \mathbf{y}_{\text{post}} - \mathbf{y}_{\text{post}}^0. \quad (2.2)$$

The average treatment effect on the treated (ATT) is then given by:

$$\widehat{\text{ATT}} := \frac{1}{n} \sum_{t=T_0+1}^{T_0+n} (y_t - \hat{y}_t^0) = \frac{1}{n} \mathbf{1}^\top \hat{\boldsymbol{\alpha}}, \quad (2.3)$$

where  $\mathbf{1} \in \mathbb{R}^n$  is a vector of ones.

### 2.2. The Model

The SHC method introduced by [Chen et al. \(2024\)](#) constructs a counterfactual using the treated unit's own pre-treatment time series.<sup>2</sup> SHC requires a few assumptions: [Assumption 2.1](#), [Assumption 2.2](#), and [Assumption 2.3](#).

<sup>2</sup>SCM estimates the counterfactual by solving:

$$\mathbf{w}^* = \underset{\mathbf{w} \in \Delta^{N_0}}{\text{argmin}} \|\mathbf{y}_1 - \mathbf{Y}_0 \mathbf{w}\|_2^2, \quad \forall t \in \mathcal{T}_1, \quad (2.4)$$

where  $\mathbf{Y}_0 \in \mathbb{R}^{T_0 \times N_0}$  is the donor matrix of unexposed units and  $\Delta^{N_0} = \{\mathbf{w} \in \mathbb{R}_{\geq 0}^{N_0} : \|\mathbf{w}\|_1 = 1\}$  is the probability simplex.

ASSUMPTION 2.1. (ERROR PROPERTIES) *The error term satisfies  $\mathbb{E}[\varepsilon_t] = 0$  and  $\mathbb{E}[\varepsilon_t^2] < \infty$  for all  $t \in \mathcal{T}$ .*

Assumption 2.1 says unmodeled fluctuations or noise in the outcome are random.

ASSUMPTION 2.2. (LATENT COMPONENT SMOOTHNESS AND MATCHING CONDITION) *The latent trend  $\tilde{y}(t)$  is  $(H + 1)$  times differentiable for some integer  $H \geq 3$ , with derivative bounded as  $|\tilde{y}^{(H+1)}(t)| < \varepsilon$  uniformly over  $t \in \mathcal{T}_1$ . Additionally,  $\exists \mathbf{w}^* \in \Delta^{N-1}$  such that  $\mathbf{y}_{eval} = \tilde{\mathbf{Y}}_{pre} \mathbf{w}^*$ .*

Assumption 2.2 allows us to approximate the post-treatment latent component using a local linear extrapolation of pre-treatment values and posits that perfect match exists within the donor pool for the evaluation window, much as Abadie et al. (2010) assume about donor pool units matching to the treated unit.

ASSUMPTION 2.3. (FEASIBILITY) *There exists a weight vector  $\mathbf{w} \in \Delta^{N-1}$  such that  $\hat{\mathbf{y}}_{eval} = \tilde{\mathbf{Y}}_{pre} \mathbf{w}$ , with  $\|\mathbf{w}\|_0 \leq s \ll N$ . Let  $\tilde{\mathbf{Y}}_{pre}^{(s)}$  denote the submatrix formed by the active columns of  $\tilde{\mathbf{Y}}_{pre}$  indexed by  $\text{supp}(\mathbf{w})$ ; then its Gram matrix satisfies  $(\tilde{\mathbf{Y}}_{pre}^{(s)})^\top \tilde{\mathbf{Y}}_{pre}^{(s)} \succ 0$ .*

Chen et al. (2024) note that Assumption 2.3 is the sample-variant of Assumption 2.2. It says that the SHC weights are identifiable, sparse, and numerically stable, similar to the assumptions described by Abadie (2021).

The first step of the SHC method smooths  $\mathbf{y}_{pre}$  using local linear regression, where we fit the regression model:

$$y_j \approx a_t + b_t(j - t), \quad \text{for } j = 1, \dots, T_0, \quad (2.5)$$

with local intercept  $a_t \in \mathbb{R}$  and slope  $b_t \in \mathbb{R}$ . Define the centered time vector  $\mathbf{s}_t = (1 - t, 2 - t, \dots, T_0 - t)^\top \in \mathbb{R}^{T_0}$ , and the design matrix  $\mathbf{X}_t = [\mathbf{1} \ \mathbf{s}_t] \in \mathbb{R}^{T_0 \times 2}$ . The diagonal kernel weight matrix is:

$$\mathbf{W}_t = \text{diag}(w_{1t}, w_{2t}, \dots, w_{T_0t}), \quad w_{jt} \propto \exp\left(-\frac{1}{2} \left(\frac{j - t}{h}\right)^2\right), \quad (2.6)$$

where  $(h > 0)$  is a bandwidth parameter governing smoothness. The local coefficients  $\boldsymbol{\beta}_t = (a_t, b_t)^\top$  solve quadratic programming problem that is strictly convex

$$\boldsymbol{\beta}_t = \underset{\boldsymbol{\beta} \in \mathbb{R}^2}{\text{argmin}} \left\| \mathbf{W}_t^{1/2} (\mathbf{y}_{pre} - \mathbf{X}_t \boldsymbol{\beta}) \right\|_2^2. \quad (2.7)$$

Fortunately, this optimization admits a closed-form solution for the coefficients:

$$\boldsymbol{\beta}_t = (\mathbf{X}_t^\top \mathbf{W}_t \mathbf{X}_t)^{-1} \mathbf{X}_t^\top \mathbf{W}_t \mathbf{y}_{pre}. \quad (2.8)$$

The smoothed outcome at time  $t$  is  $\tilde{y}_t = a_t = \mathbf{e}_1^\top \boldsymbol{\beta}_t$ , where  $\mathbf{e}_1 = (1, 0)^\top$ . Let  $\tilde{\mathbf{y}} = (\tilde{y}_1, \dots, \tilde{y}_{T_0})^\top \in \mathbb{R}^{T_0}$  denote the full smoothed series. After we have smoothed over the pre-treatment period before the evaluation period, we build our donor matrix of historical control segments/units.

In the second step, SHC constructs  $N = T_0 - m - n + 1$  overlapping temporal segments of length  $m$ . Each segment  $\tilde{\mathbf{y}}_{[i, i+m-1]}$  is a contiguous subvector of the smoothed pre-treatment series. These comprise the donor matrix  $\tilde{\mathbf{Y}}_{\text{pre}} := [\tilde{\mathbf{y}}_{[1, m]} \quad \tilde{\mathbf{y}}_{[2, m+1]} \quad \cdots \quad \tilde{\mathbf{y}}_{[N, N+m-1]}] \in \mathbb{R}^{m \times N}$ .

**EXAMPLE 2.1. (CONSTRUCTING  $\tilde{\mathbf{Y}}_{\text{pre}}$ )** Suppose we have  $T_0 = 60$  months of pre-treatment data and we wish to construct donor segments of length  $m = 12$  months in order to predict a post-treatment horizon of  $n = 6$  months. The number of eligible segments is:  $N = T_0 - m - n + 1 = 60 - 12 - 6 + 1 = 43$ . Each donor segment is a vector of 12 consecutive smoothed pre-treatment outcomes.

In the third step, we solve the following program to obtain the optimal weights:

$$\begin{aligned} \mathbf{w}^* = \underset{\mathbf{w} \in \mathbb{R}^N}{\operatorname{argmin}} \quad & \left\| \tilde{\mathbf{y}}_{\text{eval}} - \tilde{\mathbf{Y}}_{\text{pre}} \mathbf{w} \right\|_2^2 + \varsigma \left\| \mathbf{C}_0^\top \mathbf{w} \right\|_2^2 \\ \text{subject to} \quad & \mathbf{w} \geq \mathbf{0}, \quad \mathbf{1}^\top \mathbf{w} = 1 \end{aligned} \quad (2.9)$$

Chen et al. (2024) note that in the general case when we have more donor units than we do the length of the segments ( $N > m$ ), the optimization becomes an ill-posed problem. The regularization term stabilizes the optimization by penalizing weight allocations along directions of low variance in the donor pool, reducing sensitivity to collinear segments. Formally,  $\mathbf{C}_0$  contains the eigenvectors of the Grammian matrix  $\mathbf{G} \in \mathbb{R}^{N \times N}$ .  $\mathbf{C}_0$  corresponds to the eigenvalues less than  $\tau > 0$ . These directions capture low-variance combinations of donor segments, and the penalty shrinks weights in those directions.<sup>3</sup> The corresponding weight vector  $\mathbf{w}^*$  is used to compute the counterfactual by applying it to the shifted donor segments:

$$\tilde{\mathbf{Y}}_{\text{post}} = [\tilde{\mathbf{y}}_{[1+m, m+n]} \quad \tilde{\mathbf{y}}_{[2+m, m+n+1]} \quad \cdots \quad \tilde{\mathbf{y}}_{[N+m, N+m+n-1]}] \in \mathbb{R}^{n \times N}. \quad (2.10)$$

This yields in-sample and out-of-sample predictions for the SHC estimator. The full counterfactual vector is then:  $\mathbf{y}^0 = \tilde{\mathbf{Y}} \mathbf{w}^* \in \mathbb{R}^{m+n}$ .

**Bandwidth Selection via Cross-Validation** To select the bandwidth parameter  $h$  used in local linear smoothing of  $\mathbf{y}_{\text{pre}}$ , I implement a *leave-one-out cross-validation* (LOOCV) procedure over a grid of candidate values  $h \in [h_{\min}, h_{\max}] \subset \mathbb{R}_{>0}$ . This procedure chooses the bandwidth that minimizes out-of-sample prediction error for the smoothed pre-treatment series  $\tilde{\mathbf{y}} \in \mathbb{R}^{T_0}$ , as defined in Equation~(2.8).

For each candidate  $h$ , and for each time point  $t \in \mathcal{T}_1$ , define Gaussian kernel weights

$$w_{\tau t}^{(h)} = \exp \left( -\frac{1}{2} \left( \frac{\tau - t}{h} \right)^2 \right), \quad \tau \in \mathcal{T}_1 \setminus \{t\}, \quad (2.11)$$

used to construct the diagonal matrix  $\mathbf{W}_t^{(-t)} \in \mathbb{R}^{(T_0-1) \times (T_0-1)}$ , omitting the index  $t$ . Let  $\mathbf{X}_t^{(-t)} \in \mathbb{R}^{(T_0-1) \times 2}$  and  $\mathbf{y}_{\text{pre}}^{(-t)} \in \mathbb{R}^{T_0-1}$  denote the design matrix and response vector with time  $t$  excluded.

<sup>3</sup>The  $\varsigma$  (sigma) term is a small positive constant used to stabilize the solution in cases of near multicollinearity.

Then the local linear coefficients  $\beta_t^{(-t)} = (a_t^{(-t)}, b_t^{(-t)})^\top$  are obtained by solving Equation (2.7). The leave-one-out smoothed value at time  $t$  is then:

$$\tilde{y}_t^{(-t)} = a_t^{(-t)} = \mathbf{e}_1^\top \beta_t^{(-t)}, \quad \text{where } \mathbf{e}_1 = (1, 0)^\top. \quad (2.12)$$

To pick the ideal bandwidth, we minimize:

$$h^* = \underset{h \in [h_{\min}, h_{\max}]}{\operatorname{argmin}} \left\{ \operatorname{CV}(h) := \frac{1}{T_0} \sum_{t=1}^{T_0} \left( y_t - \tilde{y}_t^{(-t)} \right)^2 \right\}. \quad (2.13)$$

Once selected, the optimal bandwidth  $h^*$  is used to construct the full smoothed pre-treatment series.

**Control Group Selection** Chen et al. (2024) note that we should have a parsimonious approach to which historical controls we use, rather than choosing all of them. In their examples, SHC typically requires less than 28 controls. They advocate for a forward selection procedure to choose the optimal control group. I implement a slightly modified variant of this method to choose the ideal number of historical controls. Let  $\mathcal{N} = \{1, 2, \dots, N\}$  index the full set of donor segments in the pre-treatment matrix  $\tilde{\mathbf{Y}}_{\text{pre}} \in \mathbb{R}^{m \times N}$ . At each step the algorithm selects the donor index  $i_j \in \mathcal{N} \setminus \mathcal{S}_{j-1}$  that, when added to the current donor set  $\mathcal{S}_{j-1}$ , minimizes the in-sample mean squared error (MSE) between the evaluation window and in-sample fit:

$$\mathcal{S}_j = \mathcal{S}_{j-1} \cup \left\{ \underset{i \in \mathcal{N} \setminus \mathcal{S}_{j-1}}{\operatorname{argmin}} \left\| \tilde{\mathbf{y}}_{\text{eval}} - \tilde{\mathbf{Y}}_{\text{pre}}^{(\mathcal{S}_{j-1} \cup \{i\})} \mathbf{w}^{(j)} \right\|_2^2 + \varsigma \left\| \mathbf{C}_0^\top \mathbf{w} \right\|_2^2 \right\}, \quad \mathcal{S}_0 = \emptyset, \quad (2.14)$$

where  $\tilde{\mathbf{Y}}_{\text{pre}}^{(\mathcal{S})}$  denotes the submatrix of  $\tilde{\mathbf{Y}}_{\text{pre}}$  with columns indexed by  $\mathcal{S}$ , and the weights  $\mathbf{w}^{(j)} \in \Delta^j$  are the optimal simplex-constrained weights for the selected donor set  $\mathcal{S}_j$ . This procedure greedily expands the donor set one segment at a time, always choosing the segment that yields the greatest marginal reduction in in-sample error.

To avoid overfitting, Chen et al. (2024) manually choose the number of donors the forward selection procedure should choose. In their case, they experiment with the single best historical segment (which rarely does well), the best 15, 20, and 27 historical control units. However, this is a subjective decision that will yield different results depending on what we select. To mitigate researcher degrees of freedom, I view the number of historical segments as a tuning parameter. I employ a forward selection-based algorithm to choose the number of controls. For my implementation of SHC, I employ an early stopping rule based on a modified Bayesian information criterion (BIC), borrowing inspiration from Shi and Huang (2023). For each donor set size  $j$ , I compute:

$$\operatorname{BIC}(j) = m \cdot \log(\operatorname{MSE}_j) + \lambda j, \quad (2.15)$$

where  $\operatorname{MSE}_j$  is the in-sample error using donor set  $\mathcal{S}_j$ , and  $\lambda = \log(m)$  penalizes the inclusion of too many donors. The selection procedure terminates when  $\operatorname{BIC}(j)$  increases for two consecutive steps. The final donor set  $\mathcal{S}^*$  is then taken to be  $\mathcal{S}_{j^*}$ , where:

$$j^* = \underset{j}{\operatorname{argmin}} \operatorname{BIC}(j). \quad (2.16)$$

This approach adaptively selects a sparse donor subset that closely matches the treated

unit's pre-intervention trajectory. The selected donor set  $\mathcal{S}^*$  is used to compute the SHC model predictions. I would like to mention that this forward selection method holds much in common with recently proposed methods from the literature. For example, [Shi and Huang \(2023\)](#) propose the forward selected panel data approach which selects the comparison group for the panel data approach by [Hsiao et al. \(2012\)](#) using an aggressive forward selection procedure. [Li \(2024\)](#), correcting for overfitting in the former, proposed the forward Difference-in-Differences estimator, where a forward selection approach is used to select the donor pool for the DiD model. Similar advances have been made SCM, as developed by [Cerulli \(2024\)](#), where a similar selection mechanism is used to estimate the treatment effect. It would be interesting, but is far outside the scope of this work, to compare these estimators to one another and see when their in-sample fits and point estimates agree with one another.

**Inference** Quantifying uncertainty around the counterfactual predictions is essential for valid inference. [Chen et al. \(2024\)](#) do not provide any inferential method to use aside from in-time placebos. Following recent work in post-modeling inference by [Cattaneo et al. \(2025\)](#), I construct agnostic conformal prediction intervals around the counterfactual series. These intervals are distribution-free, require no parametric assumptions, and are valid under exchangeability of residuals between the pre- and post-treatment periods (or, good in-sample fit).

Let  $\mathbf{y}_{\text{eval}}^0 \in \mathbb{R}^m$  denote the SHC-fitted values during the evaluation period and define the residuals as:

$$\boldsymbol{\varepsilon}_{\text{eval}} := \mathbf{y}_{\text{eval}} - \mathbf{y}_{\text{eval}}^0. \quad (2.17)$$

I then compute the sample mean  $\bar{\varepsilon}$  and variance  $\hat{\sigma}^2$  of these residuals:

$$\bar{\varepsilon} = m^{-1} \sum_{t \in \mathcal{T}_{\text{eval}}} \varepsilon_t, \quad \hat{\sigma}^2 = (m-1)^{-1} \sum_{t \in \mathcal{T}_{\text{eval}}} (\varepsilon_t - \bar{\varepsilon})^2. \quad (2.18)$$

To construct prediction intervals for the post-treatment counterfactual  $\mathbf{y}_{\text{post}}^0 \in \mathbb{R}^n$ , I assume the residuals are sub-Gaussian and apply a conformal bound based on Hoeffding-type concentration inequalities. For a miscoverage rate  $\alpha \in (0, 1)$ , the interval half-width is given by:

$$c_\alpha = \sqrt{2\hat{\sigma}^2 \log \left( \frac{2}{\alpha} \right)}. \quad (2.19)$$

Then, the agnostic  $(1 - \alpha)$  prediction interval for each  $t \in \mathcal{T}_2$  is given by:

$$\left[ \hat{y}_t^0 + \bar{\varepsilon} - c_\alpha, \hat{y}_t^0 + \bar{\varepsilon} + c_\alpha \right]. \quad (2.20)$$

Stacking these yields lower and upper bound vectors:

$$\mathbf{L} := \mathbf{y}_{\text{post}}^0 + \bar{\varepsilon} \cdot \mathbf{1} - c_\alpha \cdot \mathbf{1}, \quad \mathbf{U} := \mathbf{y}_{\text{post}}^0 + \bar{\varepsilon} \cdot \mathbf{1} + c_\alpha \cdot \mathbf{1}, \quad (2.21)$$

which define the conformal confidence band around the counterfactual trajectory.

This approach allows finite-sample valid coverage under the assumption that the pre- and post-treatment residuals are exchangeable. Intuitively, if the SHC model fits the pre-treatment series well (i.e., residuals are small and centered), then its extrapolations into

the post-treatment period are expected to carry similarly bounded uncertainty. In this sense, the residual distribution learned from the evaluation window is used to calibrate uncertainty for the out-of-sample outcomes.

These prediction intervals reflect uncertainty in the estimated counterfactual. Hence, if the observed outcome falls far outside the prediction interval, this is interpreted as evidence of a treatment effect. Unlike traditional confidence intervals around treatment effect estimates (e.g., using standard errors), the conformal intervals here do not assume a data-generating model. I use a default coverage level of 90% (i.e.,  $\alpha = 0.1$ ) in all applications.

### 3. DATA

To evaluate the impact of Hawaii’s COVID-19 border closure on its tourism economy, I use monthly time series spanning multiple economic dimensions, focusing on four primary outcomes: two related to tourism demand and two capturing broader economic impacts. All outcomes are transformed into annualized year-over-year (YoY) growth rates. This transformation mitigates nonstationary trends—crucial for the validity of the SHC estimator [Chen et al. \(2024\)](#)—and allows treatment effects to be interpreted directly as percentage point changes.

The first two outcomes capture the demand side of Hawaii’s tourism sector. *Visitor Days* measures the total number of person-days spent by visitors in a given month, integrating both the volume of arrivals and their length of stay. *Occupancy* refers to the statewide hotel occupancy rate, defined as the percentage of available hotel rooms that are occupied. Together, these indicators provide complementary measures of tourism demand: the former captures total visitor presence, while the latter reflects realized capacity utilization across the hospitality sector. Both series are sourced from Hawaii’s Department of Business, Economic Development & Tourism (DBEDT) and span from January 1990 to December 2020.

The remaining outcomes capture economic effects, one sector-specific and one broader in scope. *Accommodation Employment* tracks the number of employees in Hawaii’s accommodation industry, a labor-intensive sector directly tied to tourism activity. *The Coincident Economic Activity Index* summarizes statewide macroeconomic conditions based on four indicators: nonfarm payroll employment, the unemployment rate, average weekly hours worked in manufacturing, and wage and salary disbursements. This index is benchmarked to match the trend of gross state product, making it a comprehensive indicator of the state’s economic health. Both economic indicators are sourced from the Federal Reserve Economic Data (FRED) system and cover January 1991 to December 2020.

Hawaii’s mandatory quarantine policy took effect in late March 2020. I define the intervention to begin in March 2020, corresponding to time  $t = 362$ , and formalize the treatment indicator as:

$$d_t := \mathbf{1}\{t \geq 362\},$$


so that  $d_t = 1$  for all months after the policy’s implementation and  $d_t = 0$  otherwise. The




final pre-treatment period is February 2020, yielding a pre-treatment sample of  $T_0 = 361$  months.

#### 4. RESULTS

In Figure we visualize the outcomes of interest over time from January of 1991 to December 2020. We can see that the growth rate trends for all outcomes were relatively smooth across the full pre-intervention series, exhibiting seasonality and relatively low variance. We also see that the outcomes of interest precipitously dropped when the mandatory quarantine policy was implemented (aside from unemployment claims). The plot also suggests that [Assumption 2.1](#) and [Assumption 2.2](#) may be valid.



Here are the initial SHC results. For these results,  $m = 36$  (3 years of evaluation data), but the results remain robust up until 8 years of evaluation data.



The plot reports the ATT and pre-treatment RMSE for each outcome of interest. Between March and December 2020, Hawaii experienced an 86.1 percentage point decline in the annualized growth rate of visitor days, relative to its synthetic historical control. This estimate reflects a near-complete collapse in tourism activity and far exceeds typical year-over-year fluctuations (pre-treatment RMSE = 1.1%).

Hotel occupancy rates declined by 76.7 percentage points in annualized growth, indicating a severe contraction in the hotel sector. Given the relative stability of pre-treatment occupancy rates, this represents a highly atypical deviation, unlikely to be explained by seasonal or cyclical variation.


Employment in the leisure and hospitality sector fell by 48.5 percentage points in year-over-year growth terms. While labor market indicators are typically more volatile, the pre-treatment RMSE of 0.4% underscores the magnitude of this decline. These job losses were concentrated in a sector that comprises a substantial share of Hawaii's economy, reinforcing the interpretation of the border closure as an intentional economic suspension.

Initial unemployment claims rose by 762.5 percentage points year-over-year, relative to expected levels. That is, claims were more than eight times higher than would be predicted under the counterfactual. This represents a sudden and extreme labor market shock, and the stability of this ATT across more extended evaluation windows (up to eight years) provides further evidence that the effect is both robust and attributable to the policy.

Even accounting for finite-sample prediction uncertainty, the observed collapse in tourism is extreme relative to the synthetic historical distribution. The observed values fall far

outside of the 90% conformal prediction intervals, suggesting that the results are far outside of what we'd expect under a no-quarantine scenario.

Next we do in-time placebos. I backdate the treatment to June of 2019, and predict until December 2019.



To assess the credibility of our identification strategy, we conduct a placebo test by assigning a false treatment date of June 2019 and evaluating outcomes through December 2019. Because this window well predates the onset of the COVID-19 pandemic, a well-specified model should yield no substantial treatment effects. Across all outcomes, estimated effects are indeed close to zero. For Visitor Days, the ATT is  $-0.000$  with an RMSE of 0.014. Hotel Occupancy shows an ATT of 0.030 and an RMSE of 0.011. For Leisure and Hospitality Employment, the ATT is  $-1.282$  with an RMSE of 0.504. Initial Unemployment Claims yield an ATT of  $-5.46$  and an RMSE of 5.8.

These placebo estimates support the model's internal validity. The tourism variables, in particular, show excellent pre-treatment fit and effectively zero estimated impact during the placebo period, as expected. The labor market outcomes, by contrast, exhibit noisier placebo effects and visibly weaker pre-treatment fit. This limitation stems from the fact that the SHC estimator relies on interpolation in principle: it can only construct counterfactuals within the convex hull of past time segments. As such, if the evaluation window lacks similar past dynamics, predictive accuracy may suffer.

Even so, the conformal prediction intervals for this placebo test largely cover the observed outcomes. This suggests that any discrepancies between prediction and the observed

values lie within expected model error. Altogether, the results of this falsification test reinforce confidence in the causal interpretation of the main findings.

## REFERENCES

- Abadie, A. (2021, 06). Using synthetic controls: Feasibility, data requirements, and methodological aspects. *Journal of Economic Literature* 59(2), 391–425.
- Abadie, A., A. Diamond, and J. Hainmueller (2010). Synthetic control methods for comparative case studies: Estimating the effect of california’s tobacco control program. *Journal of the American Statistical Association* 105(490), 493–505.
- Bilgel, F. (2022, 05). Effects of covid-19 lockdowns on social distancing in turkey. *The Econometrics Journal* 25(3), 781–805.
- Breidenbach, P. and T. Mitze (2021, 07). Large-scale sport events and covid-19 infection effects: evidence from the german professional football ‘experiment’. *The Econometrics Journal* 25(1), 15–45.
- Bulle, B. G., D. Shen, D. Shah, and A. E. Hosoi (2022). Public health implications of opening national football league stadiums during the covid-19 pandemic. *Proceedings of the National Academy of Sciences* 119(14), e2114226119.
- Callaway, B. and T. Li (2023). Policy evaluation during a pandemic. *Journal of Econometrics* 236(1), 105454.
- Cattaneo, M. D., Y. Feng, F. Palomba, and R. Titiunik (2025). scpi: Uncertainty quantification for synthetic control methods. *Journal of Statistical Software* 113(1), 1–38.
- Cerulli, G. (2024). Optimal initial donor selection for the synthetic control method. *Economics Letters* 244, 111976.
- Chen, Y.-T., J.-C. Yang, and T.-T. Yang (2024, September). Synthetic historical control for policy evaluation. Available at SSRN: <https://ssrn.com/abstract=4995085>.
- Friedson, A. I., D. McNichols, J. J. Sabia, and D. Dave (2021). Shelter-in-place orders and public health: Evidence from california during the covid-19 pandemic. *Journal of Policy Analysis and Management* 40(1), 258–283.
- Gibson, J. and X. Sun (2023). A synthetic control analysis of us state level covid-19 stay-at-home orders on new jobless claims. *Journal of Economics and Finance* 47(1), 1–14.
- Hsiao, C., H. Steve Ching, and S. Ki Wan (2012). A panel data approach for program evaluation: Measuring the benefits of political and economic integration of hong kong with mainland china. *Journal of Applied Econometrics* 27(5), 705–740.
- Ivanova, I. (2020, May). Nearly 30% of hawaii’s workforce filed jobless claims. here’s a state-by-state breakdown of unemployment woes. *CBS News*. Accessed: 2025-07-14.
- Ke, X. and C. Hsiao (2022). Economic impact of the most drastic lockdown during covid-19 pandemic—the experience of hubei, china. *Journal of Applied Econometrics* 37(1), 187–209.
- Ke, X. and C. Hsiao (2023). Data subject to multiple treatment effects — disentangle the impacts of global pandemic and a specific disease control policy. *The Singapore Economic Review* 68(05), 1507–1527.
- Li, K. T. (2024). Frontiers: A simple forward difference-in-differences method. *Marketing Science* 43(2), 239–468.
- Shi, Z. and J. Huang (2023). Forward-selected panel data approach for program evaluation. *Journal of Econometrics* 234(2), 512–535.
- Yang, X. (2021). Does city lockdown prevent the spread of covid-19? new evidence from the synthetic control method. *Global Health Research and Policy* 6, 1–14.