

Tensor categories

Jonathan Gruber

January 30, 2023

Introduction

In many fields of mathematics, one is naturally led to study tensor products of certain objects (e.g. sheaves in algebraic geometry, cobordisms in topology, modules in commutative algebra). All of these notions fit into the framework of *monoidal categories*, which gives an abstract definition of what a tensor product structure on a category should be. The aim of this course is to give an introduction to monoidal categories and tensor categories, the latter being certain monoidal categories endowed with some extra structure. (They are abelian and have a linear structure that is compatible with the tensor product, over some field.)

The first half of the course deals mainly with category-theoretical notions, starting from the definition of a monoidal category and then discussing important additional properties such as rigidity (the existence of duals) and braidings (functorial isomorphisms $X \otimes Y \cong Y \otimes X$). In the second half, we will turn our attention to tensor categories. Our prime example is the category of $\mathbf{Rep}_{\mathbb{k}}(G)$ finite-dimensional representations of a group G over a field \mathbb{k} , and we will discuss reconstruction theorems that allow us to recover a group (or Hopf algebra) from the corresponding category of representations, together with its monoidal structure and the *forgetful functor* that sends a representation to the underlying vector space. This gives rise to bijections between certain types of groups and Hopf algebras, up to isomorphism, and certain kinds of tensor categories, up to monoidal equivalence, which are broadly referred to as *Tannaka duality*.

Author's note

These notes are the my own synopsis of material that has been collected from many different sources, but most importantly, from [EGNO15]. Further important references include [ML98, DM82, EGNO09] and some websites such as nLab, MathStackExchange, and Wikipedia. No originality is claimed, except in the presentation of the material, and all mistakes should be considered my responsibility.

These notes are also a work in progress. If you find any mistakes or typos and if you have comments or suggestions, please let me know.

I would like to thank Johannes Flake for helpful discussions and encouragement, and for making his own notes available.

1 Monoidal categories

Definition 1.1. A *category* \mathcal{C} consists of the following data:

- (1) a class $\mathrm{Ob}(\mathcal{C})$ of *objects* of \mathcal{C} ;
- (2) for every pair of objects $X, Y \in \mathrm{Ob}(\mathcal{C})$, a set $\mathrm{Hom}_{\mathcal{C}}(X, Y)$ of *homomorphisms* from A to B ;
- (3) for every triple of objects $X, Y, Z \in \mathrm{Ob}(\mathcal{C})$, a *composition map*

$$\circ: \mathrm{Hom}_{\mathcal{C}}(Y, Z) \times \mathrm{Hom}_{\mathcal{C}}(X, Y) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(X, Z)$$

such that the following axioms hold:

(a) for $X, Y, Z, W \in \text{Ob}(\mathcal{C})$ and homomorphisms $f: X \rightarrow Y$, $g: Y \rightarrow Z$, $h: Z \rightarrow W$, we have

$$(h \circ g) \circ f = h \circ (g \circ f).$$

(b) for $X \in \text{Ob}(\mathcal{C})$, there exists an *identity homomorphism* id_X such that $\text{id}_X \circ f = f$ and $g \circ \text{id}_X = g$ for all $Y \in \text{Ob}(\mathcal{C})$ and homomorphisms $f: Y \rightarrow X$ and $g: X \rightarrow Y$.

Remark 1.2. (1) The homomorphisms in a category are often simply referred to as *morphisms*.

(2) As in points (a) and (b) of the definition, we often denote a homomorphism $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ by an arrow $f: X \rightarrow Y$.

(3) A morphism $f: X \rightarrow Y$ is called an *isomorphism* if there exists a morphism $g: Y \rightarrow X$ such that $f \circ g = \text{id}_Y$ and $g \circ f = \text{id}_X$. In that case, we write $g = f^{-1}$ and $X \cong Y$.

Example 1.3. We list some important examples of categories:

- **Set**: the category of sets, with maps between sets as homomorphisms;
- **Grp**: the category of groups with group homomorphisms;
- **AbGrp**: the category of abelian groups with group homomorphisms;
- **Vect $_{\mathbb{k}}$** : the category of finite-dimensional \mathbb{k} -vector spaces with \mathbb{k} -linear maps, for a given field \mathbb{k} ;
- **A -Mod**: the category of A -modules with A -module homomorphisms, for a given algebra A ; we write **A -mod** for the subcategory of finite-dimensional A -modules.

Definition 1.4. A *functor* $F: \mathcal{C} \rightarrow \mathcal{D}$ between categories \mathcal{C} and \mathcal{D} assigns

- (1) to every object $X \in \text{Ob}(\mathcal{C})$ an object $F(X) \in \text{Ob}(\mathcal{D})$;
- (2) to every homomorphism $f: X \rightarrow Y$ in \mathcal{C} a homomorphism $F(f): F(X) \rightarrow F(Y)$ in \mathcal{D} ;

in such a way that

$$F(\text{id}_X) = \text{id}_{F(X)} \quad \text{and} \quad F(f \circ g) = F(f) \circ F(g).$$

Remark 1.5. Given two categories \mathcal{C} and \mathcal{D} , we can form the product category $\mathcal{C} \times \mathcal{D}$ whose objects are the pairs (X, Y) of objects $X \in \text{Ob}(\mathcal{C})$ and $Y \in \text{Ob}(\mathcal{D})$, and where

$$\text{Hom}_{\mathcal{C} \times \mathcal{D}}((X, Y), (Z, W)) = \text{Hom}_{\mathcal{C}}(X, Z) \times \text{Hom}_{\mathcal{D}}(Y, W)$$

for $X, Z \in \text{Ob}(\mathcal{C})$ and $Y, W \in \text{Ob}(\mathcal{D})$. The composition and the identity morphisms are defined component-wise in the obvious way. A functor from $\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ to some other category \mathcal{E} is often called a *bifunctor*.

Definition 1.6. Let \mathcal{C} and \mathcal{D} be categories and let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{C} \rightarrow \mathcal{D}$ be functors. A *natural transformation* $\eta: F \rightarrow G$ is a family of morphisms $\eta_A: F(A) \rightarrow G(A)$ in \mathcal{D} , for every object A of \mathcal{C} , such that for every morphism $f: A \rightarrow B$ in \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc} F(A) & \xrightarrow{\eta_A} & G(A) \\ F(f) \downarrow & & \downarrow G(f) \\ F(B) & \xrightarrow{\eta_B} & G(B) \end{array}$$

For an object A of \mathcal{C} , we call η_A the component of η at A . A natural transformation is called a *natural isomorphism* if all of its components are isomorphisms.

Remark 1.7. The functors between two categories \mathcal{C} and \mathcal{D} form a category $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$ whose morphisms are natural transformations between functors. The composition of natural transformations is defined componentwise.

Definition 1.8. A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is called an equivalence if there exists a functor $G: \mathcal{D} \rightarrow \mathcal{C}$ such that $F \circ G$ is naturally isomorphic to $\text{id}_{\mathcal{D}}$ and $G \circ F$ is naturally isomorphic to $\text{id}_{\mathcal{C}}$.

Definition 1.9. A *monoidal category* is a tuple $(\mathcal{C}, \mathbf{1}, \otimes, \alpha, \lambda, \rho)$, where

- \mathcal{C} is a category,
- $\mathbf{1}$ is an object of \mathcal{C} , called the *unit object*,
- $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is a bifunctor called the *tensor product*,
- $a: - \otimes (- \otimes -) \rightarrow (- \otimes -) \otimes -$ is a natural isomorphism, called the *associativity constraint*,
- $\lambda: \mathbf{1} \otimes - \rightarrow \text{id}_{\mathcal{C}}$ and $\rho: - \otimes \mathbf{1} \rightarrow \text{id}_{\mathcal{C}}$ are natural isomorphisms, called the (left and right) *unitors*,

subject to the following axioms:

Pentagon axiom: For all objects A, B, C, D of \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc}
 A \otimes (B \otimes (C \otimes D)) & \xrightarrow{\text{id}_A \otimes \alpha_{B,C,D}} & A \otimes ((B \otimes C) \otimes D) \\
 \alpha_{A,B,C \otimes D} \swarrow & & \searrow \alpha_{A,B \otimes C,D} \\
 (A \otimes B) \otimes (C \otimes D) & & (A \otimes (B \otimes C)) \otimes D \\
 \alpha_{A \otimes B,C,D} \searrow & & \swarrow \alpha_{A,B,C} \otimes \text{id}_D \\
 & ((A \otimes B) \otimes C) \otimes D &
 \end{array}$$

In other words, we have

$$(\alpha_{A,B,C \otimes D}) \circ (\alpha_{A,B \otimes C,D}) \circ (\text{id}_A \otimes \alpha_{B,C,D}) = (\alpha_{A \otimes B,C,D}) \circ (\alpha_{A,B,C} \otimes \text{id}_D).$$

Unit axiom / triangle axiom: For all objects A, B of \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc}
 A \otimes (\mathbf{1} \otimes B) & \xrightarrow{\alpha_{A,\mathbf{1},B}} & (A \otimes \mathbf{1}) \otimes B \\
 \text{id}_A \otimes \lambda_B \searrow & & \swarrow \rho_A \otimes \text{id}_B \\
 & A \otimes B &
 \end{array}$$

In other words, we have

$$(\rho_A \otimes \text{id}_B) \circ \alpha_{A,\mathbf{1},B} = \text{id}_A \otimes \lambda_B.$$

Remark 1.10. (1) When no confusion is possible, we simply write \mathcal{C} instead of $(\mathcal{C}, \mathbf{1}, \otimes, \alpha, \lambda, \rho)$. We also say that $(\mathbf{1}, \otimes, \alpha, \lambda, \rho)$ is a *monoidal structure* on the category \mathcal{C} .

(2) Being monoidal is **not** a property of a given category, but an *additional structure*. A category can admit more than one monoidal structure. (See the examples below.)

- (3) The fact that \otimes is a bifunctor means that for suitable morphisms a, b, c, d in \mathcal{C} , we have

$$(a \otimes c) \circ (b \otimes d) = (a \circ b) \otimes (c \circ d).$$

- (4) Instead of assuming the existence of $\mathbf{1}$ with natural transformations λ and ρ subject to the unit axiom, one can start from the (seemingly weaker, but actually equivalent) assumptions that there exists an object $\mathbf{1}$ with an isomorphism $\iota: \mathbf{1} \otimes \mathbf{1} \rightarrow \mathbf{1}$ such that the functors $\mathbf{1} \otimes -$ and $- \otimes \mathbf{1}$ are equivalences. (No additional assumptions on the isomorphism ι are necessary.) See Sections 2.1 and 2.2 of [EGNO15] for more details.

Example 1.11. (1) The category **Set** of sets has a monoidal structure where the tensor product is given by the cartesian product and the unit object is a singleton $\{\bullet\}$.

- (2) The category **Grp** of groups has a monoidal structure where the tensor product is given by the cartesian product and the unit object is the trivial group $\{1\}$.
- (3) The category **AbGrp** of abelian groups has a monoidal structure where the tensor product is given by the usual tensor product $- \otimes_{\mathbb{Z}} -$ and the unit object is the group \mathbb{Z} of integers. It also inherits a different monoidal structure from the category **Grp**; see the previous point.
- (4) The category **Vect** $_{\mathbb{k}}$ of vector spaces over a field \mathbb{k} has a monoidal structure where the tensor product is given by the usual tensor product $- \otimes_{\mathbb{k}} -$ and the unit object is the one-dimensional vector space \mathbb{k} .
- (5) The category **Rep** $_{\mathbb{k}}(G)$ of finite dimensional representations of a group G over a field \mathbb{k} admits a monoidal structure where the tensor product is the usual tensor product of representations and the unit object is the trivial one-dimensional representation \mathbb{k} . More precisely, if we identify **Rep** $_{\mathbb{k}}(G)$ with the category of finite-dimensional modules over the group algebra $\mathbb{k}[G]$ then the action of $\mathbb{k}[G]$ on the tensor product $M \otimes N$ of two $\mathbb{k}[G]$ -modules M and N is uniquely determined by $g \cdot (m \otimes n) = gm \otimes gn$ for $g \in G$, $m \in M$ and $n \in N$.

In the following, we refer to the objects of **Rep** (G) as G -modules.

- (6) For a category \mathcal{C} , the category **End** $(\mathcal{C}) = \mathbf{Fun}(\mathcal{C}, \mathcal{C})$ of endofunctors of \mathcal{C} has a monoidal structure, where the tensor product is given by the composition of functors and the unit object is the identity functor $\text{id}_{\mathcal{C}}$. The associativity constraints and unitors are identity natural transformations.

The two next examples will seem quite trivial for now, but they will become more interesting later when we add extra structure:

- (7) Let G be a monoid and A an abelian group. Then we can define a monoidal category \mathcal{C}_A^G with objects $\text{Ob}(\mathcal{C}_A^G) = \{\delta_g \mid g \in G\}$ indexed by G and homomorphisms

$$\text{Hom}_{\mathcal{C}_A^G}(\delta_g, \delta_h) = \begin{cases} A & \text{if } g = h, \\ \emptyset & \text{otherwise,} \end{cases}$$

for $g, h \in G$. The tensor product is defined by $\delta_g \otimes \delta_h = \delta_{gh}$ and by $a \otimes a' = aa' \in A$ for $g, h \in G$ and $a, a' \in A$, the unit object is δ_e (where $e \in G$ is the unit element) and the associativity constraints and unitors are identity maps.

- (8) Let G be a monoid and let **Vect** $_{\mathbb{k}}^G$ be the category of finite-dimensional G -graded \mathbb{k} -vector spaces $V = \bigoplus_{g \in G} V_g$, with homomorphisms given by grading-preserving linear maps. (That is, for $V = \bigoplus_g V_g$ and $W = \bigoplus_g W_g$ two G -graded vector spaces, the homomorphisms from V to W in **Vect** $_{\mathbb{k}}^G$ are the linear maps $f: V \rightarrow W$ that satisfy $f(V_g) \subseteq W_g$ for all $g \in G$.) Then the

monoidal structure on $\mathbf{Vect}_{\mathbb{k}}$ induces a monoidal structure $\mathbf{Vect}_{\mathbb{k}}^G$, where the grading on the tensor product of G -graded vector spaces V and W is given by

$$(V \otimes W)_g = \bigoplus_{hh'=g} V_h \otimes W_{h'}$$

for $g \in G$. The tensor product of two homomorphisms in $\mathbf{Vect}_{\mathbb{k}}^G$ is just the usual tensor product of linear maps. The unit object is the one-dimensional vector space $\mathbb{k} = \mathbb{k}_e$ whose unique non-zero grading piece is indexed by the unit object $e \in G$. The associativity constraint and the unitors come from the category $\mathbf{Vect}_{\mathbb{k}}$.

Observe that there is a faithful functor $i_{\mathbb{k}}^G: \mathcal{C}_{\mathbb{k}^\times}^G \rightarrow \mathbf{Vect}_{\mathbb{k}}^G$ with $i_{\mathbb{k}}^G(\delta_g) = \mathbb{k}_g$ the one-dimensional vector space with grading concentrated in degree g , for $g \in G$, and with the obvious definition on homomorphisms. This functor is compatible with the tensor product (up to a natural isomorphism); it is an example of a *monoidal functor* (to be defined shortly).

In our final example, we demonstrate that there are monoidal categories with a less obvious choice of associativity constraint.

- (9) Let G be a monoid, let A an abelian group and let ω be a 3-cocycle for G with values in A , i.e. a map $\omega: G^{\times 3} \rightarrow A$ with

$$(1.1) \quad \omega(g_1 g_2, g_3, g_4) \omega(g_1, g_2, g_3 g_4) = \omega(g_2, g_3, g_4) \omega(g_1, g_2 g_3, g_4) \omega(g_1, g_2, g_3)$$

for $g_1, g_2, g_3, g_4 \in G$. Then we can define a monoidal category $\mathcal{C}_A^{G, \omega}$ with underlying category \mathcal{C}_A^G and with tensor product and unit object defined as in point (7), but with associativity constraint α^ω defined by

$$\alpha_{g,h,k}^\omega = \omega(g, h, k): \delta_g \otimes (\delta_h \otimes \delta_k) = \delta_{ghk} \longrightarrow \delta_{ghk} = (\delta_g \otimes \delta_h) \otimes \delta_k$$

for $g, h, k \in G$. Observe that the 3-cocycle condition implies that $\mathcal{C}_A^{G, \omega}$ satisfies the pentagon axiom. (In fact, a map $\omega: G^{\times 3} \rightarrow A$ defines an associativity constraint for \mathcal{C}_A^G if and only if ω is a 3-cocycle.) The unitors are defined by $\lambda_g = \omega(e, e, g)$ and $\rho_g = \omega(g, e, e)^{-1}$ for $g \in G$, and the unit axiom becomes the equation $\omega(g, e, h) = \omega(g, e, e) \cdot \omega(e, e, h)$ for $g, h \in G$ (which also follows from (1.1) by setting $g_2 = g_3 = e$).

Given a 3-cocycle $\omega: G^{\times 3} \rightarrow \mathbb{k}^\times$ with values in the multiplicative group of a field \mathbb{k} , we can extend the associativity constraint α^ω on $\mathcal{C}_{\mathbb{k}^\times}^G$ to an associativity constraint α^ω on $\mathbf{Vect}_{\mathbb{k}}^G$ via

$$\alpha_{\mathbb{k}_g, \mathbb{k}_h, \mathbb{k}_k}^\omega = \omega(g, h, k) \cdot \alpha_{\mathbb{k}_g, \mathbb{k}_h, \mathbb{k}_k},$$

for $g, h, k \in G$, extended by additivity, where α denotes the ‘usual’ associativity constraint in $\mathbf{Vect}_{\mathbb{k}}^G$. (Note that every object of $\mathbf{Vect}_{\mathbb{k}}^G$ is a direct sum of objects of the form \mathbb{k}_g with $g \in G$.)

Remark 1.12. Given a monoidal category $\mathcal{C} = (\mathcal{C}, \mathbf{1}, \otimes, \alpha, \lambda, \rho)$, we define the *opposite monoidal category* $\mathcal{C}^{\text{op}} = (\mathcal{C}, \mathbf{1}, \otimes^{\text{op}}, \alpha^{\text{op}}, \rho, \lambda)$ with the same underlying category, but tensor product defined by $X \otimes^{\text{op}} Y = Y \otimes X$ and $f \otimes^{\text{op}} g = g \otimes f$ for objects X, Y and homomorphisms f, g in \mathcal{C} , and with associativity constraint given by $\alpha_{X,Y,Z}^{\text{op}} = \alpha_{Z,Y,X}^{-1}$ for objects X, Y, Z of \mathcal{C} .

This is not to be confused with the *reverse category* \mathcal{C}^{rev} with $\text{Hom}_{\mathcal{C}^{\text{rev}}}(X, Y) = \text{Hom}_{\mathcal{C}}(X, Y)$. The latter can also be endowed with a canonical monoidal structure. Note that \mathcal{C}^{rev} is often also called the opposite category of \mathcal{C} ; we use non-standard terminology here to avoid confusion with the opposite monoidal category defined above.

Lemma 1.13. *For all objects A, B of \mathcal{C} , we have*

$$\rho_{A \otimes B} \circ \alpha_{A, B, \mathbf{1}} = \text{id}_A \otimes \rho_B \quad \text{and} \quad \lambda_{A \otimes B} \circ \alpha_{\mathbf{1}, A, B} = \lambda_A \otimes \text{id}_B.$$

Proof. Consider the following diagram, where all arrows are isomorphisms:

$$\begin{array}{ccccc}
A \otimes (B \otimes (\mathbf{1} \otimes D)) & \xrightarrow{\text{id}_A \otimes \alpha_{B,\mathbf{1},D}} & A \otimes ((B \otimes \mathbf{1}) \otimes D) \\
\downarrow \alpha_{A,B,\mathbf{1} \otimes D} & \searrow \text{id}_A \otimes (\text{id}_B \otimes \lambda_D) & \swarrow \text{id}_A \otimes (\rho_B \otimes \text{id}_D) & \downarrow \alpha_{A,B \otimes \mathbf{1},D} \\
& A \otimes (B \otimes D) & \\
& \downarrow \alpha_{A,B,D} & \\
& (A \otimes B) \otimes D & \\
\swarrow \text{id}_{A \otimes B} \otimes \lambda_D & & \swarrow (\text{id}_A \otimes \rho_B) \otimes \text{id}_D \\
(A \otimes B) \otimes (\mathbf{1} \otimes D) & & (A \otimes (B \otimes \mathbf{1})) \otimes D \\
\searrow \alpha_{A \otimes B,\mathbf{1},D} & \uparrow \rho_{A \otimes B} \otimes \text{id}_D & \searrow \alpha_{A,B,\mathbf{1}} \otimes \text{id}_D \\
& ((A \otimes B) \otimes \mathbf{1}) \otimes D &
\end{array}$$

The external pentagon commutes by the pentagon axiom, the quadrangles commute by naturality of the associativity constraint α , and the top triangle and the bottom left triangle commute by the triangle axiom. Since all arrows are isomorphisms, this implies that the bottom right triangle commutes. Setting $D = \mathbf{1}$ and using the natural isomorphism $\rho: - \otimes \mathbf{1} \rightarrow \text{id}_{\mathcal{C}}$, it follows that the following diagram commutes:

$$\begin{array}{ccc}
A \otimes (B \otimes \mathbf{1}) & \xrightarrow{\alpha_{A,B,\mathbf{1}}} & (A \otimes B) \otimes \mathbf{1} \\
\searrow \text{id}_A \otimes \rho_B & & \swarrow \rho_{A \otimes B} \\
& A \otimes B &
\end{array}$$

This proves the first claim, the second claim can be proven analogously. \square

Definition 1.14. Let $(\mathcal{C}, \mathbf{1}, \otimes, \alpha, \lambda, \rho)$ and $(\mathcal{C}', \mathbf{1}', \otimes', \alpha', \lambda', \rho')$ be two monoidal categories. A monoidal functor from \mathcal{C} to \mathcal{C}' is a triple $(F, \varphi, \varepsilon)$, where $F: \mathcal{C} \rightarrow \mathcal{C}'$ is a functor, $\varphi: F(- \otimes -) \rightarrow F(-) \otimes' F(-)$ is a natural isomorphism and $\varphi: F(\mathbf{1}) \rightarrow \mathbf{1}'$ is an isomorphism, such that the following diagrams commute:

$$\begin{array}{ccc}
F(A \otimes (B \otimes C)) & \xrightarrow{F(\alpha_{A,B,C})} & F((A \otimes B) \otimes C) \\
\searrow \varphi_{A,B \otimes C} & & \searrow \varphi_{A \otimes B,C} \\
F(A) \otimes' F(B \otimes C) & & F(A \otimes B) \otimes' F(C) \\
\searrow \text{id}_{F(A)} \otimes \varphi_{B,C} & & \swarrow \varphi_{A,B} \otimes \text{id}_{F(C)} \\
F(A) \otimes' (F(B) \otimes' F(C)) & \xrightarrow{\alpha'_{F(A),F(B),F(C)}} & (F(A) \otimes' F(B)) \otimes' F(C)
\end{array}$$

$$\begin{array}{ccc}
F(A \otimes \mathbf{1}) & \xrightarrow{\varphi_{A,\mathbf{1}}} & F(A) \otimes' F(\mathbf{1}) \\
\downarrow F(\rho_A) & & \downarrow \text{id}_{F(A)} \otimes' \varepsilon \\
F(A) & \xleftarrow{\rho'_{F(A)}} & F(A) \otimes' \mathbf{1}'
\end{array}
\qquad
\begin{array}{ccc}
F(\mathbf{1} \otimes A) & \xrightarrow{\varphi_{\mathbf{1},A}} & F(\mathbf{1}) \otimes' F(A) \\
\downarrow F(\lambda_A) & & \downarrow \varepsilon \otimes' \text{id}_{F(A)} \\
F(A) & \xleftarrow{\lambda'_{F(A)}} & \mathbf{1}' \otimes' F(A)
\end{array}$$

A *monoidal natural transformation* between monoidal functors $(F, \varphi, \varepsilon)$ and $(F', \varphi', \varepsilon')$ is a natural transformation $\psi: F \rightarrow F'$ such that

$$\varphi' \circ \psi_{-\otimes-} = \psi \otimes' \psi \circ \varphi \quad \text{and} \quad \varepsilon = \varepsilon' \circ \psi_{\mathbf{1}},$$

i.e. the following diagrams commute for all objects A, B of \mathcal{C} :

$$\begin{array}{ccc}
F(A \otimes B) & \xrightarrow{\varphi_{A,B}} & F(A) \otimes' F(B) \\
\downarrow \psi_{A \otimes B} & & \downarrow \psi_A \otimes' \psi_B \\
F'(A \otimes B) & \xrightarrow{\varphi'_{A,B}} & F'(A) \otimes' F'(B)
\end{array}
\qquad
\begin{array}{ccc}
F(\mathbf{1}) & \xrightarrow{\psi_{\mathbf{1}}} & F'(\mathbf{1}) \\
& \searrow \varepsilon & \downarrow \varepsilon' \\
& & \mathbf{1}'
\end{array}$$

Remark 1.15. (1) Being monoidal for a functor is an additional structure, and not a property. However, being monoidal for a natural transformation is a property.

(2) The composition of two monoidal functors $(F, \varphi, \varepsilon): \mathcal{C} \rightarrow \mathcal{D}$ and $(F', \varphi', \varepsilon'): \mathcal{D} \rightarrow \mathcal{E}$ can be considered as a monoidal functor with structure maps defined as follows, for objects X, Y of \mathcal{C} :

$$\begin{aligned}
F'(F(X \otimes Y)) &\xrightarrow{F'(\varphi_{X,Y})} F'(F(X) \otimes F(Y)) \xrightarrow{\varphi'_{F(X), F(Y)}} F'(F(X)) \otimes' F'(F(Y)) \\
F'(F(\mathbf{1})) &\xrightarrow{F'(\varepsilon)} F'(\mathbf{1}) \xrightarrow{\varepsilon'} \mathbf{1}.
\end{aligned}$$

(3) Given a monoidal functor $(F, \varphi, \varepsilon): \mathcal{C} \rightarrow \mathcal{D}$ such that F is an equivalence of categories, one can show that it is possible to choose a monoidal functor $(G, \psi, \epsilon): \mathcal{D} \rightarrow \mathcal{C}$ such that there are monoidal natural isomorphisms $F \circ G \rightarrow \text{id}_{\mathcal{D}}$ and $G \circ F \rightarrow \text{id}_{\mathcal{C}}$. In that case, we say that \mathcal{C} and \mathcal{D} are *monoidally equivalent*. For more details, see Proposition 4.4.2 in [SR72].

Example 1.16. (1) For a group G and a field \mathbb{k} , the forgetful functor $F: \mathbf{Rep}_{\mathbb{k}}(G) \rightarrow \mathbf{Vect}_{\mathbb{k}}$ which sends a G -module to the underlying vector space is monoidal. (The structure maps φ and ε are identity maps.) For a G -module V and for $g \in G$, let us write $\varphi(g)_V \in \text{End}_{\mathbb{k}}(V)$ for the action of g on V . Then $\varphi(g)$ defines a natural transformation from the functor F to itself; we write $\varphi(g) \in \text{End}(F)$. (This follows from the fact that for a homomorphism $f: V \rightarrow W$ of G -modules, the equality $\varphi(g)_W \circ F(f) = F(f) \circ \varphi(g)_V$ is equivalent to $g \cdot f(v) = f(g \cdot v)$ for $v \in V$.) The natural transformation $\varphi(g)$ is monoidal because $\varphi(g)_{V \otimes W} = \varphi(g)_V \otimes \varphi(g)_W$ and $\varphi_{\mathbb{k}}(g) = \text{id}_{\mathbb{k}}$, by definition of the tensor product of G -modules and of the trivial G -module. This example will play an important role later in the course.

Conversely, every vector space can be considered as a G -module with the trivial action of G , and this gives rise to a monoidal functor $e: \mathbf{Vect}_{\mathbb{k}} \rightarrow \mathbf{Rep}_{\mathbb{k}}(G)$.

(2) For a commutative ring R , there is a functor $F: \mathbf{Set}^{\text{rev}} \rightarrow R\text{-}\mathbf{Mod}$ that sends a set X to the free R -module $F(X) := R^X = \text{Map}(X, R)$. At the level of homomorphisms, we define $F(f)$ via $g \mapsto g \circ f$, for maps $f: X \rightarrow Y$ and $g: Y \rightarrow R$. This functor is monoidal with respect to the monoidal structure on \mathbf{Set} via the Cartesian product and on $R\text{-}\mathbf{Mod}$ via the usual tensor product of R -modules.

- (3) The total cohomology functor $H^*: \mathbf{coch}(\mathbf{Vect}_{\mathbb{k}}) \rightarrow \mathbf{Vect}_{\mathbb{k}}^{\mathbb{Z}}$ from the category of cochain complexes of \mathbb{k} -vector spaces to the category of graded \mathbb{k} -vector spaces is monoidal with respect to the usual derived tensor product on $\mathbf{coch}(\mathbf{Vect}_{\mathbb{k}})$ by the Künneth theorem: For two cochain complexes X_{\bullet} and Y_{\bullet} , we have

$$H^i(X_{\bullet} \otimes Y_{\bullet}) \cong \bigoplus_{j+k=i} H^j(X_{\bullet}) \otimes H^k(Y_{\bullet}),$$

matching the definition of the tensor product of \mathbb{Z} -graded vector spaces.

- (4) Let G and H be groups, let A be an abelian group and let $\omega: G^{\times 3} \rightarrow A$ and $\pi: H^{\times 3} \rightarrow A$ be 3-cocycles. Suppose that there is a monoidal functor $(F, \varphi, \varepsilon)$ from $\mathcal{C}_A^{G, \omega}$ to $\mathcal{C}_A^{H, \pi}$, for some $\varphi: - \otimes - \rightarrow - \otimes -$ and $\varepsilon \in \text{End}_{\mathcal{C}_A^G}(\delta_e) = A$. Then F defines a map $f: G \rightarrow H$ via $F(\delta_g) = \delta_{f(g)}$ for $g \in G$, and f is a homomorphism because

$$\delta_{f(gh)} = F(\delta_{gh}) = F(\delta_g \otimes \delta_h) \cong F(\delta_g) \otimes F(\delta_h) = \delta_{f(g)} \otimes \delta_{f(h)} = \delta_{f(g)f(h)}$$

for $g, h \in G$. Furthermore, φ defines a map $\varphi: G \times G \rightarrow A$ via

$$A \ni \varphi(g, h) := \varphi_{g, h}: F(\delta_{gh}) = F(\delta_g \otimes \delta_h) \rightarrow F(\delta_g) \otimes F(\delta_h) = \delta_{f(g)} \delta_{f(h)} = \delta_{f(gh)}$$

for $g, h \in G$, and by the definition of monoidal functors, we have

$$\varphi(g, h) \varphi(gh, k) \omega(g, h, k) = \underbrace{\pi(f(g), f(h), f(k))}_{=f^*\pi(g, h, k)} \varphi(h, k) \varphi(g, hk)$$

for all $g, h, k \in G$, that is

$$f^*\pi^{-1}\omega(g, h, k) = \varphi(h, k) \cdot \varphi(g, hk) \cdot \varphi(g, h)^{-1} \cdot \varphi(gh, k)^{-1} = d^2\varphi(g, h, k).$$

In other words, the 3-cocycle $\omega f^*\pi^{-1} = d^2\varphi$ is a 3-coboundary, so ω and $f^*\pi$ define the same element of the third cohomology group $H^3(G, A)$. (The latter is defined as the quotient of the group of 3-cocycles by the group of 3-coboundaries.) This (and the discussion in point (9) of Example 1.11) relates the equivalence classes of monoidal structures on \mathcal{C}_A^G to $H^3(G, A)$. Similarly, one can relate the equivalence classes of monoidal structures on $\mathbf{Vect}_{\mathbb{k}}^G$ to $H^3(G, \mathbb{k}^{\times})$. For more details, see Section 2.6 in [EGNO15].

2 Module categories

Unless otherwise stated, we continue to assume in this section that $(\mathcal{C}, \mathbf{1}, \otimes, \alpha, \lambda, \rho)$ is a monoidal category (which we usually abbreviate by \mathcal{C}).

Definition 2.1. A *right \mathcal{C} -module category* is a quadruple $(\mathcal{M}, \otimes_a, \beta, \vartheta)$, where

- \mathcal{M} is a category,
- $\otimes_a: \mathcal{M} \times \mathcal{C} \rightarrow \mathcal{M}$ is a bifactor, called the *action*,
- $\beta: - \otimes_a (- \otimes -) \rightarrow (- \otimes_a -) \otimes_a -$ is a natural isomorphism, called the *associativity constraint*,
- $\vartheta: - \otimes_a \mathbf{1} \rightarrow \text{id}_{\mathcal{M}}$ is a natural isomorphism, called the *unit*,

subject to the following axioms:

Pentagon axiom: For all objects M of \mathcal{M} and A, B, C of \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc}
M \otimes_a (A \otimes (B \otimes C)) & \xrightarrow{\text{id}_M \otimes \alpha_{A,B,C}} & M \otimes_a ((A \otimes B) \otimes C) \\
\downarrow \beta_{M,A,B \otimes C} & & \downarrow \beta_{M,A \otimes B,C} \\
(M \otimes_a A) \otimes_a (B \otimes C) & & (M \otimes_a (A \otimes B)) \otimes_a C \\
\searrow \beta_{M \otimes_a A,B,C} & & \swarrow \beta_{M,A,B} \otimes \text{id}_C \\
& ((M \otimes_a A) \otimes_a B) \otimes_a C &
\end{array}$$

In other words, we have

$$(\beta_{M \otimes_a A,B,C}) \circ (\beta_{M,A,B \otimes C}) = (\beta_{M,A,B} \otimes \text{id}_C) \circ (\beta_{M,A \otimes B,C}) \circ (\text{id}_M \otimes \alpha_{A,B,C}).$$

Unit axiom / triangle axiom: For all objects M of \mathcal{M} and A of \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc}
M \otimes_a (\mathbf{1} \otimes A) & \xrightarrow{\beta_{M,\mathbf{1},A}} & (M \otimes_a \mathbf{1}) \otimes_a A \\
\searrow \text{id}_M \otimes \lambda_A & & \swarrow \vartheta_M \otimes \text{id}_B \\
& M \otimes_a A &
\end{array}$$

In other words, we have

$$(\vartheta_M \otimes \text{id}_A) \circ \beta_{M,\mathbf{1},A} = \text{id}_M \otimes \lambda_A.$$

Remark 2.2. (1) We can analogously define a *left \mathcal{C} -module category* to be a tuple $(\mathcal{M}, \otimes_a, \beta, \vartheta)$ with an action bifunctor $\otimes_a: \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$, an associativity constraint

$$\beta: (- \otimes -) \otimes_a - \rightarrow - \otimes_a (- \otimes_a -)$$

and a unitor $\vartheta: \mathbf{1} \otimes_a - \rightarrow \text{id}_{\mathcal{M}}$.

(2) Being a module category over \mathcal{C} is not a property of a given category but an additional structure.

Example 2.3. (1) \mathcal{C} is a right \mathcal{C} -module category if we set $\otimes_a = \otimes$, $\beta = \alpha$ and $\vartheta = \rho$.

(2) For two categories \mathcal{C} and \mathcal{D} , the category $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$ is a right $\mathbf{End}(\mathcal{C})$ -module category, where \otimes_a is defined by composition of functors:

$$(F, G) \mapsto F \otimes_a G := F \circ G, \quad (\eta, \nu) \mapsto \eta \otimes_a \nu := \eta \nu$$

for functors $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{C} \rightarrow \mathcal{C}$ and natural transformations $\eta: F \rightarrow F'$ and $\nu: G \rightarrow G'$.

(3) For a field \mathbb{k} and a group G , the category $\mathbf{Rep}_{\mathbb{k}}(G)$ is a right $\mathbf{Vect}_{\mathbb{k}}(G)$ -module category. Indeed, viewing a \mathbb{k} -vector space as a trivial G -module gives rise to a (monoidal) functor $e: \mathbf{Vect}_{\mathbb{k}}(G) \rightarrow \mathbf{Rep}_{\mathbb{k}}(G)$, and we can define $- \otimes_a - = - \otimes e(-)$

Lemma 2.4. *Given a left module category $(\mathcal{M}, \otimes_a, \beta, \vartheta)$ over \mathcal{C} , there is a monoidal functor*

$$(F, \varphi, \varepsilon): \mathcal{C} \longrightarrow \mathbf{End}(\mathcal{M})$$

with $F = \text{id}_{\mathcal{C}} \otimes_a -$,

$$\varphi_{A,B} = \beta_{A,B,-}: F(A \otimes B) = (A \otimes B) \otimes_a - \xrightarrow{\sim} A \otimes_a (B \otimes_a -) = F(A) \circ F(B)$$

and $\varepsilon = \vartheta: \mathbf{1} \otimes_a - \rightarrow \text{id}_{\mathcal{M}}$.

Proof. This is straightforward to check using the definitions. \square

Lemma 2.5. A monoidal functor $(F, \varphi, \varepsilon): \mathcal{C} \rightarrow \mathbf{End}(\mathcal{M})$ gives rise to a left \mathcal{C} -module category structure $(\mathcal{M}, \otimes_a, \beta, \vartheta)$ via $- \otimes_a - = F(-)(-)$,

$$\beta_{A,B,M} = (\varphi_{A,B})_M: (A \otimes B) \otimes_a M = F(A \otimes B)(M) \xrightarrow{\sim} (F(A) \circ F(B))(M) = A \otimes_a (B \otimes_a M)$$

and $\vartheta = \varepsilon: \mathbf{1} \otimes_a - = F(\mathbf{1})(-) \xrightarrow{\sim} \text{id}_{\mathcal{M}}$.

Remark 2.6. Combining Lemmas 2.4 and 2.5, we see that there is a one-to-one correspondence between left \mathcal{C} -module structures on a category \mathcal{M} and monoidal functors $\mathcal{C} \rightarrow \mathbf{End}(\mathcal{M})$.

Definition 2.7. Let $(\mathcal{M}, \otimes_a, \beta, \vartheta)$ and $(\mathcal{M}', \otimes'_a, \beta', \vartheta')$ be two right \mathcal{C} -module categories. A right \mathcal{C} -module functor from \mathcal{M} to \mathcal{M}' is a pair (F, γ) , where $F: \mathcal{M} \rightarrow \mathcal{M}'$ is a functor and

$$\gamma: F(- \otimes_a -) \longrightarrow F(-) \otimes'_a -$$

is a natural isomorphism such that the following diagrams commute for all objects X, Y of \mathcal{C} and M of \mathcal{M} :

$$\begin{array}{ccc} F(M \otimes_a (X \otimes Y)) & \xrightarrow{F(\beta_{M,X,Y})} & F((M \otimes_a X) \otimes_a Y) \\ \downarrow \gamma_{M,X \otimes Y} & & \searrow \gamma_{M \otimes_a X, Y} \\ & & F(M \otimes_a X) \otimes'_a Y \\ & & \swarrow \gamma_{M,X} \otimes'_a \text{id}_Y \\ F(M) \otimes'_a (X \otimes Y) & \xrightarrow{\beta'_{F(M),X,Y}} & (F(M) \otimes'_a X) \otimes'_a Y \end{array}$$

$$\begin{array}{ccc} F(M \otimes_a \mathbf{1}) & \xrightarrow{\gamma_{M,\mathbf{1}}} & F(M) \otimes'_a \mathbf{1} \\ \searrow F(\vartheta_M) & & \swarrow \vartheta'_{F(M)} \\ & F(M) & \end{array}$$

A right \mathcal{C} -module natural transformation from a right \mathcal{C} -module functor $(F, \gamma): \mathcal{M} \rightarrow \mathcal{M}'$ to a right \mathcal{C} -module functor $(F', \gamma'): \mathcal{M} \rightarrow \mathcal{M}'$ is a natural transformation $\psi: F \rightarrow F'$ such that the following diagram commutes for all objects X of \mathcal{C} and M of \mathcal{M} :

$$\begin{array}{ccc} F(M \otimes_a X) & \xrightarrow{\gamma_{M,X}} & F(M) \otimes'_a X \\ \downarrow \psi_{M \otimes_a X} & & \downarrow \psi_M \otimes'_a \text{id}_X \\ F'(M \otimes_a X) & \xrightarrow{\gamma'_{M,X}} & F'(M) \otimes'_a X \end{array}$$

We write $\mathbf{End}_{\text{mod-}\mathcal{C}}(\mathcal{M})$ for the monoidal category of right \mathcal{C} -module endofunctors of a right \mathcal{C} -module category \mathcal{M} , with homomorphisms given by the right \mathcal{C} -module natural transformations. The tensor product is defined as the composition of functors. (There is a canonical choice of structure maps which endows the composition of two right \mathcal{C} -module functors with the structure of a right \mathcal{C} -module functor, cf. part (2) of Remark 1.15)

3 Strictness and coherence

Definition 3.1. A monoidal category \mathcal{C} is called *strict* if

$$X \otimes (Y \otimes Z) = (X \otimes Y) \otimes Z \quad \text{and} \quad \mathbf{1} \otimes X = X = X \otimes \mathbf{1}$$

for all objects X, Y, Z of \mathcal{C} (note that we require equalities and not isomorphisms) and if all associativity constraints and unitors are identity maps.

Example 3.2. The monoidal category $\mathbf{End}(\mathcal{D})$ of endofunctors of a category \mathcal{D} is strict, and so is the category $\mathbf{End}_{\text{mod-}\mathcal{C}}(\mathcal{M})$ of right \mathcal{C} -module endofunctors of a right \mathcal{C} -module category \mathcal{M} .

Theorem 3.3. *Every monoidal category is monoidally equivalent to a strict monoidal category.*

Proof (sketch). We show that \mathcal{C} is equivalent to the strict category $\mathcal{C}' := \mathbf{End}_{\text{mod-}\mathcal{C}}(\mathcal{C})$ of right \mathcal{C} -module endofunctors of \mathcal{C} . We can define a functor $F: \mathcal{C} \rightarrow \mathcal{C}'$ by

$$F(A) = (A \otimes -, \alpha_{A, -, -}) \quad \text{and} \quad F(f) = f \otimes -$$

for every object A of \mathcal{C} and every morphism $f: A \rightarrow B$ in \mathcal{C} . A functor $G: \mathcal{C}' \rightarrow \mathcal{C}$ is given by

$$G(X) = X(\mathbf{1}) \quad \text{and} \quad G(\eta) = \eta_{\mathbf{1}}$$

for a right \mathcal{C} -module endofunctor (X, γ) of \mathcal{C} and a right \mathcal{C} -module natural transformation $\eta: X \rightarrow Y$. Then one can endow F and G with structure maps to make them monoidal functors, and check that $G \circ F = \text{id}_{\mathcal{C}}$ and $F \circ G$ is naturally isomorphic to the identity functor on \mathcal{C}' . Indeed, given a right \mathcal{C} -module endofunctor (X, γ) of \mathcal{C} and an object A of \mathcal{C} , we have an isomorphism

$$F \circ G((X, \gamma))(A) = X(\mathbf{1}) \otimes A \xrightarrow{\gamma_{\mathbf{1}, A}^{-1}} X(\mathbf{1} \otimes A) \xrightarrow{X(\lambda_A)} X(A)$$

which gives rise to a right \mathcal{C} -module natural isomorphism

$$\varphi_{(X, \gamma)}: F \circ G((X, \gamma)) = (X(\mathbf{1}) \otimes -, \alpha_{X(\mathbf{1}, -, -)}) \longrightarrow (X, \gamma)$$

that is natural in (X, γ) . This yields the desired (monoidal) natural isomorphism $F \circ G \rightarrow \text{id}_{\mathbf{End}_{\text{mod-}\mathcal{C}}(\mathcal{C})}$. The details of the proof are left to the reader. \square

Example 3.4. In this example, we construct a strict monoidal category that is monoidally equivalent to $\mathbf{Vect}_{\mathbb{k}}$ for a field \mathbb{k} . Let $\mathbf{Mat}_{\mathbb{k}}$ be the category whose objects are the natural numbers

$$\text{Ob}(\mathbf{Mat}_{\mathbb{k}}) := \mathbb{N} = \{0, 1, 2, \dots\},$$

with homomorphisms from $m \in \mathbb{N}$ to $n \in \mathbb{N}$ given by the set

$$\text{Hom}_{\mathbf{Mat}_{\mathbb{k}}}(m, n) := \text{Mat}_{n \times m}(\mathbb{k})$$

of $n \times m$ -matrices over \mathbb{k} (i.e. matrices with m columns and n rows). Composition is given by matrix multiplication. We can define a monoidal structure on \mathbf{Mat} by $m \otimes n = m \cdot n$ for $m, n \in \mathbb{N}$ at the level of objects. The tensor product of two matrices $A = (a_{ij}) \in \text{Mat}_{n \times m}(\mathbb{k})$ and $B = (b_{ij}) \in \text{Mat}_{n' \times m'}(\mathbb{k})$ is the *Kronecker product*

$$A \otimes B = \begin{pmatrix} a_{11}B & \cdots & a_{1m}B \\ \vdots & \ddots & \vdots \\ a_{n1}B & \cdots & a_{nm}B \end{pmatrix} \in \text{Mat}_{(nn') \times (mm')}(\mathbb{k}).$$

and the unit object is $1 \in \mathbb{N}$. (It is straightforward to verify that the Kronecker product is associative.)

We can define a functor $F: \mathbf{Mat}_{\mathbb{k}} \rightarrow \mathbf{Vect}_{\mathbb{k}}$ by $F(m) = \mathbb{k}^{\oplus m}$ for $m \in \mathbb{N}$, at the level of objects. For $A \in \text{Mat}_{n \times m}(\mathbb{k})$, we define $F(A) \in \text{Hom}_{\mathbb{k}}(\mathbb{k}^{\oplus m}, \mathbb{k}^{\oplus n})$ to be left multiplication by A on the column vector space $\mathbb{k}^{\oplus m}$. It is straightforward to see that F is fully faithful and essentially surjective, hence an equivalence of categories, and that F is monoidal.

Theorem 3.5. *Let A_1, \dots, A_n be objects of \mathcal{C} and let P and Q be two parenthesized tensor products of A_1, \dots, A_n (in this order, but not necessarily with the same parenthesization), possibly with copies of the unit object $\mathbf{1}$ inserted in different places. Let $f: P \rightarrow Q$ and $g: P \rightarrow Q$ be two isomorphisms that are obtained by composing tensor products of identity morphisms, associativity constraints, unitors and their respective inverses. Then $f = g$.*

Proof. This follows from the strictness theorem. More specifically, let $F: \mathcal{C} \rightarrow \mathcal{C}_0$ be a monoidal equivalence from \mathcal{C} to a strict monoidal category \mathcal{C}_0 . Then we have $F(P) = F(Q)$ and $F(f) = F(g)$. Since F induces a bijection between $\text{Hom}_{\mathcal{C}}(P, Q)$ and $\text{Hom}_{\mathcal{C}_0}(F(P), F(Q))$, it follows that $f = g$. \square

Remark 3.6. An example of a setting in which we can use the coherence theorem 3.5 is Lemma 1.13: For objects A and B of \mathcal{C} , the homomorphisms

$$\rho_{A \otimes B} \circ \alpha_{A, B, \mathbf{1}}: A \otimes (B \otimes \mathbf{1}) \longrightarrow A \otimes B \quad \text{and} \quad \text{id}_A \otimes \rho_B: A \otimes (B \otimes \mathbf{1}) \longrightarrow A \otimes B$$

coincide. This does not yield an alternative proof of Lemma 1.13 because the lemma is used in the proof of the strictness theorem 3.3, which is used in turn in the proof of the coherence theorem 3.5.

Another example is that for objects A, B, C, D, E of \mathcal{C} , all of the isomorphism between

$$A \otimes (B \otimes (C \otimes (D \otimes E))) \quad \text{and} \quad (((A \otimes B) \otimes C) \otimes D) \otimes E$$

that are obtained as compositions of tensor products of identity homomorphisms and associativity constraints coincide.

4 Duals and rigidity

We continue to assume that $(\mathcal{C}, \otimes, \mathbf{1}, \alpha, \lambda, \rho)$ is a monoidal category. In order to simplify the notation, we will often ignore associativity constraints and unitors in the following.

In this section, we want to discuss the notion of dual objects in monoidal categories. We start with a motivating example.

Example 4.1. Let \mathbb{k} be a field, let V be a finite-dimensional \mathbb{k} -vector space and let $V^* = \text{Hom}_{\mathbb{k}}(V, \mathbb{k})$ be the dual space of V . Then there is a canonical evaluation map

$$\text{ev}_V: V^* \otimes V \longrightarrow \mathbb{k} \quad \text{with} \quad \xi \otimes v \longmapsto \xi(v).$$

Furthermore, for a basis $\{v_1, \dots, v_n\}$ of V with dual basis $\{v_1^*, \dots, v_n^*\}$ of V^* (defined via $v_i^*(v_j) = \delta_{ij}$), there is a coevaluation map

$$\text{coev}_V: \mathbb{k} \longrightarrow V \otimes V^*, \quad \lambda \longmapsto \lambda \cdot \sum_{i=1}^n v_i \otimes v_i^*,$$

and this map is independent of the choice of basis. Using the definitions, one easily checks that the composition

$$V \cong \mathbb{k} \otimes V \xrightarrow{\text{coev}_V \otimes \text{id}_V} V \otimes V^* \otimes V \xrightarrow{\text{id}_V \otimes \text{ev}_V} V \otimes \mathbb{k} \cong V$$

coincides with the identity map on V , that is $(\text{id}_V \otimes \text{ev}_V) \circ (\text{coev}_V \otimes \text{id}_V) = \text{id}_V$.

Now suppose that we are given a vector space W with linear maps

$$e: W \otimes V \longrightarrow \mathbb{k} \quad \text{and} \quad c: \mathbb{k} \longrightarrow V \otimes W.$$

Then we can define linear maps $f: W \rightarrow V^*$ and $g: V^* \rightarrow W$ via

$$f(w) = e(w \otimes -) \quad \text{and} \quad g(\xi) = (\xi \otimes \text{id}_W) \circ c(1),$$

for $w \in W$ and $\xi \in V^*$. Now one can further show that f and g are mutually inverse if and only if

$$(\text{id}_V \otimes e) \circ (c \otimes \text{id}_V) = \text{id}_V \quad \text{and} \quad (e \otimes \text{id}_W) \circ (\text{id}_W \otimes c) = \text{id}_W.$$

This motivates the definition of duals below.

Definition 4.2. A *left dual* of an object X of \mathcal{C} is an object X^* of \mathcal{C} together with homomorphisms

$$\text{ev}_X: X^* \otimes X \longrightarrow \mathbf{1} \quad \text{and} \quad \text{coev}_X: \mathbf{1} \longrightarrow X \otimes X^*,$$

called *evaluation* and *coevaluation*, such that the compositions

$$X \cong \mathbf{1} \otimes X \xrightarrow{\text{coev}_X \otimes \text{id}_X} X \otimes X^* \otimes X \xrightarrow{\text{id}_X \otimes \text{ev}_X} X \otimes \mathbf{1} \cong X$$

and

$$X^* \cong X^* \otimes \mathbf{1} \xrightarrow{\text{id}_{X^*} \otimes \text{coev}_X} X^* \otimes X \otimes X^* \xrightarrow{\text{ev}_X \otimes \text{id}_{X^*}} \mathbf{1} \otimes X^* \cong X^*$$

afford the identity homomorphisms on X and X^* , respectively. The equalities

$$\text{id}_X = (\text{id}_X \otimes \text{ev}_X) \circ (\text{coev}_X \otimes \text{id}_X) \quad \text{and} \quad \text{id}_{X^*} = (\text{ev}_X \otimes \text{id}_{X^*}) \circ (\text{id}_{X^*} \otimes \text{coev}_X)$$

are called the *zig-zag relations*.

A *right dual* of X is an object *X of \mathcal{C} together with homomorphisms

$$\text{ev}'_X: X \otimes X^* \longrightarrow \mathbf{1} \quad \text{and} \quad \text{coev}'_X: \mathbf{1} \longrightarrow X^* \otimes X,$$

called *evaluation* and *coevaluation*, such that the compositions

$$X \cong X \otimes \mathbf{1} \xrightarrow{\text{id}_X \otimes \text{coev}'_X} X \otimes X^* \otimes X \xrightarrow{\text{ev}'_X \otimes \text{id}_X} \mathbf{1} \otimes X \cong X$$

and

$$X^* \cong \mathbf{1} \otimes X^* \xrightarrow{\text{coev}'_X \otimes \text{id}_{X^*}} X^* \otimes X \otimes X^* \xrightarrow{\text{id}_{X^*} \otimes \text{ev}'_X} X^* \otimes \mathbf{1} \cong X^*$$

afford the identity homomorphisms on X and X^* , respectively.

Example 4.3. (1) The dual space V^* of a finite-dimensional vector space V is a left (and right) dual of V in $\mathbf{Vect}_{\mathbb{k}}$.

(2) For a G -module M in $\mathbf{Rep}_{\mathbb{k}}(G)$, the dual space M^* becomes a G -module via $(g \cdot \xi)(v) = \xi(g^{-1} \cdot v)$. The linear maps $\text{ev}_M: M^* \otimes M \rightarrow \mathbb{k}$ and $\text{coev}_M: \mathbb{k} \rightarrow M \otimes M^*$ are homomorphisms of G -modules, so M^* is a left (and right) dual of M in $\mathbf{Rep}_{\mathbb{k}}(G)$.

(3) Let G be a monoid. If for $g \in G$, the one-dimensional graded vector space \mathbb{k}_g has a left dual then g has an inverse in G , since a tensor product $(\bigoplus_h V_h) \otimes \mathbb{k}_g$ admits a non-zero homomorphism to \mathbb{k}_e only if $hg = e$ for some $h \in G$. If G is a group then every G -graded vector space $V = \bigoplus_g V_g$ has a left (and right) dual, given by the dual space V^* with grading defined by $V_g^* = (V_{g^{-1}})^*$, for $g \in G$.

Lemma 4.4. *If an object X of \mathcal{C} admits a left (or right) dual then the latter is unique up to isomorphism.*

Proof. Let X_1^* and X_2^* be two left dual objects of X and denote by e_1, e_2, c_1, c_2 the corresponding evaluation and coevaluation homomorphisms. We define two homomorphisms

$$f: X_1^* \cong X_1^* \otimes \mathbf{1} \xrightarrow{\text{id}_{X_1^*} \otimes c_2} X_1^* \otimes X \otimes X_2^* \xrightarrow{e_1 \otimes \text{id}_{X_2^*}} \mathbf{1} \otimes X_2^* \cong X_2^*$$

and

$$g: X_2^* \cong X_2^* \otimes \mathbf{1} \xrightarrow{\text{id}_{X_2^*} \otimes c_1} X_2^* \otimes X \otimes X_1^* \xrightarrow{e_2 \otimes \text{id}_{X_1^*}} \mathbf{1} \otimes X_1^* \cong X_1^*$$

and consider the following commutative diagram:

$$\begin{array}{ccccc}
X_1^* & \xrightarrow{\text{id}_{X_1^*} \otimes c_1} & X_1^* \otimes X \otimes X_1^* & & \\
\downarrow \text{id}_{X_1^*} \otimes c_2 & & \downarrow \text{id}_{X_1^*} \otimes c_2 \otimes \text{id}_{X \otimes X_1^*} & \searrow \text{id}_{X_1^* \otimes X \otimes X_1^*} & \\
X_1^* \otimes X \otimes X_2^* & \xrightarrow{\text{id}_{X_1^* \otimes X \otimes X_2^*} \otimes c_1} & X_1^* \otimes X \otimes X_2^* \otimes X \otimes X_1^* & \xrightarrow{\text{id}_{X_1^* \otimes X} \otimes e_2 \otimes \text{id}_{X_1^*}} & X_1^* \otimes X \otimes X_1^* \\
\downarrow e_1 \otimes \text{id}_{X_2^*} & & \downarrow e_1 \otimes \text{id}_{X_2^* \otimes X \otimes X_1^*} & & \downarrow e_1 \otimes \text{id}_{X_1^*} \\
X_2^* & \xrightarrow{\text{id}_{X_2^*} \otimes c_1} & X_2^* \otimes X \otimes X_1^* & \xrightarrow{e_2 \otimes \text{id}_{X_1^*}} & X_1^*
\end{array}$$

The squares commute by the bifactoriality of the tensor product and the triangle commutes because X_2^* is a left dual of X . Hence the composition along the top right boundary of the diagram coincides with the composition along the bottom left boundary. The former is the identity on X_1^* because X_1^* is a left dual of X , and the latter equals $g \circ f$. Analogously, one sees that $\text{id}_{X_2^*}^* = f \circ g$, and the claim follows. \square

Remark 4.5. More specifically, the left dual of an object of \mathcal{C} is unique up to a unique isomorphism in the following sense: In the notation of Lemma 4.4, assume that we have an homomorphism $h: X_1^* \rightarrow X_2^*$ such that $e_1 = e_2 \circ (h \otimes \text{id}_X)$ and $c_2 = (\text{id}_X \otimes h) \circ c_1$. Then one can show that h coincides with the isomorphism $f = (e_1 \otimes \text{id}_{X_2^*}) \circ (\text{id}_{X_1^*} \otimes c_2)$ from the proof of Lemma 4.4.

Lemma 4.6. *If an object X of \mathcal{C} has a left dual X^* then X is a right dual of X^* , that is ${}^*(X^*) = X$. Analogously, if X has a right dual then $({}^*X)^* = X$.*

Proof. For the first claim, set $\text{ev}'_{X^*} = \text{ev}_X$ and $\text{coev}'_{X^*} = \text{coev}_X$. The second claim is analogous. \square

Lemma 4.7. *Let X and Y be objects of \mathcal{C} with left duals. Then $Y^* \otimes X^*$ is a left dual of $X \otimes Y$. Similarly, if X and Y have right duals then ${}^*Y \otimes {}^*X$ is a right dual of $X \otimes Y$.*

Proof. We define

$$\text{ev}_{X \otimes Y}: Y^* \otimes X^* \otimes X \otimes Y \xrightarrow{\text{id}_{Y^*} \otimes \text{ev}_X \otimes \text{id}_{X^*}} Y^* \otimes \mathbf{1} \otimes Y \cong Y^* \otimes Y \xrightarrow{\text{ev}_Y} \mathbf{1}$$

and

$$\text{coev}_{X \otimes Y}: \mathbf{1} \xrightarrow{\text{coev}_X} X \otimes X^* \cong X \otimes \mathbf{1} \otimes X^* \xrightarrow{\text{id}_X \otimes \text{coev}_Y \otimes \text{id}_{X^*}} X \otimes Y \otimes Y^* \otimes X^*,$$

omitting associativity constraints and unitors. It is straightforward to check that these homomorphisms satisfy the zig-zag relations. \square

Lemma 4.8. *Let \mathcal{C} and \mathcal{D} be monoidal categories and let $(F, \varphi, \varepsilon): \mathcal{C} \rightarrow \mathcal{D}$ be a monoidal functor. If an object X of \mathcal{C} has a left dual X^* then $F(X^*)$ is a left dual of $F(X)$.*

Proof. Consider the homomorphisms

$$\text{ev}_{F(X)}: F(X^*) \otimes F(X) \xrightarrow{\varphi_{X^*, X}^{-1}} F(X \otimes X^*) \xrightarrow{F(\text{ev}_X)} F(\mathbf{1}_{\mathcal{C}}) \xrightarrow{\varepsilon} \mathbf{1}_{\mathcal{D}}$$

and

$$\text{coev}_{F(X)}: \mathbf{1}_{\mathcal{D}} \xrightarrow{\varepsilon^{-1}} F(\mathbf{1}_{\mathcal{C}}) \xrightarrow{F(\text{coev}_X)} F(X \otimes X^*) \xrightarrow{\varphi_{X, X^*}} F(X) \otimes F(X^*).$$

It is straightforward to verify that $\text{ev}_{F(X)}$ and $\text{coev}_{F(X)}$ satisfy the zig-zag relations, using the zig-zag relations for ev_X and coev_X . \square

Definition 4.9. A functor $G: \mathcal{D} \rightarrow \mathcal{C}$ is called *right adjoint* of a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ if there exists a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(-, G(-)) \cong \mathrm{Hom}_{\mathcal{D}}(F(-), -)$$

of functors from $\mathcal{C} \times \mathcal{D}$ to **Set**. In that case, we also say that F is *left adjoint* to G and write $F \dashv G$.

Remark 4.10. We have $F \dashv G$ if and only if there exist natural transformations

$$\varepsilon: FG \longrightarrow \mathrm{id}_{\mathcal{D}} \quad \text{and} \quad \eta: \mathrm{id}_{\mathcal{C}} \longrightarrow GF,$$

called the *unit* and the *counit* of the adjunction $F \dashv G$, such that the compositions

$$F \xrightarrow{\mathrm{id}_F \eta} FGF \xrightarrow{\varepsilon \mathrm{id}_F} F \quad \text{and} \quad G \xrightarrow{\eta \mathrm{id}_G} GFG \xrightarrow{\mathrm{id}_G \varepsilon} G$$

are equal to id_F and id_G , respectively. The equalities $\mathrm{id}_F = (\varepsilon \mathrm{id}_F) \circ (\mathrm{id}_F \eta)$ and $\mathrm{id}_G = (\mathrm{id}_G \varepsilon) \circ (\eta \mathrm{id}_G)$ are called the *zig-zag relations*.

Given a natural isomorphism

$$\psi: \mathrm{Hom}_{\mathcal{C}}(-, G(-)) \longrightarrow \mathrm{Hom}_{\mathcal{D}}(F(-), -),$$

we can define

$$\varepsilon_X = \psi_{X, F(X)}^{-1}(\mathrm{id}_{F(X)}): X \longrightarrow GF(X) \quad \text{and} \quad \eta_Y = \psi_{G(Y), Y}(\mathrm{id}_{G(Y)}): FG(Y) \longrightarrow Y$$

for all objects X of \mathcal{C} and Y of \mathcal{D} . Conversely, given a unit $\varepsilon: FG \rightarrow \mathrm{id}_{\mathcal{D}}$ and a counit $\eta: \mathrm{id}_{\mathcal{C}} \rightarrow GF$, we obtain a natural isomorphism

$$\psi: \mathrm{Hom}_{\mathcal{C}}(-, G(-)) \longrightarrow \mathrm{Hom}_{\mathcal{D}}(F(-), -)$$

via

$$\psi_{X, Y}(f) = \varepsilon_X \circ F(f): F(X) \xrightarrow{F(f)} FG(Y) \xrightarrow{\varepsilon_X} Y$$

for objects X of \mathcal{C} and Y of \mathcal{D} and a homomorphism $f: X \rightarrow G(Y)$ in \mathcal{D} .

Example 4.11. In the monoidal category **End**(\mathcal{D}) of endofunctors of a category \mathcal{D} , a left dual of a functor $F: \mathcal{D} \rightarrow \mathcal{D}$ is the same as a left adjoint of F , and a right dual is the same as a right adjoint.

Lemma 4.12. *If an object X of \mathcal{C} has a left dual X^* then there are adjunctions*

$$- \otimes X \dashv - \otimes X^* \quad \text{and} \quad X^* \otimes - \dashv X \otimes -.$$

In particular, for objects Y and Z of \mathcal{C} , there are isomorphisms

$$\mathrm{Hom}_{\mathcal{C}}(Y \otimes X, Z) \rightarrow \mathrm{Hom}_{\mathcal{C}}(Y, Z \otimes X^*) \quad \text{and} \quad \mathrm{Hom}_{\mathcal{C}}(X^* \otimes Y, Z) \rightarrow \mathrm{Hom}_{\mathcal{C}}(Y, X \otimes Z)$$

which are natural in Y and Z .

Proof. The unit and the counit of the adjunction $- \otimes X \dashv - \otimes X^*$ are given by

$$\varepsilon_Y: (Y \otimes X^*) \otimes X \xrightarrow{\alpha_{Y, X^*, X}} Y \otimes (X^* \otimes X) \xrightarrow{\mathrm{id}_Y \otimes \mathrm{ev}_X} Y$$

and

$$\eta_Y: Y \xrightarrow{\mathrm{coev}_X \otimes \mathrm{id}_Y} (X \otimes X^*) \otimes Y \xrightarrow{\alpha_{X, X^*, Y}^{-1}} X \otimes (X^* \otimes Y).$$

The zig-zag relations follow from the defining properties of the evaluation and coevaluation homomorphisms, and the first isomorphism between Hom-sets is immediate from Remark 4.10. The second adjunction and isomorphism of Hom-sets are obtained analogously. \square

Example 4.13. Consider the category $\mathbf{AbGrp} = \mathbf{Mod}_{\mathbb{Z}}$ of abelian groups (or \mathbb{Z} -modules), with the usual tensor product $\otimes = \otimes_{\mathbb{Z}}$ over \mathbb{Z} and the unit object \mathbb{Z} . Observe that for any two abelian groups A and B , the set $\mathrm{Hom}_{\mathbb{Z}}(A, B)$ can be considered as an abelian group via pointwise addition, and that $\mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}, A) \cong A$ (by evaluation at $1 \in \mathbb{Z}$). If the group $A = \mathbb{Z}/3\mathbb{Z}$ has a left dual A^* then

$$A^* \cong \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}, A^*) \cong \mathrm{Hom}(A, \mathbb{Z}) = 0$$

by Lemma 4.12, contradicting the zig zag relations. Hence $\mathbb{Z}/3\mathbb{Z}$ does not admit a left dual in \mathbf{AbGrp} .

Remark 4.14. Before we continue discussing duals, some reminders about category theory are in order. Given two categories \mathcal{C} and \mathcal{D} , a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is called *faithful* (or *full*) if the map

$$\mathrm{Hom}_{\mathcal{C}}(X, Y) \xrightarrow{f \mapsto F(f)} \mathrm{Hom}_{\mathcal{D}}(F(X), F(Y))$$

is injective (respectively surjective). The functor F is called *fully faithful* if it is full and faithful, and it is called *essentially surjective* if for every object Y of \mathcal{D} , there exists an object X of \mathcal{C} such that $F(X) \cong Y$. One can show that a functor is fully faithful and essentially surjective if and only if it is an equivalence.

Now for every object X of \mathcal{C} , we have a functor

$$\mathrm{Hom}_{\mathcal{C}}(X, -): \mathcal{C} \longrightarrow \mathbf{Set}$$

and for a homomorphism $f: X \rightarrow Y$ in \mathcal{C} , there is a natural transformation

$$\mathrm{Hom}(f, -): \mathrm{Hom}_{\mathcal{C}}(Y, -) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, -)$$

with components

$$\mathrm{Hom}(f, Z): \mathrm{Hom}_{\mathcal{C}}(Y, Z) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, Z), \quad g \mapsto g \circ f.$$

These data give rise to a functor

$$\mathcal{C}^{\mathrm{rev}} \longrightarrow \mathbf{Fun}(\mathcal{C}, \mathbf{Set})$$

which is fully faithful by Yoneda's lemma. In particular, given two objects X and Y of \mathcal{C} such that the functors $\mathrm{Hom}_{\mathcal{C}}(X, -)$ and $\mathrm{Hom}_{\mathcal{C}}(Y, -)$ are naturally isomorphic, there exists an isomorphism $X \cong Y$ in \mathcal{C} .

Remark 4.15. Using Lemma 4.12, we can give an alternative proof of Lemma 4.4: For an element X of \mathcal{C} with two left duals X_1^* and X_2^* , there is a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(X_1^*, -) \cong \mathrm{Hom}_{\mathcal{C}}(\mathbf{1}, X \otimes -) \cong \mathrm{Hom}_{\mathcal{C}}(X_2^*, -),$$

and Yoneda's lemma implies that $X_1^* \cong X_2^*$.

Potential topics for talks

The topics that appear in gray have already been chosen.

- (1) pivotal categories, traces and dimension (?);
- (2) Frobenius-Perron dimension;
- (3) monoidal categories by generators and relations (via diagrams);
- (4) the Temperley-Lieb category (and knot invariants);
- (5) tensor triangular geometry;

- (6) the geometric Satake equivalence;
- (7) negligible morphisms and semisimplification;
- (8) the Drinfel'd double and the Drinfel'd center;
- (9) \mathfrak{sl}_n -webs;
- (10) interpolation categories;
- (11) ...

References

- [DM82] Pierre Deligne and J. S. Milne. Tannakian categories. Hodge cycles, motives, and Shimura varieties, Lect. Notes Math. 900, 101-228 (1982)., 1982.
- [EGNO09] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. Tensor categories. Lecture notes for a course taught by P. Etingof at MIT, available at http://mtm.ufsc.br/~ebatista/2016-2/tensor_categories.pdf, 2009.
- [EGNO15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*, volume 205 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [ML98] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [SR72] Neantro Saavedra Rivano. Catégories tannakiennes. *Bull. Soc. Math. Fr.*, 100:417–430, 1972.