

Random variables

A random variable is any uncertain outcome. Examples:

- ▶ Whether a coin flip comes up heads or tails.
- ▶ The winning team in the NCAA basketball tournament.
- ▶ The value of Apple stock next year.
- ▶ The name of the next new person you meet.

A probability distribution describes a random variable as a list of two things.

- ▶ The sample space: the set of possible outcomes (“events”) for the random variable
- ▶ The probability for each event in the sample space.

Examples

Soon, we'll study a few examples of common probability distributions.

- ▶ Binomial
- ▶ Poisson
- ▶ Normal

For now, let's focus on a simple example: the outcome of rolling two dice.

Examples

Suppose you roll two dice. Let X be the sum of the dice.

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

Examples

Here the the probability distribution for X .

x_k	$P(X = x_k)$
2	1/36
3	2/36
4	3/36
5	4/36
6	5/36
7	6/36
8	5/36
9	4/36
10	3/36
11	2/36
12	1/36

Expected value

The expected value, or mean, is one measure for the center of a probability distribution. Suppose the random variable X has sample space x_1, \dots, x_M . Then

$$E(X) = \sum_{k=1}^M x_k \cdot P(X = x_k)$$

It is the weighted average of the possible outcomes, where the weight on each event is that event's probability.

Dice example:

$$E(X) = 2 \cdot \frac{1}{36} + 3 \cdot \frac{2}{36} + \dots + 11 \cdot \frac{2}{36} + 12 \cdot \frac{1}{36} = 7.$$

Variance

The variance measures the spread, or dispersion, of a probability distribution. Again, suppose the random variable X has sample space x_1, \dots, x_M . Then

$$\text{var}(X) = E([X - E(X)]^2) = \sum_{k=1}^M (x_k - E(X))^2 \cdot P(X = x_k)$$

It is the weighted average of the squared deviations from the mean, where the weight on each event is that event's probability.

Dice example:

$$\text{var}(X) = (2 - 7)^2 \cdot \frac{1}{36} + (3 - 7)^2 \cdot \frac{2}{36} + \dots + (12 - 7)^2 \cdot \frac{1}{36} \approx 5.83.$$