

Perspective: The Convergence of Coronavirus Disease 2019 (COVID-19) and Food Insecurity in the United States

Jason M Nagata,¹ Hilary K Seligman,^{2,3} and Sheri D Weiser²

¹Department of Pediatrics, University of California, San Francisco, San Francisco, CA, USA; ²Department of Medicine, University of California, San Francisco, San Francisco, CA, USA; and ³Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, CA, USA

ABSTRACT

During the coronavirus disease 2019 (COVID-19) pandemic, food insecurity has doubled overall and tripled among households with children in the United States. Food insecurity and COVID-19 may exacerbate one another through bidirectional links, leading to a syndemic, or sequential disease clusters, which exacerbate one another. Experiencing food insecurity may be associated with macronutrient and micronutrient deficiencies, which can weaken host defenses, thus increasing susceptibility to COVID-19. Food insecurity is associated with chronic medical conditions, which may afford a higher risk of severe COVID-19 illness. People experiencing food insecurity may have increased exposure to COVID-19 while procuring food. People with COVID-19 may be unable to work, generate income, and procure food while quarantined, which may exacerbate food insecurity. Clinicians should screen for food insecurity during the COVID-19 pandemic and provide referrals to food-assistance programs when appropriate. Policymakers should expand benefits for the Supplemental Nutrition Assistance Program (SNAP) and the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) to address increases in the depth and breadth of food insecurity during the COVID-19 pandemic. *Adv Nutr* 2021;12:287–290.

Keywords: COVID-19, food insecurity, health policy, nutrition, screening

Introduction

During the coronavirus disease 2019 (COVID-19) pandemic, food insecurity is estimated to have doubled overall and tripled among households with children in the United States (1). Although there is increasing recognition that social-distancing measures may contribute to unemployment and food insecurity, less attention has been focused on how food insecurity may relate to COVID-19. Food insecurity and COVID-19 may exacerbate one another

through bidirectional links leading to a syndemic (2), where conditions interact synergistically to exacerbate negative health impacts of each.

COVID-19 May Exacerbate Food Insecurity

People with COVID-19 may be unable to work for income due to symptoms or, even if asymptomatic, due to self-isolation measures. People with COVID-19 may also accrue significant health care costs associated with testing and treatment, which may displace budgets for food. Stigma or fear of contagion associated with having the illness may lead to further isolation and an inability to rely on regular support networks, which can exacerbate food insecurity.

Widespread unemployment and reduced hours due to the public health response to COVID-19 are associated with increased food insecurity rates in the general population; the lowest-wage positions (e.g., retail, food services) are most vulnerable to job losses (3). Food banks across the country report skyrocketing demand, with upwards of 30% being first-time users of the charitable food system. California's online Supplemental Nutrition Assistance Program (SNAP) application program, GetCalFresh, provides real-time insight into the increasing demand for SNAP benefits, with a daily

JMN is supported by the American Heart Association (CDA34760281). HKS is supported by the Centers for Disease Control and Prevention (6U48DP006374). SDW is supported by the National Institute of Allergy and Infectious Diseases (K24AI134326).

Author disclosures: HKS receives funding from Feeding America, a 501c3 nonprofit dedicated to ending hunger in the United States. All other authors report no conflicts of interest. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Perspective articles allow authors to take a position on a topic of current major importance or controversy in the field of nutrition. As such, these articles could include statements based on author opinions or point of view. Opinions expressed in Perspective articles are those of the author and are not attributable to the funder(s) or the sponsor(s) or the publisher, Editor, or Editorial Board of *Advances in Nutrition*. Individuals with different positions on the topic of a Perspective are invited to submit their comments in the form of a Perspectives article or in a Letter to the Editor.

Address correspondence to JMN (email: jasonmnagata@gmail.com).

Abbreviations used: COVID-19, coronavirus disease 2019; PEB-T, Pandemic Electronic Benefits Transfer; SNAP, Supplemental Nutrition Assistance Program; WIC, Special Supplemental Nutrition Program for Women, Infants, and Children.

application rate in April nearly 4 times higher than usual; almost half of the applicants (48%) were first-time applicants, and 57% reported job loss in the last 30 d (compared with 16% in January).

Food procurement in food-insecure households often requires more time, cognitive bandwidth, and logistical coordination than in food-secure households. Quarantine, self-isolation, and shelter in place are important to protect public health, but at the same time are exacerbating these food-procurement challenges.

Food Insecurity May Lead to Susceptibility to COVID-19

Food insecurity may lead to susceptibility to COVID-19 through biological and behavioral mechanisms. First, food insecurity may lead to micronutrient and macronutrient deficiencies, which can weaken host defense mechanisms and contribute to immunologic decline. Such immunologic decline can increase susceptibility to COVID-19 and morbidity among those with COVID-19 (4). Food insecurity is associated with chronic medical conditions, including chronic kidney disease, obstructive pulmonary disease, obesity, coronary artery disease (5, 6), and diabetes (7, 8). In the case of diabetes, longitudinal studies have shown that food insecurity is prospectively associated with subsequent elevations in diabetes risk (7, 8). These same medical conditions are associated with an increased risk of severe illness from COVID-19 (9). Food insecurity also contributes to stress, anxiety, and depression (10–12), which can also impair immune function and increase susceptibility to infection (6). Finally, food insecurity is associated with smoking (11), which is also a risk factor for severe illness, intensive care unit admission, need for mechanical ventilation, and death from COVID-19 (13).

Behavioral pathways by which food insecurity can increase susceptibility to COVID-19 include needing to access foods through alternate distribution channels, such as food pantries where food may need to be obtained more frequently or potentially through more crowded venues, or needing to access food more frequently due to financial limitations. Furthermore, poverty and food insecurity are closely linked. Poverty is a risk factor for food insecurity and people experiencing poverty may have a decreased ability to self-isolate due to shared living spaces, crowded housing, or caring for dependent children or elders. People with low incomes may also work in essential jobs, which increases COVID-19 exposure (such as public transportation, food retail/delivery, and nonclinical health care workers such as hospital janitors) (14).

Food Insecurity in Vulnerable Populations

Vulnerable and marginalized populations are disproportionately affected by both food insecurity and COVID-19 (4). In the United States, people living in poverty and racial and ethnic minorities have been disproportionately affected by both COVID-19 and food insecurity (15). COVID-19 outbreaks have emerged in homeless shelters and prison populations. This is also true globally. Low-income countries

may be the hardest hit by food insecurity during the COVID-19 pandemic and may also lack the health infrastructure and workforce to care for a surge in patients with COVID-19.

School closures during the COVID-19 pandemic have important food insecurity implications for low-income children and their families. Over 30 million children rely on the National School Lunch Program and School Breakfast Program, and meals and snacks from schools fulfill up to two-thirds of children's daily nutritional needs (16). Without access to school meals (or a replacement meal provided by the school) during the pandemic, low-income families have the financial burden of providing additional meals for their children (16). Parents may also not be able to work and generate income while providing childcare for their children during school closures, which may exacerbate food insecurity. Early in the pandemic, policymakers recognized these concerns and responded with alternative distribution mechanisms for school meals, the Pandemic Electronic Benefits Transfer (P-EBT) program, and unemployment insurance. However, as the pandemic and school closures continue, renewals, extensions, and longer-term policies will need to be considered.

Health Care Provider Actions

Health care professionals can help address this syndemic of COVID-19 and food insecurity. Although screening for food insecurity is recommended at least annually by many professional medical organizations (American Academy of Pediatrics, American Academy of Family Physicians, American College of Physicians, and American College of Obstetrics and Gynecology) (17), clinicians should screen for food insecurity at every in-person and telehealth visit during the COVID-19 pandemic. A brief, 2-item screener has 97% sensitivity and 83% specificity for food insecurity (Table 1) (18). Dietitians or social workers can provide referrals to food resources such as food pantries; SNAP; the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC); or the P-EBT program. Eligibility criteria and geographic availability vary substantially.

For people with COVID-19 or at a high risk of severe disease, clinicians and social workers can help patients navigate home food-delivery resources, leveraging innovative programming that is currently undergoing massive expansion. Such programming includes, for example, use of SNAP benefits online, home delivery of groceries from the food bank, and home-delivered meals. Again, eligibility criteria and geographic availability vary.

Policy Recommendations

The Families First Coronavirus Response Act (H.R. 6201), signed on 18 March 2020, lifted work and work-training requirements, and temporarily suspended time limits for SNAP enrollment for able-bodied adults without dependents who are not working, in recognition of rising unemployment rates (19). The Coronavirus Aid, Relief, and Economic Security (CARES) Act (H.R. 748), signed on 27 March 2020 (20), appropriated nearly \$16 billion for SNAP benefits

TABLE 1 Actions for health care providers to address food insecurity during the COVID-19 pandemic¹

- Screen for food insecurity at each clinical visit during the COVID-19 pandemic. A response of “sometimes” or “often” true to either question should trigger a clinical response.
“Within the past 12 months, we worried whether our food would run out before we got money to buy more.”
Reponses: never true, sometimes true, often true (18)
“Within the past 12 months, the food we bought just didn’t last, and we didn’t have the money to get more.”
Reponses: never true, sometimes true, often true (18)
- Provide referrals and support for programs such as community food distribution programs (21); the Supplemental Nutrition Assistance Program (SNAP) (22); the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) (23); and the National School Lunch Program (24) and School Breakfast Program (25).
- Prescribe medically tailored meals to those with food insecurity and medical conditions.

¹ COVID-19, coronavirus disease 2019.

for newly eligible households (SNAP is an entitlement program, meaning that anyone qualifying under program rules can receive benefits). However, these bills did not expand SNAP benefits to address increases in the depth and breadth of food insecurity. Current proposals for expansion include raising the maximum SNAP benefit by 15%. This increase would support the food security of households enrolled in SNAP while quickly pushing money into local economies, helping to mitigate the effect of the economic crisis.

For WIC, the Families First Coronavirus Response Act authorized an additional \$500 million and allowed for additional flexibility—for example, it waived the need to be physically present for enrollment (to reduce potential exposure to COVID-19 at enrollment sites) and allowed for substitution of WIC-approved products (to accommodate COVID-19–related gaps in the food supply). Congress should support the WIC Benefit Flexibility During COVID-19 Act (H.R. 6726), which would increase the cash value of WIC’s fruit and vegetable vouchers to \$35 to enable more healthy dietary intake (26). Similar to new flexibilities in the SNAP program, WIC participants should also be able to purchase groceries online and use alternative food-delivery models to mitigate COVID-19 exposure.

Conclusions

Food insecurity and COVID-19 are intertwined processes that exacerbate one another and disproportionately affect vulnerable populations. Particularly in the current crisis, clinicians should be regularly screening patients for food insecurity and supporting them in accessing food resources. Policymakers should expand legislation to address food insecurity and poverty as part of their efforts to halt the COVID-19 pandemic.

Acknowledgments

We appreciate the support of Tracy Fox, Food, Nutrition, & Policy Consultants LLC, for her review of the policy summary as well as Vanessa Machen, Jacqueline Genovese, and Samuel Benabou for their review of the article. The authors’ responsibilities were as follows—JMN: drafted the manuscript; SDW and HKS: reviewed and critically revised

the manuscript; and all authors: read and approved the final manuscript.

References

1. Schanzebach D, Pitts A. How much has food insecurity risen? Evidence from the Census Household Pulse Survey [Internet]. 2020. [cited 2020 Aug 22]. Available from: <https://www.ipr.northwestern.edu/documents/reports/ipr-rapid-research-reports-pulse-hh-data-10-june-2020.pdf>.
2. Carlson CJ, Mendenhall E. Preparing for emerging infections means expecting new syndemics. *Lancet North Am Ed* 2019;394:297.
3. Lund S, Ellingrud K, Hancock B, Manyika J, Dua A. The near-term impact of coronavirus on workers [Internet]. 2020. [cited 2020 Aug 22]. Available from: <https://www.mckinsey.com/industries/public-sector/our-insights/lives-and-livelihoods-assessing-the-near-term-impact-of-covid-19-on-us-workers>.
4. Weiser SD, Young SL, Cohen CR, Kushel MB, Tsai AC, Tien PC, Hatcher AM, Frongillo EA, Bangsberg DR. Conceptual framework for understanding the bidirectional links between food insecurity and HIV/AIDS. *Am J Clin Nutr* [Internet]. 2011;94:1729S–39S. Available from: <http://www.ajcn.org/content/94/6/1729S.abstract>.
5. Nagata JM, Palar K, Gooding HC, Garber AK, Bibbins-Domingo K, Weiser SD. Food insecurity and chronic disease in US young adults: findings from the National Longitudinal Study of Adolescent to Adult Health. *J Gen Intern Med* [Internet]. 2019;34:2756–62. [cited 2020 Aug 21]. Available from: <https://doi.org/10.1007/s11606-019-05317-8>.
6. Seligman HK, Berkowitz SA. Aligning programs and policies to support food security and public health goals in the United States. *Annu Rev Public Health* 2019;40:319–37.
7. Berkowitz SA, Gao X, Tucker KL. Food-insecure dietary patterns are associated with poor longitudinal glycemic control in diabetes: results from the Boston Puerto Rican Health Study. *Diabetes Care* [Internet]. 2014;37:2587–92. [cited 2020 Aug 21]. Available from: <https://pubmed.ncbi.nlm.nih.gov/24969578/>.
8. Berkowitz SA, Karter AJ, Corbie-Smith G, Seligman HK, Ackroyd SA, Barnard LS, Atlas SJ, Wexler DJ. Food insecurity, food “deserts,” and glycemic control in patients with diabetes: a longitudinal analysis. *Diabetes Care* [Internet]. 2018;41:1188–95. [cited 2020 Aug 21]. Available from: <https://pubmed.ncbi.nlm.nih.gov/29555650/>.
9. Chow N, Fleming-Dutra K, Gierke R, Hall A, Hughes M, Pilishvili T, Ritchey M, Roguski K, Skoff T, Ussery E. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019—United States, February 12–March 28, 2020. *MMWR Morb Mortal Wkly Rep* [Internet]. 2020;69:382–6. [cited 2020 Aug 21]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32240123/>.
10. Nagata JM, Palar K, Gooding HC, Garber AK, Whittle HJ, Bibbins-Domingo K, Weiser SD. Food insecurity is associated with poorer mental health and sleep outcomes in young adults. *J Adolesc Health* [Internet]. 2019;65(6):805–11. [cited 2020 Aug 21]. Available from: <https://doi.org/10.1016/j.jadohealth.2019.08.010>.

11. Nagata JM, Palar K, Gooding HC, Garber AK, Whittle HJ, Tabler J, Bibbins-Domingo K, Weiser SD. Food insecurity, sexual risk, and substance use among young adults. *J Adolesc Health* 2020. [cited 2020 Aug 21]. Available from: <https://doi.org/10.1016/j.jadohealth.2020.05.038>.
12. Nagata JM, Gomberg S, Hagan MJ, Heyman MB, Wojcicki JM. Food insecurity is associated with maternal depression and child pervasive developmental symptoms in low-income Latino households. *J Hunger Environ Nutr* 2019;14(4):526–39, doi:10.1080/19320248.2018.1434101.
13. Vardavas CI, Nikitara K. COVID-19 and smoking: a systematic review of the evidence. *Tobacco Induced Diseases* [Internet] 2020. [cited 2020 Aug 21]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32206052/>.
14. Parks CA, Nugent NB, Fleischacker SE, Yaroch AL. Food system workers are the unexpected but under protected COVID heroes. *J Nutr* 2020;150(8):2006–8.
15. Webb Hooper M, Nápoles AM, Pérez-Stable EJ. COVID-19 and racial/ethnic disparities. *JAMA* [Internet] 2020. [cited 2020 May 12]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32391864>.
16. Dunn CG, Kenney E, Fleischacker SE, Bleich SN. Feeding low-income children during the CoviD-19 pandemic. *N Engl J Med* [Internet] 2020;382:e40. [cited 2020 Jul 7]. Available from: <http://www.nejm.org/doi/10.1056/NEJMp2005638>.
17. Patil SP, Craven K, Kolasa K. Food insecurity: how you can help your patients. *Am Fam Physician* [Internet] 2018;98:143–5. [cited 2020 May 1]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30215895>.
18. Hager ER, Quigg AM, Black MM, Coleman SM, Heeren T, Rose-Jacobs R, Cook JT, de Cuba SA, Casey PH, Chilton M, et al. Development and validity of a 2-item screen to identify families at risk for food insecurity. *Pediatrics* 2010;126:26.
19. Lowey NM. H.R. 6201—116th Congress (2019–2020): Families First Coronavirus Response Act [Internet]. 2020. [cited 2020 Aug 22]. Available from: <https://www.congress.gov/bill/116th-congress/house-bill/6201>.
20. Courtney J. H.R. 748—116th Congress (2019–2020): CARES Act [Internet]. 2020. [cited 2020 Aug 22]. Available from: <https://www.congress.gov/bill/116th-congress/house-bill/748>.
21. US Department of Agriculture. USDA foods [Internet]. [cited 2020 Aug 22]. Available from: <https://www.fns.usda.gov/usda-foods>.
22. US Department of Agriculture. Supplemental Nutrition Assistance Program (SNAP) [Internet]. [cited 2020 Aug 22]. Available from: <https://www.fns.usda.gov/snap/supplemental-nutrition-assistance-program>.
23. US Department of Agriculture. Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) [Internet]. [cited 2020 Aug 22]. Available from: <https://www.fns.usda.gov/wic>.
24. US Department of Agriculture. National School Lunch Program [Internet]. [cited 2020 Aug 22]. Available from: <https://www.fns.usda.gov/nslp>.
25. US Department of Agriculture. School Breakfast Program [Internet]. [cited 2020 Aug 22]. Available from: <https://www.fns.usda.gov/sbp/school-breakfast-program>.
26. Schrier K. H.R. 6726—116th Congress (2019–2020): WIC Benefit Flexibility During COVID-19 Act [Internet]. 2020. [cited 2020 Aug 22]. Available from: <https://www.congress.gov/bill/116th-congress/house-bill/6726>.