

MIT Introduction to Statistics 18.05 Reading 3

John Hancock

February 8, 2017

Contents

1	References and License	1
2	What is $P(S_2 S_1^c)$	1

1 References and License

We are answering questions in the material from MIT OpenCourseWare course 18.05, Introduction to Probability and Statistics.

Please see the references section for detailed citation information.

The material for the course is licensed under the terms at <http://ocw.mit.edu/terms>.

2 What is $P(S_2 | S_1^c)$

S_1 = 'first card is a spade' S_2 = 'second card is a spade'

Therefore S_2^c = 'second card is not a spade'

We will calculate

$$P(S_2 | S_1^c) \tag{1}$$

That is, we will calculate the probability that the first card is a spade given that the second card is not a spade.

We will apply the same method Orloff and Bloom use in [1], section 3, "Multiplication Rule."

If the first card is a spade then of the 51 cards remaining, $3 \times 13 = 39$ cards are not spades.

Therefore

$$P(S_2 | S_1^c) = \frac{39}{51} \approx 0.765 \tag{2}$$

We use the multiplication rule from [1] to compute the same probability.

We apply the multiplication rule letting $A = S_1$ and $B = S_2^c$

$$P\left(S_1 \mid S_2^c\right) = \frac{P\left(S_1 \cap S_2^c\right)}{P\left(S_2^c\right)} \quad (3)$$

First we calculate $P\left(S_1 \cap S_2^c\right)$.

There are

References

- [1] Jeremy Orloff and Jonathan Bloom. *Conditional Probability, Independence and Bayes Theorem Class 3, 18.05, Spring 2014*. Available at https://ocw.mit.edu/courses/mathematics/18-05-introduction-to-probability-and-statistics-spring-2014/readings/MIT18_05S14_Reading3.pdf (Spring 2014).