# Disentangling Sources of SN Ia Variation

Jared Hand<sup>1</sup>

<sup>1</sup> University of Pittsburgh, jsh89@pitt.edu

## 1. INTRODUCTION

Resolving outstanding issues standardizing Type Ia supernova (SN Ia) for use as cosmic distance measures will be the focus of my fifth year as a physics graduate student at the University of Pittsburgh. Host galaxy dust attenuation is coupled to both the observed time-independent (color) variation in SN Ia populations and an established bias between SN Ia and host galaxy properties, neither of which are appropriately accounted for during lightcurve standardization. Resolution requires directly disentangling dust-independent SN Ia color variation from host attenuation and quantifying attenuation's contribution to the observed host bias. These two approaches implemented using powerful statistical tools and techniques that form my dissertation's foundation.

## 2. RESEARCH STATEMENT

The extreme brightness of SNe Ia are standardized using correlations between peak brightness and both SED 'blueness' and explosion duration to reduce the already naturally low scatter in peak brightness, allowing their use as cosmic 'rulers' to constrain cosmological models and explore the properties of dark energy (Perlmutter et al. 1999). A lack of theoretical understanding SN Ia necessitates these transient objects be parameterized by generative empirical models trained from representative SN samples. The current linear Tripp standardization technique and commonly used SN Ia models cannot differentiate between time-independent wavelength (color) variation intrinsic to the SN Ia population and color variation from host galaxy dust attenuation (Mandel et al. 2017), and overcorrects for a known bias where brighter SN Ia prefer younger, less massive hosts with higher star formation rates (SFR) (Sullivan et al. 2010; Rigault et al. 2020). SFR and stellar age are highly correlated with both each other and dust attenuation This host bias propagating through Tripp standardization increases cosmological measurement uncertainty and is driven both by mishandling of dust effects on SN Ia color variation and ignoring the established host bias during standardization (Brout & Scolnic 2021; Rose et al. 2021).

#### 2.1. Determining the Connection between Dust and the Host Bias:

My first project focuses on systematics arising from how we measure host properties, with my first and recently resubmitted paper alleviating concerns of bias arising from chosen observation or fitting technique. This first project now is focused on quantifying dependence of host SFR and stellar age on host dust attenuation measurement using different observational SFR tracers such as H $\alpha$  emission, which provides an instantaneous (< 10 Myr) SFR measurement, and UV flux, which provides an extended (10 – 100 Myr) SFR measurement. Our planned integration of sophisticated galaxy attenuation curve modeling from Salim

et al. (2018) and Narayanan et al. (2018) will be a dramatic improvement from past SN Ia analyses. Numerous covariates including latent model parameters intrinsic to SN Ia standardization necessitates efficient posterior analysis which will be done using a hierarchical Bayesian framework implemented with a Hamiltonian Monte Carlo sampler built into the software package Stan Carpenter et al. (2017).

# 2.2. Separating Sources of SN Ia Color Variation

My second project's genesis is from a DOE SCGSR award I received in 2019 to work with Dr. Alex Kim of the Lawrence Berkeley National Laboratory using spectroscopic SN Ia time series to build a hierarchical Bayesian model with Gaussian processes to agnostically disentangle the intrinsic color variation of SNe Ia from host dust effects. This project too makes heavy use of Stan. Parameterizing color variation using a set of multiplicative color variation curves that makes no strong prior assumption on color curve shape, our model exploits mathematical symmetries to simultaneously recover dust attenuation effects and intrinsic color variation. The our model's wavelength resolution is adjustable, and time-varying color variation is being implemented to create a fully descriptive SN Ia empirical model. We are in the process of preparing our first manuscript for this project.

Mentoring Undergraduate Researchers: Joining the host bias analysis project are two undergraduates of the Michael Wood-Vasey research group who I will mentor during my remaining time as a PhD candidate. This will include preparing to present their work to the Dark Energy Science collaboration and guidance in manuscript preparation and writing. The Z. Daniels Fellowship would increase my available time to mentor these students.

**Dissertation Progress Notes:** My second committee meeting and DOE award were both delayed by health issues during 2020 and the COVID-19 pandemic. The second meeting is now planned for June 2021 and the DOE award, restarted remotely in January 2021, will conclude July 2021.

## REFERENCES

- Brout, D., & Scolnic, D. 2021, ApJ, 909, 26, doi: 10.3847/1538-4357/abd69b
- Carpenter, B., Gelman, A., Hoffman, M. D., et al. 2017, Journal of Statistical Software, Articles, 76, 1, doi: 10.18637/jss.v076.i01
- Mandel, K. S., Scolnic, D. M., Shariff, H., Foley, R. J., & Kirshner, R. P. 2017, ApJ, 842, 93, doi: 10.3847/1538-4357/aa6038
- Narayanan, D., Conroy, C., Davé, R., Johnson, B. D., & Popping, G. 2018, ApJ, 869, 70, doi: 10.3847/1538-4357/aaed25
- Perlmutter, S., Aldering, G., Goldhaber, G., et al. 1999, ApJ, 517, 565, doi: 10.1086/307221

- Rigault, M., et al. 2020, Astron. Astrophys., 644, A176,
  - doi: 10.1051/0004-6361/201730404
- Rose, B. M., Rubin, D., Strolger, L., & Garnavich, P. M. 2021, ApJ, 909, 28, doi: 10.3847/1538-4357/abd550
- Salim, S., Boquien, M., & Lee, J. C. 2018, ApJ, 859, 11,
  - doi: 10.3847/1538-4357/aabf3c
- Sullivan, M., Conley, A., Howell, D. A., et al. 2010, MNRAS, 406, 782,doi: 10.1111/j.1365-2966.2010.16731.x