

Chap 8 Notes: Confidence Interval Estimation

J. Harner A. Billings

Department of Statistics
West Virginia University

Stat 211 Fall 2007

Outline

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design

Case : $\mu_D = \mu_X - \mu_Y$

Statistical Estimation and Confidence Interval:

- Measurements should be accurate, i.e., close to the true or actual value.
- Measurements have two sources of error:
 1. random measurement error
 2. bias (systematic measurement error)
- So

measured value = true value + bias + random error

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(From Sec. 5.1)

Law of Large Numbers:

If the sample size mean \bar{X} should be close to the population mean μ .

- This implies that when n is large enough, the random errors in the data tend to "cancel out", so that

$$\begin{aligned}\bar{X} &\simeq \text{true value} + \text{bias} \\ &\simeq \mu + \text{bias}\end{aligned}$$

- Good methods and measuring instruments are necessary to try to eliminate bias.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Example:

Estimate mean height of population of 8th grade male gym students, using sample (random) of 100.

Measured height = True height + bias + random error
here,

bias : wearing tennis shoes

random error : stance (straight, slouched, etc)

and

$$\bar{X} \simeq \mu + \text{bias}$$

\bar{X} : sample mean height

μ : population mean height

bias : tennis shoes height

- Eliminate bias by removing shoes.

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Estimating Population Parameters:

- When the sample size n is large, the sample mean \bar{X} and sample standard deviation S should be reasonably good estimators of the population mean μ and population standard deviation σ .
- ▶ \bar{X}, S are (sample) statistics.
- ▶ μ, σ are (population) parameters.

Capital \bar{X}, S — random variables

Little \bar{x}, s — computed or observed numerical values

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Random sample and Estimates

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- A repeated trials random experiment, or just random experiment, consists of n independent random variables $X_1, X_2, X_3, \dots, X_n$ each with the same probability distribution (identically distributed); n is the number of replications. The collection $X_1, X_2, X_3, \dots, X_n$ of random variables is called a random sample.

Example:

In this example;

Population = all conceivable tosses;

Random sample is 100 tosses;

each toss is independent of all others.

each X has same probability distribution.

X	1	...	6
p(x)	$\frac{1}{6}$...	$\frac{1}{6}$

$n = 100$ tosses.

Sampling with replacement from a conceptual population.

Chap 8 Notes:

Confidence Interval Estimation

Sect 8.1: An Introduction To The Statistical Estimation Problem

Sect 8.2: The Central Limit Theorem

Sect 6.3: Large-Sample confidence Interval for the Population Mean μ

Sect 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sect 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sect 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sect 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sect 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

Random Sample from Population

(Survey sampling)

- Random sampling from a population usually involves sampling without replacement - we usually don't want to have the same subject in the sample twice. Thus the observations are not independent. (Recall our "marble" example from Chap. 4)
- But when the population size N is large, we can "think of" the observation $X_1, X_2, X_3, \dots, X_n$ as being independent.

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Example:

1. Box with 500 Red, 500 yellow.
Sample without replacement.
- Does first selection "affect" second selection?

$$P(R_2|R_1) = \frac{499}{999} = 0.4995$$

$$P(R_2|B_1) = \frac{500}{999} = 0.5005$$

- **Yes** - but not much!
Even less "affect" if N is bigger.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Large Population Random Sample Rule:

- When obtaining a random sample from a real population of size N , which is large compared to the sample of size n , the random sample $X_1, X_2, X_3, \dots, X_n$ can be modeled as being independent and identically distributed.
- Here, "large" means $N \geq 20n$ or the size of the sample should be 5% or less the size of the population.

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

J. Harner, A. Billings

- Sample Mean \bar{X} is a random variable.

Example:

X = distance to home

[illegible]Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population

Means, $\mu_x = \mu_y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case :

$$\mu_D = \mu_X - \mu_Y$$

- Since \bar{X} is a random variable, \bar{X} has a distribution, which will have a mean and standard deviation. Let $\mu_{\bar{X}}$ be the mean of \bar{X} 's distribution, and $\sigma_{\bar{X}}$ be the the Standard deviation of \bar{X} 's distribution.
- Recall μ and σ are the mean and standard deviation of the distribution from which we obtain the data (the X 's), called the **Parents distribution**.
- $\sigma_{\bar{X}}$, the standard deviation of the distribution of all possible values of \bar{X} , is also called **standard Error of \bar{X}** , denoted as SE(\bar{X}).

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population

Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case :

$\mu_D = \mu_X - \mu_Y$

- The distribution of all possible values of \bar{X} is called the **sampling Distribution** of \bar{X} .
- **But some questions remain:**
 - What values do $\mu_{\bar{X}}$ and $SE(\bar{X})$ assume?
 - what is the shape of the sampling distribution of \bar{X} ?
- We can use the 5-step Method to simulate the probability distribution of possible values of \bar{X} (sampling distribution).

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

**Sec 8.2: The Central Limit
Theorem**

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- See the following diagram¹.




Figure 8.4 Experimental histogram of 10,000 simulated \bar{x} -values from a fair die population, $n = 25$.

- Histogram suggests that:
1. sampling distribution of \bar{X} is approximately normal.
 2. $E(\bar{X}) = \mu_{\bar{X}} \simeq \mu = 3.5$
(from Sec 5.3)
 3. $SE(\bar{X}) = \sigma_{\bar{X}} = 0.34 < \sigma = 1.71$
(from Sec 5.3)

¹refer to page Page: 438 of the Text Book

Central Limit Theorem for \bar{X} (CLT):

- ▶ Will the sampling distribution of \bar{X} always look (approximately) normal?
- ▶ what is the value of $\mu_{\bar{X}}$? $SE(\bar{X})$?
- When the sample size n is large ($n \geq 20$ for this book), we can say something about the sampling distribution of \bar{X} :

CLT: Suppose $X_1, X_2, X_3, \dots, X_n$ is a random sample from a population with mean μ and standard deviation σ . Then the sampling distribution of \bar{X} has

$$\begin{aligned}\mu_{\bar{X}} = E(\bar{X}) &= \mu \\ \sigma_{\bar{X}} = SE(\bar{X}) &= \frac{\sigma}{\sqrt{n}}\end{aligned}$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- IF $n \geq 20$, the sampling distribution of \bar{X} will be approximately normal.

Also

$$P(\bar{X} \leq x) = P\left(Z < \frac{x - \mu}{\sigma/\sqrt{n}}\right)$$

- ▶ What if $n < 20$?

- IF $n < 20$, the sampling distribution of \bar{X} is not necessarily normal, or even approximately normal.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population

Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :

$\mu_D = \mu_X - \mu_Y$

Example:

Suppose random sample of size $n = 25$
 from population with $\mu = 50$ and $\sigma = 10$.
 Describe the Sampling Distribution of \bar{X} .

Population Dist.			Sampling Dist. of \bar{X}		
μ	σ	n	$\mu_{\bar{X}}$	$SE(\bar{X})$	Shape
50	10	16			
50	10	25			
50	10	49			
50	10	100			
75	12	9			
42	24	64			

- $SE(\bar{X}) = \frac{\sigma}{\sqrt{n}}$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

**Sect 8.2: The Central Limit
Theorem**

Sect 6.3: Large-Sample
confidence Interval for the
Population Mean μ


Sect 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sect 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sect 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sect 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.


Sect 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$



Population Distribution — $N, \mu=50, \sigma=10$

\bar{X} 's Distribution ($n=25$) — $N, \mu_{\bar{X}}=50, \sigma_{\bar{X}}=2$

\bar{X} 's Distribution ($n=100$) — $N, \mu_{\bar{X}}=50, \sigma_{\bar{X}}=1$



answer depends on the shape of the distribution of the sampled population, as shown by Figure 6.10. Generally speaking, the greater the skewness of the sampled population distribution, the larger the sample size must be before the normal distribution is an adequate approximation for the sampling distribution of \bar{x} . For most sampled populations, sample sizes of $n \approx 30$ will suffice for the normal approximation to be reasonable. We will use the normal approximation for the sampling distribution of \bar{x} when the sample size is at least 30.

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case :

$$\mu_D = \mu_X - \mu_Y$$

Example:

Suppose weight of an apple follows a probability distribution with mean 100 grams and standard deviation 16 grams.

- a. Assume that the probability distribution is Normal. find the probability that a randomly selected apple weighs more than 105 grams.

$$\begin{aligned} Z &= \frac{X - \mu}{\sigma} \\ &= \frac{105 - 100}{16} \\ &= 0.3125 \end{aligned}$$



Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

$$\begin{aligned}
 P(X > 105) &= P(Z > 0.31) \\
 &= 1 - P(Z < 0.31) \\
 &= 1 - 0.6217 \\
 &= 0.3783
 \end{aligned}$$

- b. No longer assume that apple weight follows a normal distribution.

Find the probability that the mean weight of 64 randomly selected apples exceeds 105 grams

Find $P(\bar{X} > 105)$

Since n is large ($n = 64 \geq 20$), we can use the **CLT**.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

$$\begin{aligned} Z &= \frac{X - \mu}{\frac{\sigma}{\sqrt{n}}} \\ &= \frac{105 - 100}{16/\sqrt{64}} \\ &= 2.50 \end{aligned}$$



$$\begin{aligned} P(\bar{X} > 105) &= P(Z > 2.50) \\ &= 1 - P(Z < 2.50) \\ &= 1 - 0.9938 \\ &= 0.0062 \end{aligned}$$

Chap 8 Notes:
Confidence Interval
Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Confidence Interval Estimation - Introduction:

- We can use \bar{X} to estimate μ . But it is unlikely that \bar{X} is a perfect estimate of μ . (\bar{X} is a point estimate of μ)
- Better to say that μ is between two numbers, a and b, i.e.,

$$a < \mu < b$$

- ▶ This is called an interval estimate of μ
- We can state the interval equivalently as

$$\bar{X} \pm \text{error term}$$

- ▶ What is the **likelihood** that μ really is between a and b? (**Confidence level**)
- ▶ How large is the **error term**?

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small


Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

95% Confidence Interval for μ (Large Sample):

- Recall:** If n is large ($n \geq 20$), \bar{X} 's distribution is approximately normal, with mean $\mu_{\bar{X}} = \mu$ and standard deviation $SE(\bar{X}) = \frac{\sigma}{\sqrt{n}} = \sigma_{\bar{X}}$



Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- So 95% of the values of \bar{X} lie between $\mu - 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right)$ and $\mu + 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right)$

$$P\left(\mu - 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right) < \bar{X} < \mu + 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right)\right) = 0.95$$

which can be rearranged

$$P\left(\bar{X} - 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right) < \mu < \bar{X} + 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right)\right) = 0.95$$

- So the 95% **C.I.** goes from

$$\bar{X} - 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right) \text{ to } \bar{X} + 1.96 \cdot \left(\frac{\sigma}{\sqrt{n}}\right)$$

- Since we usually don't know σ , we use s .

$$\bar{X} - 1.96 \cdot \left(\frac{s}{\sqrt{n}}\right) \text{ to } \bar{X} + 1.96 \cdot \left(\frac{s}{\sqrt{n}}\right)$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Example:

Construct a 95% C.I. for mean weight of full grown dogs,

$$\bar{X} = 44 \text{ lbs}, \quad s = 8 \text{ lbs}, \quad n = 36$$

$$\begin{aligned}\bar{X} \pm 1.96 \cdot \left(\frac{s}{\sqrt{n}} \right) &= 44 \pm 1.96 \cdot \left(\frac{8}{\sqrt{36}} \right) \\ &= 44 \pm 2.61\end{aligned}$$

- So our 95% C.I. is from

41.39 lbs to 46.61 lbs, .

- We are 95% confident that the mean weight of all full grown dogs is between

41.39 lbs and 46.61 lbs, .

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

- ▶ 95% is called the **Confidence Level**.
- ▶ **Interpretation** - See Page: 455 of the Text Book.
- ▶ **Note:** Other Confidence Levels are sometimes used by researchers:

90% Use $Z = 1.645$

99% Use $Z = 2.578$

$$\bar{X} \pm \text{critical value} \times \widehat{SE}(\bar{X})$$

$$\bar{X} \pm Z \times \widehat{SE}(\bar{X})$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- The error term depends on
 1. The **sample size** n
(the larger n is, the smaller the error term.)
 2. The **standard deviation** S
(the smaller S is, the smaller the error term.)
 3. The **confidence level**
(the higher the confidence level, the larger the critical value is; so the larger the error term is.)

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Sample Size for Estimating μ

- The size, n , of the sample depends on
 - Variability of the data.** (standard deviation S)
 - Precision of the estimate**, i.e., the size of the error term $Z\left(\frac{S}{\sqrt{n}}\right)$

Example:

Suppose we wish to estimate μ with a precision of 2 units (i.e., the error term is 2) with 95% confidence. Also suppose we know $S = 12.6$. Find the sample size required.

- We must find n such that

$$Z\left(\frac{S}{\sqrt{n}}\right) = 2$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

(Contd.)

Assuming that n is "large" for a 95% C.I.

$$Z = 1.96$$

So,

$$1.96 \times \left(\frac{S}{\sqrt{n}} \right) = 2$$

$$1.96 \times \left(\frac{12.6}{\sqrt{n}} \right) = 2$$

Solving for n yields

$$n = 152.47$$

which we round up to

$$n = 153 \text{ observations}$$

We need at least 153 observations.

Chap 8 Notes:
Confidence Interval
Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

100(1- α)% C.I. for Binomial Parameter p (Large Sample):

- Suppose we have data from a binomial experiment, but we don't know the value of p.
(**Recall:** p = P(Success))
- We can use a point estimator

$$\hat{p} = \frac{X}{n}$$

to estimate p.

- We can also use an interval estimate, provided

- $n \cdot \hat{p} > 5$ and
- $n \cdot (1 - \hat{p}) > 5$

- The 100(1- α)% C.I. will have the form

$$\hat{p} \pm Z \times SE(\hat{p})$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- We can estimate $SE(\hat{p})$ by

$$SE(\hat{p}) \simeq \sqrt{\frac{\hat{p} \cdot (1 - \hat{p})}{n}}$$

- Thus for a **binomial experiment** with X successes in n trial, a $100(1-\alpha)\%$ C.I. for p is

$$\frac{X}{n} \pm Z \times \sqrt{\frac{\frac{X}{n} \cdot (1 - \frac{X}{n})}{n}}$$

or

$$\hat{p} \pm Z \times \sqrt{\frac{\hat{p} \cdot (1 - \hat{p})}{n}}$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

Example:

In a sample of 200 college students, 120 were female and 80 were male. Construct a 95% C.I. for the proportion p of all college students which are female.

- Let Success = female.

Then X = number of Successes = number of females
and $n = 200$.

$$\begin{aligned}\frac{X}{n} &\pm 1.96 \times \sqrt{\frac{\frac{X}{n} \cdot (1 - \frac{X}{n})}{n}} \\ \frac{120}{200} &\pm 1.96 \times \sqrt{\frac{\frac{120}{200} \cdot (1 - \frac{120}{200})}{200}} \\ 0.6 &\pm 1.96 \times \sqrt{(0.6) \cdot (1 - 0.6)200} \\ 0.6 &\pm 1.96 \times (0.034641) \\ 0.6 &\pm 0.0679\end{aligned}$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

**Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p**

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

So a 95% C.I. on p goes from 0.5321 to 0.6679.

- We are 95% confident that the proportion of female college students is between

0.5321 to 0.6679

Example:

Example 8.20 of the Text Book².

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

**Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p**

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

²refer to Page: 458 of the Text Book

Sample Size for Estimating p:

- Given a specified size for the error term, find n.
- **Recall:** For a 95% C.I. on p, the error term is given by

$$1.96 \times \sqrt{\frac{\hat{p} \cdot (1 - \hat{p})}{n}}$$

Example: (Contd.)

We wish to estimate p within 0.03 (error term) with 95% confidence. Our previous example had,

$$n = 200 \text{ and } X = 120$$

So,

$$\hat{p} = \frac{X}{n} = \frac{120}{200} = 0.6$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

(Contd.)

We must solve for n:

$$1.96 \times \sqrt{\frac{\hat{p} \cdot (1 - \hat{p})}{n}} = 0.03$$

$$1.96 \times \sqrt{(0.6) \cdot (1 - 0.6)n} = 0.03$$

Solving for n yields

$$n = 1024.43$$

We need n = 1025 observations.

- **Note:** IF \hat{p} is not available from a "pilot" study, set \hat{p} to 0.5. Then solve for n as in the example above.

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sect 8.2: The Central Limit
Theorem

Sect 6.3: Large-Sample
confidence Interval for the
Population Mean μ

**Sect 8.4: Large-Sample
Confidence Interval for The
Population Parameter p**

Sect 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sect 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sect 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sect 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Confidence Interval for μ (Small Sample; Data From Normal Population)

I. **Case 1** - σ known

Distribution of \bar{X} when population is Normal.

- Suppose X_1, X_2, \dots, X_n is a random sample from a Normal population with mean μ and standard deviation σ . Then the sampling distribution of \bar{X} is Normal with mean $\mu_{\bar{X}} = \mu$ and $SE(\bar{X}) = \sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$
- IF X_1, X_2, \dots, X_n is a random sample from Normal population with parameters μ and σ , then a $(1-\alpha)100\%$ C.I. is given by

$$\bar{X} \pm Z \cdot SE(\bar{X})$$

i.e.,

$$\bar{X} \pm Z \cdot \left(\frac{\sigma}{\sqrt{n}} \right)$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

Example:

A random sample of 16 observations from a Normal population with known $\sigma = 18$ yields $\bar{X} = 43.2$

Construct a 99% C.I on μ

$$\begin{aligned}\bar{X} &\pm Z \cdot \left(\frac{\sigma}{\sqrt{n}} \right) \\ 43.2 &\pm 2.578 \cdot \left(\frac{18}{\sqrt{16}} \right) \\ 43.2 &\pm 2.578 \cdot (4.5) \\ 43.2 &\pm 11.6\end{aligned}$$

Our 99% C.I. goes from 31.6 to 54.8.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

II. Case 2 - σ unknown

- When σ is unknown, we use a slightly different formula to construct 100(1- α)% C.I.

$$\bar{X} \pm t_{n-1} \times \left(\frac{S}{\sqrt{n}} \right)$$

where the value of t_{n-1} comes from a "t - distribution" with (n-1) degrees of freedom (**Table F**)³.

- The value of t depends on:

1. The Confidence Level

for a 95% C.I. use

$$\frac{\alpha}{2} = \frac{(1 - 0.95)}{2} = 0.025$$

2. The Sample size, n,

which determines the number of degrees of freedom,

$$df = n - 1$$

³refer to Page: 701 of the Text Book

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Example:

Construct a 95% C.I. for the mean weight of apples.

$$\bar{X} = 183 \text{ grams}, S = 14.1 \text{ grams}, n = 16$$

- First, find the value of t.

Here, $\frac{\alpha}{2} = 0.025$ (95% C.I.)
and $df = n-1 = 15$.

So, $t = 2.13$

Next, compute

$$\begin{aligned}\bar{X} &\pm t_{n-1} \times \left(\frac{S}{\sqrt{n}} \right) \\ 183 &\pm 2.13 \times \left(\frac{14.1}{\sqrt{16}} \right) \\ 183 &\pm 2.13 \times (3.525) \\ 183 &\pm 7.5\end{aligned}$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

**Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small**

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

So, we estimate, with 95% confidence, that the mean weight μ of apple is between 175.5 grams and 190.5 grams.

Example: (Contd.)

Suppose n or α were different. Find t :

1. $n = 10$ $\alpha = 0.05$
2. $n = 24$ 95% C.I.
3. $n = 18$ 90% C.I.

Example:

See Example 8.26 of the Text Book⁴.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

⁴refer to Page: 473 of the Text Book

t - distribution

- The t - distribution is actually a family of distributions, each determined by one parameter, the number of **degrees of freedom (df)**.
- All t-distributions possess the following properties:
 1. $\mu = 0$
 2. mound-shaped
 3. symmetric about $\mu = 0$
 4. look like the standard normal distribution (Z), except with a "flatter" middle and "fatter" tails.
 5. $t = \frac{\bar{X} - \mu}{\left(\frac{s}{\sqrt{N}}\right)}$
 6. as n increases,

$$t \longrightarrow Z$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ


Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

**Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small**

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$



$$N, \mu = 0, \sigma = 1$$

t with df = 10

t with df = 20

Degrees Of Freedom (d.f.)

- The number of degrees of freedom equals the number of independent data values used in the calculation, minus the number of restriction placed on the data.

Example:

Suppose we know $n = 5$ and $\bar{X} = 10$.

Then $df = n - 1 = 4$

Since we can "choose" the first 4 data values freely; however the fifth data value must be chosen so that $\bar{X} = 10$.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

For example, the first 4 data values might be

4 6 12 7

Then the fifth value must be

21 (to make $\bar{X} = 10$)

or,

The first 4 data values might be

8 11 15 9

and the fifth value must be

7

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

**Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small**


Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

100(1- α)% C.I. for Difference in Means of Two populations (Unpaired data), $\mu_X - \mu_Y$

- Suppose we obtain random samples from 2 distinct populations. We wish to estimate the difference in population means $\mu_X - \mu_Y$ using a 95% C.I.



- **Note:** The sizes of the two samples may be different.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

- Case I: Small Samples from Normal populations**

100(1- α)% C.I. for $\mu_X - \mu_Y$

$$(\bar{X} - \bar{Y}) \pm t.S_P \cdot \sqrt{\frac{1}{n} + \frac{1}{m}}$$

► where

\bar{X} = mean of sample 1

n = size of sample 1

\bar{Y} = mean of sample 2

m = size of sample 2

t has df = $n + m - 2$

$$S_P = \sqrt{\frac{(n-1).S_X^2 + (m-1).S_Y^2}{n+m-2}}$$

► S_P is called the **"pooled standard deviation"**.

S_X^2 = variance of sample 1

S_Y^2 = variance of sample 2

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Example:

Mental task Problem

$$n = 15 \text{ males} \quad \bar{X} = 18.3 \text{ min} \quad S_X = 3.4 \text{ min}$$

$$m = 16 \text{ females} \quad \bar{Y} = 12.6 \text{ min} \quad S_Y = 2.9 \text{ min}$$

Find a 95% C.I. for mean difference in time between males and females $\mu_X - \mu_Y$.

- First Compute

$$\begin{aligned} S_P &= \sqrt{\frac{(n-1) \cdot S_X^2 + (m-1) \cdot S_Y^2}{n+m-2}} \\ &= \sqrt{\frac{(15-1) \cdot 3.4^2 + (16-1) \cdot 2.9^2}{15+16-2}} \\ &= \sqrt{\frac{287.99}{29}} \\ &= \sqrt{9.931} \\ &= 3.15 \end{aligned}$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

For 95%,

$$\frac{\alpha}{2} = \frac{0.05}{2} = 0.025$$

$$df = n + m - 2 = 29$$

So, $t = 2.045$

$$(\bar{X} - \bar{Y}) \pm t.S_P.\sqrt{\frac{1}{n} + \frac{1}{m}}$$

$$(18.3 - 12.6) \pm 2.045.(3.15).\sqrt{\frac{1}{15} + \frac{1}{16}}$$

$$5.7 \pm 6.442.\sqrt{0.129167}$$

$$5.7 \pm 2.31$$

Our 95% C.I. goes from 3.4 min to 8.0 min.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

**Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$**

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- **This means:**

With 95% Confidence, it takes males between 3.4 min and 8.0 min more than females, "on the average" to complete the task.

Assumptions:

1. Measurements in each population are approximately normally distributed.
2. The population standard deviations are equal, i.e.,

$$\sigma_X = \sigma_Y$$

Example:

See Example 8.28 of the Text Book.⁵

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

⁵refer to the Page: 479 of the Text Book

• Case II: Large Samples

If both sample sizes, n and m are large, we can construct a $100(1 - \alpha)\%$ C.I. on $\mu_X - \mu_Y$, as follows

$$(\bar{X} - \bar{Y}) \pm Z \cdot \sqrt{\frac{S_X^2}{n} + \frac{S_Y^2}{m}}$$

Example:

College statistics professor wants to estimate the difference in performance on EXAM 1 of students who had two or more high school math courses and those students who took fewer than two math courses in high school.

Summary of data as follows

Two or more math courses:

$$n = 35 \quad \bar{X} = 84.2 \quad S_X = 10.2$$

Fewer than 2 math courses:

$$m = 45 \quad \bar{Y} = 73.1 \quad S_Y = 14.3$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

(Contd.)

Find 90% C.I. for the difference in mean scores $\mu_X - \mu_Y$.

- A $100(1-\alpha)\%$ C.I. on $\mu_X - \mu_Y$ is given by,

$$(\bar{X} - \bar{Y}) \pm Z \cdot \sqrt{\frac{S_X^2}{n} + \frac{S_Y^2}{m}}$$

For a 90% C.I., $Z = 1.645$

Then,

$$\begin{aligned}(84.2 - 73.1) &\pm 1.645 \cdot \sqrt{\frac{10.2^2}{35} + \frac{14.3^2}{45}} \\ 11.1 &\pm 1.645 \cdot \sqrt{7.51679} \\ 11.1 &\pm 4.51\end{aligned}$$

Our 90% C.I. goes from 6.6 to 15.6

Chap 8 Notes:
Confidence Interval
Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

**Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$**

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :

$$\mu_D = \mu_X - \mu_Y$$

- This means:**

With 90% Confidence, students who took two or more high school math courses average 6.6 to 15.6 points higher on statistics EXAM 1 than students who take fewer than two high school math courses.

Confidence Interval for The Difference between Two Population Proportions, $p_1 - p_2$

- We may be interested in estimating (C.I.) the difference in proportion for two independent populations, $p_1 - p_2$.
- Suppose n is the size of a sample from population 1, and m is the size of the sample from population 2.
- ▶ If the following conditions hold true:
 1. $n \cdot \hat{p}_1 > 5$
 2. $n \cdot (1 - \hat{p}_1) > 5$
 3. $m \cdot \hat{p}_2 > 5$
 4. $m \cdot (1 - \hat{p}_2) > 5$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

(Contd.)

- then we can construct $100(1-\alpha)\%$ C.I. on $p_1 - p_2$

$$(\hat{p}_1 - \hat{p}_2) \pm Z \cdot \sqrt{\frac{\hat{p}_1 \cdot (1 - \hat{p}_1)}{n} + \frac{\hat{p}_2 \cdot (1 - \hat{p}_2)}{m}}$$

where $\hat{p}_1 = \frac{X}{n}$ and $\hat{p}_2 = \frac{Y}{m}$

X = number of Successes in Sample 1

Y = number of Successes in Sample 2

Example: Flu vaccine

Estimate the difference in proportion of vaccinated adults who get the flu, and the proportion of unvaccinated adults who get the flu. Independent random samples from each population

Vaccinated:

$$n = 50 \quad X = \text{get flu} = 18 \quad \hat{p}_1 = \frac{X}{n} = 0.36$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

Unvaccinated:

$$m = 100 \quad Y = \text{get flu} = 48 \quad \hat{p}_2 = \frac{Y}{m} = 0.48$$

- Note that:

1. $n.\hat{p}_1 = 18 > 5$
2. $n.(1 - \hat{p}_1) = 32 > 5$
3. $m.\hat{p}_2 = 48 > 5$
4. $m.(1 - \hat{p}_2) = 52 > 5$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

Lets' construct 90% C.I. on $p_1 - p_2$, the difference in population proportions:

$$\begin{aligned}(\hat{p}_1 - \hat{p}_2) &\pm Z \cdot \sqrt{\frac{\hat{p}_1 \cdot (1 - \hat{p}_1)}{n} + \frac{\hat{p}_2 \cdot (1 - \hat{p}_2)}{m}} \\(0.36 - 0.48) &\pm 1.645 \cdot \sqrt{\frac{0.36 \cdot (1 - 0.36)}{50} + \frac{0.48 \cdot (1 - 0.48)}{100}} \\-0.12 &\pm 1.645 \cdot \sqrt{0.004608 + 0.002496} \\-0.12 &\pm 1.645 \cdot \sqrt{0.007104} \\-0.12 &\pm 1.645 \cdot (0.084285) \\-0.12 &\pm 0.13865 \\&(-0.25865, 0.01865)\end{aligned}$$

- With 90% Confidence, we estimate that the proportion of vaccinated adults who get flu is between 25.9% less and 1.9% more than the proportion of unvaccinated adults who get flu.

Chap 8 Notes:
Confidence Interval
Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population

Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :

$\mu_D = \mu_X - \mu_Y$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

- ▶ **Note:** since 0 is inside the confidence interval, a difference in proportions equal 0 is plausible. Effectively, the result of this experiment indicates that vaccination was not effective in reducing the chance of getting flu.
- ▶ **Note:** Flu vaccine is effective for the elderly and ill. Flu vaccine's effectiveness is questionable for generally healthy adults.

100(1- α)% C.I. for Mean of Paired Difference: μ_D

- "Real-world" research is often concerned with comparisons between two populations.
- For some research, it might make sense to "pair" the data values. We can then make estimate and draw inferences about the differences between pairs.
- Paired Differences:

Population 1	Population 2	Difference (d)
X_1	Y_1	$d_1 = Y_1 - X_1$
X_2	Y_2	$d_2 = Y_2 - X_2$
\vdots	\vdots	\vdots
X_n	Y_n	$d_n = Y_n - X_n$

Example:

See Example 8.30 of the Text Book⁶.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

⁶refer to Page: 484 of the Text Book

- **Note:** Paired differences also arise in "before" and "after" research.
- 100(1- α)% C.I. for Mean of Paired Difference is given by

$$\bar{d} \pm t. \left(\frac{S_d}{\sqrt{n}} \right)$$

where

$$S_d = \sqrt{\frac{\sum d^2 - \frac{(\sum d)^2}{n}}{n - 1}}$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

Example:

$$S_d = \sqrt{\frac{\sum (d - \bar{d})^2}{n - 1}}$$

Twelve subjects participated in an experiment to study the effectiveness of a certain diet, combined with a program of exercise, in reducing serum cholesterol levels. Table 6.4.1 shows the serum cholesterol levels for the 12 subjects at the beginning of the program (Before) and at the end of the program (After).

Table 6.4.1

Serum Cholesterol Levels for 12 Subjects Before and After Diet-Exercise Program

Subject	Serum Cholesterol		Difference (After-Before)
	Before (X)	After (Y)	
1	201	200	-1
2	231	236	+5
3	221	216	-5
4	260	233	-27
5	228	224	-4
6	237	216	-21
7	326	296	-30
8	235	195	-40
9	240	207	-33
10	267	247	-20
11	284	210	-74
12	201	209	+8

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for The Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

(Contd.)

$$\bar{d} = \frac{\sum d}{n} = \frac{-242}{12} = -20.17$$

$$\begin{aligned} S_d &= \sqrt{\frac{\sum d^2 - \frac{(\sum d)^2}{n}}{n-1}} \\ &= \sqrt{\frac{1076 - \frac{(-242)^2}{12}}{11}} \\ &= \sqrt{535.06} \\ &= 23.131 \end{aligned}$$

$$df = n - 1 = 12 - 1 = 11$$

► For 95% C.I.

$$\alpha = \frac{1 - 0.95}{2} = 0.025$$

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To
The Statistical Estimation
Problem

Sec 8.2: The Central Limit
Theorem

Sec 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sec 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sec 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sec 8.7: Confidence Interval
for The Difference Between
Two Population
Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sec 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :
 $\mu_D = \mu_X - \mu_Y$

(Contd.)

$$\begin{aligned}
 \bar{d} \pm t. \left(\frac{S_d}{\sqrt{n}} \right) &= -20.17 \pm 2.20. \left(\frac{23.131}{\sqrt{12}} \right) \\
 &= -20.17 \pm 2.20.(6.677) \\
 &= -20.17 \pm 14.70
 \end{aligned}$$

So our 95% C.I. on the mean difference is from -34.87 to -5.47.

Interpretation:

You Try!

Assumptions:

The differences are approximately normally distributed.

Chap 8 Notes: Confidence Interval Estimation

Sec 8.1: An Introduction To The Statistical Estimation Problem

Sec 8.2: The Central Limit Theorem

Sec 6.3: Large-Sample confidence Interval for the Population Mean μ

Sec 8.4: Large-Sample Confidence Interval for The Population Parameter p

Sec 8.6: Confidence Intervals for μ When The Population Is Normal and The Sample Size Is Small

Sec 8.7: Confidence Interval for The Difference Between Two Population Means, $\mu_X - \mu_Y$

Sec 8.8: Large-Sample Confidence Interval for The Difference $p_1 - p_2$ Between Two Population Proportions.

Sec 8.9: Confidence Interval for The Difference of Two Population means In Matched-Pairs Design Case : $\mu_D = \mu_X - \mu_Y$

Chap 8 Notes: Confidence Interval Estimation

Sect 8.1: An Introduction To
The Statistical Estimation
Problem

Sect 8.2: The Central Limit
Theorem

Sect 6.3: Large-Sample
confidence Interval for the
Population Mean μ

Sect 8.4: Large-Sample
Confidence Interval for The
Population Parameter p

Sect 8.6: Confidence Intervals
for μ When The Population Is
Normal and The Sample Size
Is Small

Sect 8.7: Confidence Interval
for The Difference Between
Two Population

Means, $\mu_X - \mu_Y$

Sect 8.8: Large-Sample
Confidence Interval for The
Difference $p_1 - p_2$ Between
Two Population Proportions.

Sect 8.9: Confidence Interval
for The Difference of Two
Population means In
Matched-Pairs Design Case :

$\mu_D = \mu_X - \mu_Y$

- **IF** $n \geq 30$ then the values of t in Table F are almost identical to the corresponding Z values in the standard normal Table E.

So for $n \geq 30$, we can construct a C.I. on mean of paired difference using

$$\bar{d} \pm Z. \left(\frac{s}{\sqrt{n}} \right)$$