
MATH1401

Fall 2021

Lecture 19

A/B Testing

Class Checklist

- **Lab 6 – Due Date : Friday 11/5 – 9 PM**

Review: Statistical Significance


Conventions About Inconsistency

- **“Inconsistent with the null”:** The test statistic is in the tail of the empirical distribution under the null hypothesis
 - **“In the tail,” first convention:**
 - The area in the tail is less than 5%
 - The result is “statistically significant”
 - **“In the tail,” second convention:**
 - The area in the tail is less than 1%
 - The result is “highly statistically significant”
-

The P-Value as an Area

Empirical distribution
of the test statistic
under the null
hypothesis

The red dot is the
observed statistic.



Definition of the P -value

Formal name: **observed significance level**

The P -value is the chance,

- if the null hypothesis is true,
 - that the test statistic
 - is equal to the value that was observed in the data
 - or is even further in the direction of the alternative.
-

A/B Testing

Comparing Two Samples

- Compare values of sampled individuals in Group A with values of sampled individuals in Group B.
- Question: Do the two sets of values come from the same underlying distribution?
- Answering this question by performing a statistical test is called **A/B testing**.

(Demo)

The Groups and the Question

- Random sample of mothers of newborns. Compare:
 - (A) Birth weights of babies of mothers who smoked during pregnancy
 - (B) Birth weights of babies of mothers who didn't smoke
 - Question: Could the difference be due to chance alone?
-

Hypotheses

- Null:
 - In the population, the distributions of the birth weights of the babies in the two groups are the same. (They are different in the sample just due to chance.)
 - Alternative:
 - In the population, the babies of the mothers who smoked weigh less, on average, than the babies of the non-smokers.
-

Test Statistic

- Group A: non-smokers
 - Group B: smokers
 - Statistic: Difference between average weights
Group B average - Group A average
 - Negative values of this statistic favor the alternative
- (Demo)
-

The Data



Non-smoker

120 oz



Non-smoker

113 oz



Smoker

128 oz



Smoker

108 oz

...



Non-smoker

...

117 oz

Shuffling Labels Under the Null



Smoker

120 oz



Non-smoker

113 oz



Non-smoker

128 oz



Smoker

108 oz

...



Smoker

117 oz

...

Shuffling Rows

Random Permutation

- `tbl.sample(n)`
 - Table of `n` rows picked randomly with replacement
- `tbl.sample()`
 - Table with same number of rows as original `tbl`, picked randomly with replacement
- `tbl.sample(n, with_replacement = False)`
 - Table of `n` rows picked randomly without replacement
- `tbl.sample(with_replacement = False)`
 - All rows of `tbl`, in random order

(Demo)

Simulating Under the Null

- If the null is true, all rearrangements of labels are equally likely
- Plan:
 - Shuffle all group labels
 - Assign each shuffled label to a birth weight
 - Find the difference between the averages of the two shuffled groups
 - Repeat

(Demo)
