

Spectral Synthesis of Accretion Disk Winds in Active Galactic Nuclei

James Matthews¹

Nick Higginbottom¹, Christian Knigge¹, Knox Long², Stuart Sim³

1- University of Southampton, 2- STScI, 3- Queen's University, Belfast

UNIVERSITY OF
Southampton




Abstract

Outflows are central to our understanding of the connection between active galactic nuclei (AGN) and their host galaxies. In particular, a wind emanating from the accretion disk that fuels the supermassive black hole can provide the kinetic luminosity required for efficient AGN feedback, whilst also having a profound effect on the observational appearance of these systems. To test the effect of these outflows on the observational properties of AGN and QSOs, we have developed a Monte Carlo ionization and radiative transfer code. We have already used our code to demonstrate that a simple, physically motivated disk wind model can produce ultraviolet absorption lines that resemble those seen in broad absorption line quasars (BALQSOs). Recently, we have incorporated Lucy's 'macro-atom' into our code. This allows us to correctly model key spectral features, such as the Lyman-alpha and Balmer emission lines, and has already been successfully applied to a Cataclysmic Variable model. Here, we present the first results of simulations using this scheme, designed to test disk-wind-based AGN unification scenarios. In particular, we address the question whether disk winds may be responsible for the broad emission lines seen in all Type I AGN/QSOs.

Radiative Transfer Code

- We use a Monte Carlo Radiative Transfer Code, named PYTHON.
- Photons are produced by an accretion disk and central object
- Heating and cooling balance include Compton processes



Different ionization & excitation options:


- a modified Saha equation which models the spectrum in a cell
- Full solution of rate matrices for 'macro atom' mode

A Benchmark Model for BALQSOs

- Higginbottom+ 2013 present a benchmark model
- successfully reproduces blue shifted BAL troughs in e.g. CIV**
- First time this has been done in a spectral synthesis code using a self-consistently calculated wind structure

The model has a number of unsolved problems:

- The unabsorbed 2-10keV X-ray luminosity is too weak, at $1\text{e}43 \text{ erg/s}$ (to avoid over-ionizing wind)
- The model fails to produce Lyman alpha emission
- The model fails to produce significant broad emission lines



Implications for AGN feedback

- Disk winds proposed as a possible AGN feedback mechanism- so-called 'Quasar-mode' feedback
- For a disk wind to be responsible for AGN feedback it must have an energetically significant kinetic luminosity, defined as
- $$L_k = \frac{1}{2} \dot{M}_w v^2$$
- Di Matteo+ (2005) and Hopkins & Elvis (2010) require $L_k/L_{bol} = 0.005-0.05$ for effective feedback
- We require the mass loss rate of the wind to be comparable to accretion rate, which gives $L_k/L_{bol} \approx 0.25$
- Consistent with idea that winds may be of sufficient kinetic power to influence the host galaxy**

The Next Step

QSO Macro atom models



- Preliminary results suggest improved treatment of Hydrogen level populations produces more Lyman alpha emission (see right)

Parameter searches: Kinematics and Geometry

- Can a similar model produce BELs with modest kinematic changes?
- Can a model with a more realistic X-ray luminosity produce BAL features?
- Clumping: Is it required? Density fluctuations could lower the ionization parameter and boost the emission measure without changing mass loss


Radiative-Hydro

- Higginbottom+ 2014 treats the Proga & Kallman 2004 model with our radiative transfer code (see density map, right)
- We find the wind is overionized and would not exist in that *particular* state
- Future plan to couple our code with ZEUS for radiative-hydro



Cataclysmic Variable Models

- 'Macro-Atoms': Line Transfer scheme developed by Lucy (2002,2003)
- NLTE solution without simplifying internal atomic transitions- good for recombination lines
- Using macro-atoms produces strong Balmer emission lines and enhances the Lyman alpha emission from the wind (Matthews+, in prep)



References & Contact

- Di Matteo et al. 2005, Nature 443, 604
Higginbottom et al. 2013, MNRAS 436, 1390
Higginbottom et al. 2014, ApJ 789, 19
Hopkins & Elvis 2010, MNRAS 401, 7
Lucy 2002, A&A 384, 725
Lucy 2003, A&A 403, 261
Matthews et al., in prep

Contact: jm8g08@soton.ac.uk



Scan to download pdf