NSIC
NATIONAL SEED INDUSTRY COUNCIL

No.	Year Approved	NSIC Registration Number	Variety Name	Line Name
1	1955	Philippine Seed Board	Tjeremas	
2	1955	Philippine Seed Board	Raminad Strain 1	
3	1955	Philippine Seed Board	Pela	
4	1955	Philippine Seed Board	B.E. 3	
5	1955	Philippine Seed Board	Magsanaya	
6	1955	Philippine Seed Board	Fortuna	
7	1955	Philippine Seed Board	Palawan	
8	1955	Philippine Seed Board	Ginandang Puti	
9	1956	Philippine Seed Board	Azucena	
10	1956	Philippine Seed Board	Nagdami	
11	1956	Philippine Seed Board	Seraup Kedi1139 Str. 482	
12	1957	Philippine Seed Board	Tjahara	
13	1957	Philippine Seed Board	Intan	
14	1957	Philippine Seed Board	Pinulot	
15	1959	Philippine Seed Board	Mangarez	
16	1959	Philippine Seed Board	Milpal	
17	1959	Philippine Seed Board	BPI-76	
18	1960	Philippine Seed Board	Dinalaga	
19	1960	Philippine Seed Board	Nengawan	
20	1960	Philippine Seed Board	PB-121	
21	1961	Philippine Seed Board	AC440 Dr. 260	
22	1962	Philippine Seed Board	Milfor6 (2)	
23	1962	Philippine Seed Board	Nang Thay	

24	1963	Philippine Seed Board	Norelon Str. 340	
25	1963	Philippine Seed Board	FR 178 A	
26	1963	Philippine Seed Board	C-12	
27	1963	Philippine Seed Board	HBDA-2	
28	1963	Philippine Seed Board	Azmil 85	
29	1963	Philippine Seed Board	Texas 317	
30	1963	Philippine Seed Board	C-18	
31	1964	Philippine Seed Board	C-74	
32	1966	Philippine Seed Board	BPI 121-407	
33	1967	Philippine Seed Board	BPI 76-1	
34	1967	Philippine Seed Board	IR-8-288-3	
35	1967	Philippine Seed Board	Azmil 86	
36	1968	Philippine Seed Board	C4-63G	
37	1968	Philippine Seed Board	IR-8	
38	1968	Philippine Seed Board	IR-5	
39	1968	Philippine Seed Board	M9-33B	
40	1968	Philippine Seed Board	BPI-76 (NS)-LOWLAND	
41	1969	Philippine Seed Board	C4-137 (PB76-137)	
42	1969	Philippine Seed Board	IR-20 (IR-532 S76)	
43	1970	Philippine Seed Board	IR-22 (IR 79-2)	
44	1971	Philippine Seed Board	IR-24 (IR-661-1-140-177)	
45	1971	Philippine Seed Board	BPI 121-407	
46	1972	Philippine Seed Board	C22-51	
47	1973	Philippine Seed Board	BPI 3-2	
48	1973	Philippine Seed Board	IR-26 (IR-661-1-140-177)	
49	1973	Philippine Seed Board	C168 (INTAN X BPI-76)	
50	1973	Philippine Seed Board	PARC 2-2	
51	1973	Philippine Seed Board	IR 1561-288-3	
52	1974	Philippine Seed Board	C-22	
53	1974	Philippine Seed Board	BPI 36-DRY SEEDING	
54	1974	Philippine Seed Board	C-168	
55	1975	Philippine Seed Board	IR-28 (IR2061-114-3-2-8-12)	
56	1975	Philippine Seed Board	IR-30 (IR1531-59-1-4)	
57	1975	Philippine Seed Board	IR-32 (IR2070-747-6)	
58	1975	Philippine Seed Board	BPI RI-2 (MRC)	
59	1976	Philippine Seed Board	RP-KN-2	

60	1976	Philippine Seed Board	IR-38	
61	1976	Philippine Seed Board	IR-34	
62	1976	Philippine Seed Board	IR-36	
63	1977	Philippine Seed Board	UPL-Ri-1	
64	1977	Philippine Seed Board	IR-42	
65	1977	Philippine Seed Board	IR-40	
66	1978	Philippine Seed Board	IR-46	
67	1978	Philippine Seed Board	IR-45	
68	1978	Philippine Seed Board	IR-44	
69	1978	Philippine Seed Board	IR-43	
70	1979	Philippine Seed Board	BPI Ri-6	
71	1979	Philippine Seed Board	UPL Ri-3	
72	1979	Philippine Seed Board	BPI Ri-1	
73	1979	Philippine Seed Board	IR-48	
74	1980	Philippine Seed Board	UPL Ri-5	
75	1980	Philippine Seed Board	IR-52	
76	1980	Philippine Seed Board	IR-54	
77	1980	Philippine Seed Board	IR-50	
78	1981	Philippine Seed Board	UPL Ri-7	
79	1981	Philippine Seed Board	BPI Ri-3	
80	1982	Philippine Seed Board	IR-56	
81	1982	Philippine Seed Board	UPL-Ri-4	
82	1983	Philippine Seed Board	BPI-Ri-10	
83	1983	Philippine Seed Board	IR-60	
84	1983	Philippine Seed Board	IR-58	
85	1984	Philippine Seed Board	IR-62	
86	1985	Philippine Seed Board	IR-65	
87	1985	Philippine Seed Board	IR-64	
88	1987	Philippine Seed Board	BPI-Ri-12	
89	1987	Philippine Seed Board	IR-66	

90	1988	Philippine Seed Board	IR-74	
91	1988	Philippine Seed Board	IR-72	
92	1988	Philippine Seed Board	IR-68	
93	1988	Philippine Seed Board	IR-70	
94	1988	Philippine Seed Board	IR-13149	
95	1990	Philippine Seed Board	PSB Ri-1	
96	1991	Philippine Seed Board	PSB Ri-2	
97	1991	Philippine Seed Board	IR-4185-111-3-2-2	
98	1992	Philippine Seed Board	PSB Rc 6	
99	1992	Philippine Seed Board	PSB Rc 8	
100	1992	Philippine Seed Board	PSB Rc 10	
101	1992	Philippine Seed Board	PSB Rc 12	
102	1992	Philippine Seed Board	PSB Rc 14	
103	1993	Philippine Seed Board	PSB Rc 16	
104	1994	NSIC 1994 Rc 18	Ala	IR51672-6662-2-1-2-3
105	1994	NSIC 1994 Rc 20	Chico	IR57301-195-3-3
106	1994	NSIC 1994 Rc 22	Liliw	C3098-41-1-1
107	1994	NSIC 1994 Rc 24	Cagayan	MRC 22095-265
108	1994	NSIC 1994 Rc 26	Magat Hybrid	IR 64 616 H 26 H
109	1995	NSIC 1995 Rc 28	Agno	IR 56381-139-2-2
110	1995	NSIC 1995 Rc 30	Agus	IR 58099-41-2-3
111	1995	NSIC 1995 Rc 32	Jaro	C3 563-B-5-1

112	1995	NSIC 1995 Rc 34	Burdagol	
113	1995	NSIC 1995 Rc 36	Ma-ayon/Ennano II	
114	1995	NSIC 1995 Rc 38	Rinara	
115	1995	NSIC 1995 Rc 40	Chayong	
116	1995	NSIC 1995 Rc 42	Baliwag	MRC22939-24-2
117	1995	NSIC 1995 Rc 44	Gohang	IR59469-B-B-3-2
118	1995	NSIC 1995 Rc 46	Sumadel	IR25976-12-2-2-2-1-1
119	1995	NSIC 1995 Rc 48	Hagonoy	IR9884-84-3-IE-PI
120	1995	NSIC 1995 Rc 50	Bicol	IR5100-AC11-1
121	1997	NSIC 1997 Rc 52	Gandara	IR59682-132-1-1-2
122	1997	NSIC 1997 Rc 54	Abra	IR60819-34-2-1
123	1997	NSIC 1997 Rc 56	Dapitan	PR23416-32
124	1997	NSIC 1997 Rc 58	Mayapa	C2732-10-2-1-1-1
125	1997	NSIC 1997 Rc 60	Tugatog	IR41431-6-1-2-3
126	1997	NSIC 1997 Rc 62	Naguilian	PR23765-23
127	1997	NSIC 1997 Rc 64	Kabacan	IR59552-21-3-2-2
128	1997	NSIC 1997 Rc 66	Agusan	PR23426-66

129	1997	NSIC 1997 Rc 68	Sacobia	IR57515PMI-8-1-1
130	1997	NSIC 1997 Rc 70	Bamban	IR60267-11-2-2-1
131	1997	NSIC 1997 Rc 3	Ginilingan Puti	Ginilingan Puti
132	1997	NSIC 1997 Rc 5	Arayat	IR47686-30-3-2
133	1997	NSIC 1997 Rc 72H	Mestizo Hybrid	IR68284H
134	1999	NSIC 1999 Rc 74	Aklan	C3559-B-18-3-2-1-1
135	1999	NSIC 1999 Rc 76H	Panay	CRH-05
136	1999	NSIC 2000 Rc 78	Pampanga	PR26305-M32
137	2000	NSIC 2000 Rc 80	Pasig	IR62141-114-3-2-2-2
138	2000	NSIC 2000 Rc 82	Peñaranda	IR64683-87-2-2-3-3
139	2000	NSIC 2000 Rc 84	Sipocot	IR65185-3B-8-3-2
140	2000	NSIC 2000 Rc 86	Matnog	IR65195-3B-8-2-3
141	2000	NSIC 2000 Rc 88	Naga	IR527-13-2B-8-2B-1-2
142	2000	NSIC 2000 Rc 90	Buguey	PR25989-2-4B
143	2000	NSIC 2000 Rc 92	Sagada	IR9202-25-1-3
144	2000	NSIC 2000 Rc 94	Hungduan	IR61336-4B-14-3-2

145	2000	NSIC 2000 Rc 96	Ibulao	IR61608-3B-20-2-2-1-1
146	2000	NSIC 2000 Rc 98	Lian	C3419-10-1-2
147	2000	NSIC 2000 Rc 100	Santiago	MRC22965-4955
148	2000	NSIC 2000 Rc 7	Banahaw	PR23336-10
149	2000	IR69726-29-1-2-2-2	Matatag 2 (Stop gap variety)	IR69726-29-1-2-2-2
150	2000	NSIC 2000 Rc 102	Mamburao	IR54068-B-60-1-3-3
151	2001	NSIC 2001 Rc 104	Balili	PR26770-PJ2
152	2001	NSIC 2001 Rc 106	Sumilao	IR61920-3B-22-1
153	2001	NSIC 2001 Rc 108	Anahawan	PR26008-8-4B
154	2001	NSIC 2001 Rc 9	Apo	IR55423-1
155	2001	NSIC 2001 Rc 11	Canlaon	PR23722-20
156	2002	NSIC 2002 Rc 110	Tubigan 1	IR71606-1--4-2-3-1-2
157	2002	NSIC 2002 Rc 112	Tubigan 2	IR72102-4-159-1-3-3-3
158	2002	NSIC 2002 Rc 114	Mestiso 2	IR75207H
159	2002	NSIC 2002 Rc 116H	Mestiso 3	IR75217H
160	2003	NSIC 2003 Rc 118	Matatag 3	IR68305-18-1-1

161	2003	NSIC 2003 Rc 120	Matatag 6	LF-31-28-1
162	2003	NSIC 2003 Rc 122	Angelica	IR61979-138-1-3-2-3
163	2004	NSIC 2004 Rc 124H	Mestiso 4	BIGANTE
164	2004	NSIC 2004 Rc 126H	Mestiso 5	MRH 005 (MAGILAS)
165	2004	NSIC 2004 Rc 128	Mabango 1	PR26645-B-7 (A)
166	2004	NSIC 2004 Rc 130	Tubigan 3	PR30244-AC-9-1
167	2004	NSIC 2004 Rc 132H	Mestiso 6	SL 8H
168	2004	NSIC 2004 Rc 13	Malagkit 1	C4063-2B-7-1-1 (G)
169	2004	NSIC 2004 Rc 15	Malagkit 2	PR25965-14-3 (G)
170	2004	NSIC 2004 Rc 17	Malagkit 3	PR25959-3-3-4-4-1 (G)
171	2005	NSIC 2005 Rc 134	Tubigan 4	PR26873-PJ21-2-1
172	2005	NSIC 2005 Rc 19	Malagkit 4	PR29332-52-1 (G)
173	2005	NSIC 2005 Rc 136H	Mestiso 7	IR78386H
174	2005	NSIC 2005 Rc 138	Tubigan 5	PR26703-3B-PJ25
175	2005	NSIC 2005 Rc 140	Tubigan 6	IR77298-5-6
176	2005	NSIC 2005 Rc 142	Tubigan 7	PR31563-AR32-19-3-3

177	2005	NSIC 2005 Rc 144	Tubigan 8	C6053-B-1-2-1-2
178	2005	NSIC 2005 Rc 146	PJ7	PR26703-3B-PJ7
179	2006	NSIC 2006 Rc 148	Mabango 2	IR71137-243-2-2-3-3
180	2006	NSIC 2006 Rc 150	Tubigan 9	PR27845-PJ27
181	2006	NSIC 2006 Rc 152	Tubigan 10	PR30138-21-2
182	2006	NSIC 2006 Rc 154	Tubigan 11	PR31561-AR32-11-83-1-4-1
183	2006	NSIC 2006 Rc 156	Tubigan 12	C6824-B-1-2
184	2006	NSIC 2006 Rc 158	Tubigan 13	IR77186-122-2-2-3
185	2006	NSIC 2006 Rc 160	Tubigan 14	PR30536-B-48-2
186	2006	NSIC 2006 Rc 162H	Mestiso 8	BIO 401
187	2006	NSIC 2006 Rc 164H	Mestiso 9	Rizalina 28
188	2007	NSIC 2007 Rc 166H	Mestiso 10	PSD 3
189	2007	NSIC 2007 Rc 168H	Mestiso 11	Arize H64
190	2008	NSIC 2008 Rc 170SR	MS 11	IR68333-R-R-B-22
191	2008	NSIC 2008 Rc 172SR	MS 13	IR68144-2B-2-2-3-2

192	2008	NSIC 2008 Rc 174H	Mestiso 12	LP 0331
193	2008	NSIC 2008 Rc 176H	Mestiso 13	LP 0353
194	2008	NSIC 2008 Rc 178H	Mestiso 14	LP 0330
195	2008	NSIC 2008 Rc 180H	Mestiso 15	BCHR 4172
196	2009	NSIC 2009 Rc 182	Salinas 1	IR63307-4B-4-3
197	2009	NSIC 2009 Rc 184	Salinas 2	PR26016-16-B-B-B
198	2009	NSIC 2009 Rc 186	Salinas 3	PR30244-AC-V2
199	2009	NSIC 2009 Rc 188	Salinas 4	PR28524-AC97-55
200	2009	NSIC 2009 Rc 190	Salinas 5	PR25997-B-B-B
201	2009	NSIC 2009 Rc 192	Sahod Ulan 1	IR74371-54-1-1
202	2009	NSIC 2009 Rc 194	Submarino 1	IR84194-139 (IR64-Sub1)
203	2009	NSIC 2009 Rc 196H	Mestiso 16	PR31191H
204	2009	NSIC 2009 Rc 198H	Mestiso 17	PR31204H
205	2009	NSIC 2009 Rc 200H	Mestiso 18	BCS 065

206	2009	NSIC 2009 Rc 202H	Mestiso 19	PRUP 7
207	2009	NSIC 2009 Rc 204H	Mestiso 20	PRUP 9
208	2009	NSIC 2009 Rc 206H	Mestiso 21	IR83199H
209	2009	NSIC 2009 Rc 208H	Mestiso 22	NK 6401
210	2009	NSIC 2009 Rc 210	Mestiso 23	PHB 71
211	2009	NSIC 2009 Rc 212	Tubigan 15	IR77495-10-2-6-2
212	2009	NSIC 2009 Rc 214	Tubigan 16	IR78566-1-2-1-2
213	2009	NSIC 2009 Rc 216	Tubigan 17	PR34141-38-1-J2
214	2009	NSIC 2009 Rc 218SR	Mabango 3	PR34201-2B-3-1-1 (A)
215	2009	NSIC 2009 Rc 220SR	Japonica 1	IR81551-15-1-2-3 (J)
216	2009	NSIC 2009 Rc 222	Tubigan 18	IR78581-12-3-2-2
217	2010	NSIC 2010 Rc 224	Tubigan 19	PR31091-17-3-1
218	2010	NSIC 2010 Rc 226	Tubigan 20	PR33373-10-1-1-B
219	2010	NSIC 2010 Rc 228H	Mestiso 24	HR4842H

220	2010	NSIC 2010 Rc 230H	Mestiso 25	IR82363H
221	2010	NSIC 2010 Rc 232H	Mestiso 26	IR8237H
222	2010	NSIC 2010 Rc 234H	Mestiso 27	NK 5017
223	2010	NSIC 2010 Rc 236H	Mestiso 28	SW 836
224	2011	NSIC 2011 Rc 238	Tubigan 21	IR78555-68-3-3-3
225	2011	NSIC 2011 Rc 240	Tubigan 22	PR31132-B-1-1-1-3-3
226	2011	NSIC 2011 Rc 21SR	Malagkit 5	PR25482-4-2 (G)
227	2011	NSIC 2011 Rc 242SR	Japonica 2	IR83515-30-1-1-3 (J)
228	2011	NSIC 2011 Rc 244H	Mestiso 29	PR36020H

229	2011	NSIC 2011 Rc 246H	Mestiso 30	IR84714H
230	2011	NSIC 2011 Rc 248H	Mestiso 31	IR80637H
231	2011	NSIC 2011 Rc 250H	Mestiso 32	IR81949H
232	2011	NSIC 2011 Rc 252H	Mestiso 33	PAC 809
233	2011	NSIC 2011 Rc 254H	Mestiso 34	PAC 835
234	2011	NSIC 2011 Rc 256H	Mestiso 35	BSG 0802
235	2011	NSIC 2011 Rc 258H	Mestiso 36	MR 5

236	2011	NSIC 2011 Rc 260H	Mestiso 37	SW 804
237	2011	NSIC 2011 Rc 262 H	Mestiso 38	PHILSCAT 3-1
238	2011	NSIC 2011 Rc 264H	Mestiso 39	MR 1
239	2011	NSIC 2011 Rc 266H	Mestiso 40	DRH 257
240	2011	NSIC 2011 Rc 268H	Mestiso 41	DRH 1531
241	2011	NSIC 2011 Rc 270H	Mestiso 42	DRH 007
242	2011	NSIC 2011 Rc 272	Sahod Ulan 2	PR34363-4-Pokkali/ AC-45-M5R-19

243	2011	NSIC 2011 Rc 274	Sahod Ulan 3	IR81412-B-B-82-1
244	2011	NSIC 2011 Rc 276	Sahod Ulan 4	C8108-B-10-2-2-1
245	2011	NSIC 2011 Rc 278	Sahod Ulan 5	IR81023-B-116-1-2
246	2011	NSIC 2011 Rc 280	Sahod Ulan 6	IR72667-16-1-B-B-3
247	2011	NSIC 2011 Rc 282	Sahod Ulan 7	C8231-B-1-1
248	2011	NSIC 2011 Rc 284	Sahod Ulan 8	IR74963-262-5-1-3-3
249	2011	NSIC 2011 Rc 286	Sahod Ulan 9	C6392-2B-3-3-1-2

250	2011	NSIC 2011 Rc 288	Sahod Ulan 10	PR25769-B-9-1
251	2011	NSIC 2011 Rc 23	Katihan 1	IR79913-B-176-B-4
252	2011	NSIC 2011 Rc 290	Salinas 6	PR28377-AC97-54
253	2011	NSIC 2011 Rc 292	Salinas 7	PR30244-AC-V19
254	2011	NSIC 2011 Rc 294	Salinas 8	PR28378-AC96-36
255	2011	NSIC 2011 Rc 296	Salinas 9	IR71896-3R-8-3-1
256	2012	NSIC 2012 Rc 298	Tubigan 23	PR34159-13-1
257	2012	NSIC 2012 Rc 300	Tubigan 24	PR31379-2B-10-1-2-1-2
258	2012	NSIC 2012 Rc 302	Tubigan 25	IR79643-39-2-2-3
259	2012	NSIC 2012 Rc 304SR	Japonica 3	PR34126-B-1(J)

260	2012	NSIC 2012 Rc 306H	Mestiso 43	MR 18
261	2013	NSIC 2013 Rc 308	Tubigan 26	PR35766-B-24-3
262	2013	NSIC 2013 Rc 310	Mestiso 44	H6129
263	2013	NSIC 2013 Rc 312H	Mestiso 45	INH97158
264	2013	NSIC 2013 Rc 314H	Mestiso 46	INH10001
265	2013	NSIC 2013 Rc 316H	Mestiso 47	PR36559H
266	2013	NSIC 2013 Rc 318H	Mestiso 48	PR35664H

267	2013	NSIC 2013 Rc 320H	Mestiso 49	PR36444H
268	2013	NSIC 2013 Rc 322H	Mestiso 50	P2010-31
269	2013	NSIC 2013 Rc 324	Salinas 10	PR31607-2-B-B-B-B
270	2013	NSIC 2013 Rc 326	Salinas 11	IR84048-B-B-1-1
271	2013	NSIC 2013 Rc 328	Salinas 12	IR62700-2B-9-2-3
272	2013	NSIC 2013 Rc 330	Salinas 13	PR37435-30-1
273	2013	NSIC 2013 Rc 332	Salinas 14	PR38566-Wagwag-V9-3-2-15-2
274	2013	NSIC 2013 Rc 334	Salinas 15	IR83410-6-B-4-1-1-2
275	2013	NSIC 2013 Rc 336	Salinas 16	IR84095-AJY3-8-SD01-B
276	2013	NSIC 2013 Rc 338	Salinas 17	PR30665-1B-1-B-B-Cg
277	2013	NSIC 2013 Rc 340	Salinas 18	IR84096-AY4-2-SD04-B
278	2013	NSIC 2103 Rc 342SR	Mabango 4	PR37299-31-69-16-2-1-2 (A)

279	2013	NSIC 2013 Rc 344SR	Mabango 5	IR04A285 (A)
280	2013	NSIC 2013 Rc 346	Sahod Ulan 11	PR34350-4-POKKALI-24-M5R-10 (DrS 88)
281	2013	NSIC 2013 Rc 348	Sahod Ulan 12	IR81047-B-106-2-4
282	2013	NSIC 2013 Rc 350H	Mestiso 51	SW 82
283	2014	NSIC 2014 Rc 25	Katihan 2	IR82635-B-B-47-2
284	2014	NSIC 2014 Rc 27	Katihan 3	IR86857-101-2-1-3
285	2014	NSIC 2014 Rc 29	Katihan 4	IR83140-B-36-B
286	2014	NSIC 2014 Rc 352	Tubigan 27	IR80694-44-1-2-2
287	2014	NSIC 2014 Rc 354	Tubigan 28	PR37274-6-33-9-1-2(AR)

288	2014	NSIC 2014 Rc 356	Tubigan 29	IR78585-98-2-2-1
289	2014	NSIC 2014 Rc 358	Tubigan 30	PR35789-B-1-1-1
290	2014	NSIC 2014 Rc 360	Tubigan 31	IR80894-18-2-2-3
291	2014	NSIC 2014 Rc 362H	Mestiso 52	PHDR 0913
292	2014	NSIC 2014 Rc 364H	Mestiso 53	PHDR 0911
293	2014	NSIC 2014 Rc 366H	Mestiso 54	PHDR 0912
294	2014	NSIC 2014 Rc 368H	Mestiso 55	PR36474H
295	2014	NSIC 2014 Rc 370H	Mestiso 56	IR81955H

296	2014	NSIC 2014 Rc 372H	Mestiso 57	BIO 452
297	2014	NSIC 2014 Rc 374H	Mestiso 58	PHDR 0910
298	2014	NSIC 2014 Rc 376H	Mestiso 59	PHDR0914
299	2014	NSIC 2014 Rc 378H	Mestiso 60	PAC 801
300	2014	NSIC 2014 Rc 380H	Mestiso 61	IR81265H
301	2014	NSIC 2014 Rc 382H	Mestiso 62	PHDR 1104
302	2014	NSIC 2014 Rc 384H	Mestiso 63	LP 106

303	2014	NSIC 2014 Rc 386H	Mestiso 64	PHDR 1102
304	2014	NSIC 2014 Rc 388H	Mestiso 65	PHDR 1101
305	2014	NSIC 2014 Rc 390	Salinas 19	IR83140-B-28-B
306	2014	NSIC 2014 Rc 392	Salinas 20	IR84675-58-4-1-B-B
307	2015	NSIC 2015 Rc 394	Tubigan 32	PR36723-B-B-3-3-3
308	2015	NSIC 2015 Rc 396	Tubigan 33	PR35766-B-24-1
309	2015	NSIC 2015 Rc 398	Tubigan 34	C8112-B-4-3-2-1-1
310	2015	NSIC 2015 Rc 400	Tubigan 35	IR06A150

311	2015	NSIC 2015 Rc 402	Tubigan 36	PR36930-B-7-3
312	2015	NSIC 2015 Rc 404H	Mestiso 66	LP 2096
313	2015	NSIC 2015 Rc 406H	Mestiso 67	INH10002
314	2015	NSIC 2015 Rc 408H	Mestiso 68	IR82391H
315	2015	NSIC 2015 Rc 410H	Mestiso 69	INH11001
316	2015	NSIC 2015 Rc 412H	Mestiso 70	LP 205
317	2015	NSIC 2015 Rc 414SR	Japonica 4	PR27958-135-1-2-(J)
318	2015	NSIC 2015 Rc 416	Sahod Ulan 13	PR33382-25-1-1-B-B (Raeline)

319	2015	NSIC 2015 Rc 418	Sahod Ulan 14	C9270-B-3-1-3-2
320	2015	NSIC 2015 Rc 420	Sahod Ulan 15	IR83383-B-B-129-4
321	2015	NSIC 2015 Rc 422	Sahod Ulan 16	PR38583-IR64-AC97WP-135-36-1-6
322	2015	NSIC 2015 Rc 424	Sahod Ulan 17	PR30952-AC10-SSD-35
323	2015	NSIC 2015 Rc 426	Sahod Ulan 18	PR30245-IR64-AC97WD-135-44-3-4
324	2015	NSIC 2015 Rc 428	Sahod Ulan 19	IR86781-3-3-1-1
325	2015	NSIC 2015 Rc 430	Sahod Ulan 20	PR37416-18-1
326	2016	NSIC 2016 Rc 432H	Mestiso 71	IR82386H
327	2016	NSIC 2016 Rc 434	Sahod Ulan 21	IR83142-B-19-B

328	2016	NSIC 2016 Rc 436	Tubigan 37	GSR IR1-5-S6-S3-D1
329	2016	NSIC 2016 Rc 438	Tubigan 38	PR37241-3-1-2-1-1
330	2016	NSIC 2016 Rc 440	Tubigan 39	PR36720-17-1-2-1
331	2016	NSIC 2016 Rc 442	Tubigan 40	IR04A115
332	2016	NSIC 2016 Rc 444H	Mestiso 72	PHDR 2112
333	2016	NSIC 2016 Rc 446H	Mestiso 73	PRUP 10
334	2016	NSIC 2016 Rc 448H	Mestiso 74	PHDR 2114

335	2016	NSIC 2016 Rc 450H	Mestiso 75	PHDR 2115
336	2016	NSIC 2016 Rc 452H	Mestiso 76	PHDR 2116
337	2016	NSIC 2016 Rc 454H	Mestiso 77	IR81958H
338	2016	NSIC 2016 Rc 456H	Mestiso 78	LPP 937
339	2016	NSIC 2016 Rc 458H	Mestiso 79	LP 357
340	2016	NSIC 2016 Rc 460	Zinc Rice	IR10M300
341	2016	NSIC 2016 Rc 462	Salinas 21	PR30025-99AC-WSAL-1086
342	2016	NSIC 2016 Rc 464	Salinas 22	IR86385-38-1-1-B
343	2016	NSIC 2016 Rc 466	Salinas 23	IR84089-7-3-AJY1-B

344	2016	NSIC 2016 Rc 468	Salinas 24	IR06M139
345	2016	NSIC 2016 Rc 470	Salinas 25	PR30245-IR64-ID-18-1-4
346	2016	NSIC 2016 Rc 472	Sahod Ulan 22	PR30245-AC-123
347	2016	NSIC 2016 Rc 474	Sahod Ulan 23	IR91648-B-20-B-3-1
348	2016	NSIC 2016 Rc 476	Sahod Ulan 24	C9305-B-9-2
349	2016	NSIC 2016 Rc 478	Sahod Ulan 25	IR84878-B-60-4-1
350	2016	NSIC 2016 Rc 480	GSR 8	GSR IR1-8-S6-S3-Y2
351	2017	NSIC 2017 Rc 482SR	Japonica 5	PR39494-NIPPONBARE-25kR-29-1-1-1-1 (J)
352	2017	NSIC 2017 Rc 484SR	Japonica 6	IR94746-24-3-2-2-2 (J)

353	2017	NSIC 2017 Rc 31SR	Malagkit 6	PR34859-B-4-1-1-2-1 (G)
354	2017	NSIC 2017 Rc 486H	Mestiso 80	LP 534
355	2017	NSIC 2017 Rc 488H	Mestiso 81	SL-19H
356	2017	NSIC 2017 Rc 490H	Mestiso 82	PHILSCAT 9
357	2017	NSIC 2017 Rc 492H	Mestiso 83	P2014-77
358	2017	NSIC 2017 Rc 494H	Mestiso 84	INH13144
359	2017	NSIC 2017 Rc 496H	Mestiso 85	PH11017

360	2017	NSIC 2017 Rc 498H	Mestiso 86	PH11014
361	2017	NSIC 2017 Rc 500H	Mestiso 87	PH11020
362	2017	NSIC 2017 Rc 502H	Mestiso 88	INH07028
363	2017	NSIC 2017 Rc 504H	Mestiso 89	IR81255H
364	2018	NSIC 2018 Rc 506	Tubigan 41	C9301-B-2-1-2-2
365	2018	NSIC 2018 Rc 508	Tubigan 42	IR82572-28-3-1-1-2
366	2018	NSIC 2018 Rc 510	Tubigan 43	C9222-B-2-2-1-2
367	2018	NSIC 2018 Rc 512	Tubigan 44	HHZ3-SAL13-Y1-SAL1

368	2018	NSIC 2018 Rc 514	Tubigan 45	GSR IR1-21-Y4-Y2-Y1
369	2018	NSIC 2018 Rc 516H	Mestiso 90	LP 402
370	2018	NSIC 2018 Rc 518H	Mestiso 91	PH1 15029
371	2018	NSIC 2018 Rc 520H	Mestiso 92	INH10008
372	2018	NSIC 2018 Rc 522H	Mestiso 93	BIO 453
373	2018	NSIC 2018 Rc 524H	Mestiso 94	BCS 065 Plus
374	2018	NSIC 2018 Rc 526H	Mestiso 95	LPP 952
375	2018	NSIC 2018 Rc 528	Salinas 26	PR38456-B-12-1-1-B

376	2018	NSIC 2018 Rc 530	Salinas 27	IR85922-4-1-1-AJY1-2-2-B
377	2018	NSIC 2018 Rc 532	Salinas 28	IR84096-AJY5-3-AJY1-B-AJY3-B
378	2018	NSIC 2018 Rc 534	Salinas 29	GSR-IR1-12-D10-S1-D1
379	2018	NSIC 2018 Rc 536	Salinas 30	PR38121-B-27-B-B-B
380	2018	NSIC 2018 Rc 538H	Mestiso 96	LPP 507
381	2018	NSIC 2018 Rc 540H	Mestiso 97	SL-20H
382	2018	NSIC 2018 Rc 542H	Mestiso 98	Rashmi

383	2018	NSIC 2018 Rc 544H	Mestiso 99	PRUP 11
384	2018	NSIC 2018 Rc 546H	Mestiso 100	BCS 064 Plus (Essentially derived from Mestiso 11)
385	2018	NSIC 2018 Rc 548H	Mestiso 101	Bigante Plus (Essentially derived from Mestiso 4)
386	2019	NSIC 2019 Rc 550H	Mestiso 102	LP 412
387	2019	NSIC 2019 Rc 552H	Mestiso 103	PRUP 12 (Essentially derived from Mestiso 19)
388	2019	NSIC 2019 Rc 554	Salinas 31	GSR IR2-6-R4-S3-Y2
389	2019	NSIC 2019 Rc 556	Salinas 32	GSR IR2-19-R2-SU2-R2

390	2019	NSIC 2019 Rc 558	Salinas 33	PR34350-2-POKKALI-AC-24-M_5R-8-DrS87
391	2019	NSIC 2019 Rc 560	Cordillera 1	PR28705-B-3-86-6-1-1-2-10
392	2019	NSIC 2019 Rc 562	Cordillera 2	PR29399-B-2-2-1
393	2019	NSIC 2019 Rc 564	Cordillera 3	PR34131-B-11-1
394	2019	NSIC 2019 Rc 566	Cordillera 4	IR73688-82-3
395	2019	NSIC 2019 Rc 568	Sahod Ulan 26	PR30952-AC10SSD-164

396	2019	NSIC 2019 Rc 570	Sahod Ulan 27	IR101465-5-33
397	2019	NSIC 2019 Rc 572	Sahod Ulan 28	PR41395-NSIC Rc9-IV-M2009DS-50-1-4
398	2019	NSIC 2019 Rc 574	Sahod Ulan 29	PR38121-B-7-B-B-B
399	2019	NSIC 2019 Rc 576	Sahod Ulan 30	IR102860-12-6-B-B
400	2019	NSIC 2019 Rc 578	Sahod Ulan 31	PR38075-B-4-B-B-B

401	2019	NSIC 2019 Rc 580	Tubigan 46	IR09A136
402	2019	NSIC 2019 Rc 582	Tubigan 47	C9386-B-7-2-3
403	2019	NSIC 2019 Rc 584SR	Japonica 7	IR101055-52-2-1-2-2 (J)
404	2019	NSIC 2019 Rc 586H	Mestiso 104	SYT 008

405	2019	NSIC 2019 Rc 588H	Mestiso 105	P2017-PA4
406	2020	NSIC 2020 Rc 590	Submarino 2	IR13F265
407	2020	NSIC 2020 Rc 592	Sahod Ulan 32	PR33374-28-1-5-B-B (Raeline 7)
408	2020	NSIC 2020 Rc 594	Sahod Ulan 33	PR33481-B-2-B (Raeline 5)
409	2020	NSIC 2020 Rc 596	Sahod Ulan 34	C9305-B-9-2-1
410	2020	NSIC 2020 Rc 598	Sahod Ulan 35	IR93349-3-B-20-5-20-IRGA-2RGA-1-B-B
411	2020	NSIC 2020 Rc 600	PangMainit 1	PR40330-4-2-7-1-2

412	2020	NSIC 2020 Rc 602	PangMainit 2	PR39923-B-3-B-1-1
413	2020	NSIC 2020 Rc 604	Salinas 34	PR45167-B-B-18-1
414	2020	NSIC 2020 Rc 606	Salinas 35	IR91716-60-BAY-1-3-2
415	2020	NSIC 2020 Rc 608	Salinas 36	IR15T1030
416	2020	NSIC 2020 Rc 610	Salinas 37	IR15T1094

417	2020	NSIC 2020 Rc 612	Salinas 38	GSR IR2-DQ25-L1-C1
418	2020	NSIC 2020 Rc 614H	Mestiso 106	LP 488
419	2020	NSIC 2020 Rc 616H	Mestiso 107	LP 702
420	2020	NSIC 2020 Rc 618H	Mestiso 108	LP 727
421	2021	NSIC 2021 Rc 620H	Mestiso 109 (HYVAR S26)	SW 1210

422	2021	NSIC 2021 Rc 622	Tubigan 48	PR38168-2B-3-1-3-1-1-2
423	2021	NSIC 2021 Rc 624	Tubigan 49	PR39142-10-3-2-1-1
424	2021	NSIC 2021 Rc 626	Tubigan 50	PR40432-14-2-1-B
425	2021	NSIC 2021 Rc 628	Tubigan 51	IR11A307
426	2021	NSIC 2021 Rc 630	Tubigan 52	IR11N313
427	2021	NSIC 2021 Rc 632	Tubigan 53	IR12L130
428	2021	NSIC 2021 Rc 634	Tubigan 54	GSR-IR2-9-L1-L3-L2
429	2021	NSIC 2021 Rc 636	Tubigan 55	GSR-IR1-5-L10-U1-L3
430	2021	NSIC 2021 Rc 638SR	Pigmented 1	PR35034-B-3-2-1-1-4-1-2-1-1-2-1 (R/NG)
431	2021	NSIC 2021 Rc 640SR	Pigmented 2	PR47521-B-3-2-1-4 (BL/NG)
432	2021	NSIC 2021 Rc 642SR	Pigmented 3	PR33277-2B-5-2-2-2-3-2-1-2-1-1-2-3-3 (R/NG)

433	2021	NSIC 2021 Rc 644SR	Pigmented 4	PR43172-2B-19-1-2 (R/NG)
434	2021	NSIC 2021 Rc 646SR	Pigmented 5	PR46441-B-B-12-2-1-1 (BL-NG)
435	2021	NSIC 2021 Rc 648	Zinc Rice 2	IR93337:37-B-15-22-1RGA-2RGA-1-B
436	2021	NSIC 2021 Rc 650	Sahod Ulan 36	IR16L1860
437	2021	NSIC 2021 Rc 652H	Mestiso 110	LP 624
438	2021	NSIC 2021 Rc 654H	Mestiso 111	LP 862
439	2021	NSIC 2021 Rc 656H	Mestiso 112	LP 524
440	2021	NSIC 2021 Rc 658H	Mestiso 113	LP 616
441	2021	NSIC 2021 Rc 660H	Mestiso 114	LP 763

442	2021	NSIC 2021 Rc 662H	Mestiso 115	LP 869
443	2021	NSIC 2021 Rc 664H	Mestiso 116	P2019-PJ6
444	2021	NSIC 2021 Rc 666H	Mestiso 117	LAV 102
445	2021	NSIC 2021 Rc 668H	Mestiso 118	PAC 801 Elite
446	2022	NSIC 2022 Rc 670	Tubigan 56	PR39502-B-7-98
447	2022	NSIC 2022 Rc 672	Tubigan 57	PR40846-92-2-2
448	2022	NSIC 2022 Rc 674	Tubigan 58	IR10A135
449	2022	NSIC 2022 Rc 676	Tubigan 59	IR101465-10-6
450	2022	NSIC 2022 Rc 678	Tubigan 60	IR13A515
451	2022	NSIC 2022 Rc 680	Tubigan 61	C9670-B-9-2-2-1
452	2022	NSIC 2022 Rc 682GR2E	Malusog 1	IR112017-GR2-E:23-40-51-B-22-26-B-B-B

453	2022	NSIC 2022 Rc 684	Zinc Rice 3	IR115928-B-42-1-3- B
454	2022	NSIC 2022 Rc 686	Salinas 39	PR48421-FR13A-IVM2012DS-1-7-4
455	2022	NSIC 2022 Rc 688H	Mestiso 119	AZ8433DT
456	2022	NSIC 2022 Rc 690H	Mestiso 120	IR134554H
457	2022	NSIC 2022 Rc 692H	Mestiso 121	LAV 105
458	2022	NSIC 2022 Rc 694H	Mestiso 122	LAV 107
459	2022	NSIC 2022 Rc 696H	Mestiso 123	LP906
460	2022	NSIC 2022 Rc 698H	Mestiso 124	LP996
461	2022	NSIC 2022 Rc 700H	Mestiso 125	P10
462	2022	NSIC 2022 Rc 702H	Mestiso 126	P17

463	2022	NSIC 2022 Rc 704H	Mestiso 127	P18
464	2022	NSIC 2022 Rc 706H	Mestiso 128	P20
465	2022	NSIC 2022 Rc 708H	Mestiso 129	PHH18011
466	2022	NSIC 2022 Rc 710H	Mestiso 130	P2021-P37
467	2022	NSIC 2022 Rc 712H	Mestiso 131	P2021-300
468	2022	NSIC 2022 Rc 714H	Mestiso 132	PR48800H
469	2022	NSIC 2022 Rc 718H	Mestiso 134	SL-39H
470	2022	NSIC 2022 Rc 720H	Mestiso 135	SL-43H
471	2022	NSIC 2022 Rc 722H	Mestiso 136	SL-201H
472	2022	NSIC 2022 Rc 724H	Mestiso 137	SL-422H
473	2022	NSIC 2022 Rc 726H	Mestiso 138	SL-606H

474	2022	NSIC 2022 Rc 728H	Mestiso 139	SL-631H
475	2023	NSIC 2023 Rc 730 SR	Pigmented 6	PR44867-2B-12-1 (RNG)
476	2023	NSIC 2023 Rc 732	Sahod Ulan 37	PR42837-NSIC Rc 222-48-B-RTD9-5-DTR2
477	2023	NSIC 2023 Rc 734	Sahod Ulan 38	C7645-2B-8-2
478	2023	NSIC 2023 Rc 736	Sahod Ulan 39	IR17L1430
479	2023	NSIC 2023 Rc 738	Tubigan 62	PR33481-5-B-2-B-B
480	2023	NSIC 2023 Rc 740	Tubigan 63	PR48421-FR13A-IVM2012DS-5-8-1
481	2023	NSIC 2023 Rc 742H	Mestiso 140 (Biorice Bonus)	Biorice 888
482	2023	NSIC 2023 Rc 744H	Mestiso 141 (Biorice Bounty)	Biorice 432
483	2023	NSIC 2023 Rc 746H	Mestiso 142 (Harvest King 158)	TN 905
484	2023	NSIC 2023 Rc 748H	Mestiso 143 (Arize 6585)	INH16085

485	2023	NSIC 2023 Rc 750H	Mestiso 144	LP 777
486	2023	NSIC 2023 Rc 752H	Mestiso 145	LP 924
487	2023	NSIC 2023 Rc 754H	Mestiso 146 (PHB 91)	P2022-9061
488	2023	NSIC 2023 Rc 756H	Mestiso 147	SL-203H
489	2023	NSIC 2023 Rc 758H	Mestiso 148	SL-222H
490	2024	NSIC 2024 Rc 760SR	Zinc Rice 4	IR17M1467
491	2024	NSIC 2024 Rc 762SR	Zinc Rice 5	IR17M1658
492	2024	NSIC 2024 Rc 764SR	Zinc Rice 6	IR17M1653
493	2024	NSIC 2024 Rc 33	Katihan 5	C9656-B-11-1-1-1-1(Upland)
494	2024	NSIC 2024 Rc 766	Salinas 40	PR42837-NSIC Rc222-18-B-RTD-1-21-DRT1
495	2024	NSIC 2024 Rc 768H	Mestiso 149	IR142464H

496	2024	NSIC 2024 Rc 770H	Mestiso 150	LP 894H
497	2024	NSIC 2024 Rc 772H	Mestiso 151	LP 994H
498	2024	NSIC 2024 Rc 774H	Mestiso 152	SL-27H
499	2024	NSIC 2024 Rc 776H	Mestiso 153	SL-771H
500	2024	NSIC 2024 Rc 778H	Mestiso 154	Vanbee 75
501	2025	NSIC 2025 Rc 780H	Mestiso 155 (Milagros Harvester 101)	Winall 111
502	2025	NSIC 2025 Rc 782H	Mestiso 156 (KRHP 299)	ESRH 003

503	2026	NSIC 2026 Rc 784	Tubigan 64 (NSIC Rc 222 Malambot)	IR20X1019
504	2026	NSIC 2026 Rc 786H	Mestiso 157 (DADI - 1H)	DADI-1H
505	2026	NSIC 2026 Rc 788H	Mestiso 158 (DADI - 2H)	DADI-2H
506	2026	NSIC 2026 Rc 790H	Mestiso 159 (DADI - 3H)	DADI-3H
507	2026	NSIC 2026 Rc 792H	Mestiso 160 (H380)	G211
508	2026	NSIC 2026 Rc 794H	Mestiso 161 (AZ8445DT)	INH16019

509	2026	NSIC 2026 Rc 796H	Mestiso 162 (AZ2241ST)	INH18071
510	2026	NSIC 2026 Rc 798H	Mestiso 163 (AZ5532ST)	INH220221
511	2026	NSIC 2026 Rc 800H	Mestiso 164 (IR138982H)	IR138982H
512	2026	NSIC 2026 Rc 802H	Mestiso 165 (LP 756)	LP 756
513	2026	NSIC 2026 Rc 804H	Mestiso 166 (LP 757)	LP 757

514	2026	NSIC 2026 Rc 806H	Mestiso 167 (LP 794)	LP 794
515	2026	NSIC 2026 Rc 808H	Mestiso 168 (MIP 5053)	MIP 5053
516	2026	NSIC 2026 Rc 810H	Mestiso 169 (MIP 5057)	MIP 5057
517	2026	NSIC 2026 Rc 812H	Mestiso 170 (MIP 5065)	MIP 5065
518	2026	NSIC 2026 Rc 814H	Mestiso 171 (TH3-5)	TH3-5

519	2026	NSIC 2026 Rc 816H	Mestiso 172 (PhilSCAT 21)	P21
520	2026	NSIC 2026 Rc 818H	Mestiso 173 (PhilSCAT 22)	P22
521	2026	NSIC 2026 Rc 820H	Mestiso 174 (PRUP 15)	PRUP 15
522	2026	NSIC 2026 Rc 822H	Mestiso 175 (RTP740FP)	RT2024-127
523	2026	NSIC 2026 Rc 824H	Mestiso 176 (RTP750FP)	RT2024-134

524	2026	NSIC 2026 Rc 826H	Mestiso 177 (UH 309)	UH 309
525	2026	NSIC 2026 Rc 828H	Mestiso 178 (WSTAR-78)	WSSTAR 78
526	2026	NSIC 2026 Rc 830H	Mestiso 179 (WSTAR-98)	WSSTAR 98

Abbreviations:
DAS = Days After Sowing
DS = Dry Season
WS = Wet Season
DSR = Direct Seeded Rice
DWSR = Direct Wet Seeded Rice
TPR = Transplanted Rice
I = Intermediate
R = Resistant
S = Susceptible
MR = Moderately Resistant
MS = Moderately Susceptible
WSB = White Stemborer
YSB = Yellow Stemborer
NCT 1 = National Cooperative Trials Phase 1
MAT = Multi-Location Adaptation Trial
IRRI = International Rice Research Institute
PhilRice = Philippine Rice Research Institute
CBES = Central Bicol Experiment Station
UPLB = University of the Philippines Los Baños

REGISTERED RICE VARIETIES

Oryza sativa L.

Type of Variety	Owner/ Breeder	Average Yield (kg	
		TPR	
		DS	WS
	BPI		
	BPI		
	BPI		
	BPI		
	BPI		
	BPI		
	BPI		
	BPI		

	BPI		
	UPLB		
	UPLB		
	UPLB		
	UPLB		
	UPLB		
	BPI		
	BPI		
	IRRI		
	UPLB		
	UP		
	IRRI		
	IRRI		
	BPI		
	UP		
	IRRI		
	IRRI		
	IRRI		
	BPI		
	UP		
	BPI		
	IRRI		
	UP		
	PARC		
	IRRI		
	UP		
	BPI		
	UP		
	IRRI		
	IRRI		
	IRRI		
	BPI		
Inbred			

Inbred	IRRI		
Inbred	IRRI		
Inbred	IRRI		
Inbred	UPLB		
Inbred	IRRI		
Inbred	IRRI	8190 cav.	
Inbred	IRRI	3970	
Inbred	IRRI	2510	
Inbred	IRRI	3520	
Inbred	IRRI	3520	
Inbred	BPI	2500	
Inbred	UPLB	980	
Inbred	BPI	4300	
Inbred	IRRI	4400	
Inbred	UPLB	2600	
Inbred	IRRI	3100	
Inbred	IRRI	4000	
Inbred	IRRI	4500	
Inbred	UPLB	3040	
Inbred	BPI	4700	
Inbred	IRRI	4500	
Inbred	UPLB	4700	
Inbred	BPI	4600	
Inbred	IRRI	4700	
Inbred	IRRI	4100	
Inbred	IRRI	4770	
Inbred	IRRI	4770	
Inbred	IRRI	5600	
Inbred	BPI	4890	
Inbred	IRRI	5190	

Inbred	IRRI	4710	
Inbred	IRRI	5000	
Inbred	IRRI	4470	
Inbred	IRRI	4810	
Inbred	IRRI	4600	
Inbred	IRRI	2390	
Inbred	IRRI	4500-5600	
Inbred	IRRI	4200-4800	
Inbred	IRRI	5100	
Inbred	IRRI	5300	
Inbred	IRRI	5070	4684
Inbred	UPLB	4200	
Inbred	UPLB	4200	
Inbred	IRRI	2700	
Inbred	IRRI	5015.5	4528.5
Inbred	IRRI	5015.33	4552.5
Inbred	UPLB	4343	4495.5
Inbred	PHILRICE		
Hybrid	IRRI	5741	4932
Inbred	IRRI	4347	4308
Inbred	IRRI	4822	4229
Inbred	UPLB	4696	3985

Inbred	PhilRice	4771	4915
Inbred	PhilRice	2730.5	
Inbred	PhilRice	2671	
Inbred	PhilRice		2799
Inbred	IRRI		
Inbred	IRRI	4181	
Inbred	IRRI	4282	
Inbred	IRRI	2744	
Inbred	IRRI	2969	
Inbred	IRRI	5290	
Inbred	IRRI	4970	
Inbred	PhilRice	5270	
Inbred	UPLB	4910	
Inbred	IRRI		3630
Inbred	PhilRice		3700
Inbred	UPLB	4960	
Inbred	PhilRice	5210	

Inbred	IRRI		3350
Inbred	IRRI		3220
Inbred		2850	
Inbred		2900	
Hybrid	IRRI	5420	
Inbred	UPLB	5352	4968
Hybrid	Cargil Phil. Inc.	5049	4632
Inbred	PhilRice	5016.5	4463
Inbred	IRRI	5035.5	4587.5
Inbred	IRRI	5078	5094.5
Inbred	IRRI	910	2671
Inbred	IRRI	976.5	2148
Inbred	IRRI	488	1448
Inbred	PhilRice	3635	3166
Inbred	IRRI		3631
Inbred	IRRI		3321

Inbred	IRRI		3675
Inbred	UPLB		4569.5
Inbred	PhilRice		4467.5
Inbred	PhilRice		
Inbred	IRRI	5794	3704
Inbred	IRRI		4303
Inbred	PhilRice	4668	3868
Inbred	IRRI	3586	2315
Inbred	PhilRice	3315	2584
Inbred	IRRI		
Inbred	PhilRice		
Inbred	IRRI	4772	4411
Inbred	IRRI	4624	4427
Hybrid	IRRI	5849	5734
Hybrid	IRRI	5847	5715
Inbred	IRRI	5050.5	4656

Inbred	PhilRice	4980	5033
Inbred	IRRI	5063	4953.5
Hybrid	Bayer Crop Science	5673	5669
Hybrid	Monsanto Philippines Inc.	6308	6139
Inbred	PhilRice	4822	4692
Inbred	PhilRice	5069	4817
Hybrid	SL Agritech	5976	5847
Inbred	UPLB	4764	4480
Inbred	PhilRIce	6665	4910
Inbred	PhilRIce	4237	4410
Inbred	PhilRice	5190	5630
Inbred	PhilRice	5191	3900
Hybrid	IRRI	6502	6827
Inbred	PhilRice	5337	5879
Inbred	PhilRice	5433	6141
Inbred	PhilRice	5403	5784

Inbred	UPLB	5124	5482
Inbred	PhilRice	5300	4182
Inbred	IRRI	4592	4610
Inbred	PhilRice	5879	5883
Inbred	PhilRice	5706	6255
Inbred	PhilRice	5670	6125
Inbred	UPLB	5521	5841
Inbred	IRRI	5703	6245
Inbred	PhilRice		
Hybrid	BIOSEED Research Phils., Inc.	6271	5703
Hybrid	HyRice Seed Corporation	6454	6272
Hybrid	Syngenta Phils., Inc.	6398	6831
Hybrid	Bayer Crop Science	6368	6909
Inbred	IRRI	4820	4269
Inbred	IRRI	5449	4192

Hybrid	PhilSCat	6723	7403
Hybrid	PhilSCat	6366	6719
Hybrid	PhilSCat	5837	6825
Hybrid	BIOSEED Research Phils., Inc.	6987	5848
Inbred	IRRI	2951	1861
Inbred	PhilRice	3554	1684
Inbred	PhilRice	3380	2927
Inbred	PhilRice	3295	3184
Inbred	PhilRice	3332	2405
Inbred	IRRI		
Inbred	IRRI	3500	2500
Hybrid	PhilRice	5984	6381
Hybrid	PhilRice	6354	6750
Hybrid	Bayer Crop Science	6225	6791

Hybrid	PhilRice-UPLB	6858	6465
Hybrid	PhilRice-UPLB	6521	6372
Hybrid	IRRI, Laguna	6231	6541
Hybrid	Syngenta Phil., Inc.	5935	5919
Hybrid	Pioneer Hi-bred	6857	5862
Inbred	IRRI	6004	6075
Inbred	IRRI	6070	5896
Inbred	PhilRice	5835	6108
Inbred	PhilRice	4905	3284
Inbred	IRRI	3421	2446
Inbred	IRRI	5965	6271
Inbred	PhilRice	5578	5851
Inbred	PhilRice	6054	6303
Hybrid	HyRice Corporation	7116	7417

Hybrid	IRRI	7202	6151
Hybrid	IRRI	7101	6926
Hybrid	SYNGENTA	6869	6115
Hybrid	SEEDWORKS	6201	6211
Inbred	IRRI	6955	5843
Inbred	PhilRice	6971	5768
Inbred	PhilRice	5171	3911
Inbred	IRRI	5298	2187
Hybrid	PhilRice	7072	6997

Hybrid	IRRI	7414	6887
Hybrid	IRRI	7086	7162
Hybrid	IRRI	6950	6049
Hybrid	Advanta International Phils.	7266	6424
Hybrid	Advanta International Phils.	5982	6144
Hybrid	Beidahuang Seed Group Co. Ltd.	6495	6154
Hybrid	Dhaanya Seeds and RVIG-PhiLRice	6958	6453

Hybrid	Seedworks Phils., Inc.	6593	6396
Hybrid	PhilRice-CLSU-PHILSCAT	6972	5919
Hybrid	Dhaanya Seeds	6697	5986
Hybrid	DEVGEN NV	6580	6065
Hybrid	DEVGEN NV	7008	5944
Hybrid	DEVGEN NV	6677	6268
Inbred	PhilRice		

Inbred	IRRI		
Inbred	UPLB		
Inbred	IRRI		
Inbred	IRRI		
Inbred	UPLB		
Inbred	IRRI		
Inbred	UPLB		

Inbred	PhilRice		
Inbred	IRRI		
Inbred	PhilRice	4159	2885
Inbred	PhilRice	3135	2912
Inbred	PhilRice	3140	2765
Inbred	IRRI	3300	3159
Inbred	PhilRice		
Inbred	PhilRice	5881	5602
Inbred	IRRI	5815	5687
Inbred	PhilRice	3521	2831

Hybrid	Methahelix Life Science	5643.75	6141
Inbred	PhilRice	6728	5480
Hybrid	Bayer Crop Science	5924	5541
Hybrid	Bayer Crop Science	5973	5308
Hybrid	Bayer Crop Science	6127	5637
Hybrid	PhilRice	6176	5754
Hybrid	PhiRice	6702	5801

Hybrid	PhilRice	6083	5465
Hybrid	Pioneer Hi-bred	6090	5552
Inbred	PhiRice	2762	2134
Inbred	IRRI	2636	2218
Inbred	IRRI	2777	2043
Inbred	PhilRice	3128	2208
Inbred	PhilRice	2508	2166
Inbred	IRRI	2765	2135
Inbred	IRRI	3675	2523
Inbred	PhilRice	2625	2345
Inbred	IRRI	2797	2265
Inbred	PhilRice	4755	3906

Inbred	IRRI	6022	4524
Inbred	PhilRice		
Inbred	PhilRice		
Hybrid	Seedworks	6426	5905
Inbred	IRRI		
Inbred	IRRI		
Inbred	IRRI		
Inbred	IRRI	5839	4546
Inbred	PhilRice	5813	5497

Inbred	IRRI	5172	4978
Inbred	PhilRice	5381	5213
Inbred	IRRI	5256	4952
Hybrid	Syngenta Phils., Inc.	6289	5921
Hybrid	Syngenta Phils., Inc.	6292	5634
Hybrid	Syngenta Phils., Inc.	6330	5703
Hybrid	PhilRice	6183	5592
Hybrid	IRRI	6309	5875

Hybrid	BIOSEED Research Phils., Inc.	5567	4953
Hybrid	Syngenta Phils., Inc.	6414	5632
Hybrid	Syngenta Phils., Inc.	6730	5883
Hybrid	Advanta India Limited	6141	5303
Hybrid	IRRI	6181	6098
Hybrid	Syngenta Phils., Inc.	6605	5776
Hybrid	Long Ping High-Tech	6271	5760

Hybrid	Syngenta Phils., Inc.	6541	6173
Hybrid	Syngenta Phils., Inc.	6404	5512
Inbred	IRRI	3730	4382
Inbred	IRRI	3366	3103
Inbred	PhilRice	5756	5310
Inbred	PhilRice	5367	4766
Inbred	UPLB	5773	5794
Inbred	IRRI	5628	6277

Inbred	PhilRice	5804	5275
Hybrid	Long Ping High-Tech	5969	5853
Hybrid	Bayer Crop Science	6081	5654
Hybrid	IRRI	5974	6238
Hybrid	Bayer Crop Science	5979	5695
Hybrid	Long Ping High-Tech	5630	5896
Inbred	PhilRice	4600	2914
Inbred	PhilRice		

Inbred	UPLB		
Inbred	IRRI		
Inbred	PhilRice		
Inbred	PhilRice		
Inbred	PhilRice		
Inbred	IRRI		
Inbred	PhilRice		
Hybrid	IRRI	6429	5819
Inbred	IRRI		

Inbred	IRRI	6183	5600
Inbred	PhilRice	6548	5344
Inbred	PhilRice	6868	5571
Inbred	IRRI	6987	5668
Hybrid	Syngenta Phils.Inc.	7311	6089
Hybrid	UPLB	6927	6330
Hybrid	Syngenta Phils.Inc.	7280	5959

Hybrid	Syngenta Phils.Inc.	7248	5759
Hybrid	Syngenta Phils.Inc.	7018	5840
Hybrid	IRRI	6945	5867
Hybrid	PhilSCAT	7104	6230
Hybrid	Long Ping High Tech	6677	6575
Inbred	IRRI	5579	4246
Inbred	PhilRice	3782	2757
Inbred	IRRI	2988	3022
Inbred	IRRI	3481	2964

Inbred	IRRI	4103	3145
Inbred	PhilRice	4330	2835
Inbred	PhilRice		3264
Inbred	IRRI		3225
Inbred	UPLB		3588
Inbred	IRRI		3765
Inbred	IRRI		3224
Inbred	PhilRice	3943	2797
Inbred	IRRI	4636	3286

Inbred	PhilRice	5128	4158
Hybird	Long Ping High Tech	7132	5885
Hybrid	SL-Agritech Corp.	6775	5717
Hybrid	PhilSCAT	7113	5421
Hybrid	Pioneer Hi-Bred	6599	5924
Hybrid	Bayer Crop Science	6496	5460
Hybrid	Syngenta Phils., Inc.	7446	5780

Hybrid	Syngenta Phils., Inc.	7129	5959
Hybrid	Syngenta Phils., Inc.	7316	6137
Hybrid	Bayer Crop Science	6945	5583
Hybrid	IRRI	6649	5979
Inbred	UPLB RVIT	5638	6027
Inbred	IRRI	5612	6343
Inbred	UPLB RVIT	5472	6229
Inbred	IRRI	5538	5475

Inbred	IRRI	5561	6588
Hybrid	Long Ping High Tech	6863	6015
Hybrid	Syngenta Phils., Inc.	6766	6000
Hybrid	Bayer Crop Science	6050	5798
Hybrid	Bioseed Research Phils., Inc.	6237	6013
Hybrid	Bayer Crop Science	7043	5630
Hybrid	PhilSCAT	6157	5954
Inbred	PhilRice	3427	2236

Inbred	IRRI	3818	1934
Inbred	IRRI	3392	2333
Inbred	IRRI	3874	2121
Inbred	PhilRice	2994	2334
Hybrid	PhilSCAT/Long Ping	6692	5979
Hybrid	SL-Agritech Corp.	6310	6494
Hybrid	PRASAD SEEDS	5817	5492

Hybrid	PhilRice/UPLB	6652	6665
Hybrid	Bayer Crop Science	6575	5700
Hybrid	Bayer Crop Science	5975	5641
Hybrid	Long Ping High Tech	6229	5779
Hybrid	PhilRice/UPLB	6736	6382
Inbred	IRRI	2910	2277
Inbred	IRRI	2730	2524

Inbred	PhilRice	2729	2043
Inbred	PhilRice	3986	
Inbred	PhilRice	4280	
Inbred	PhilRice	3833	
Inbred	PhilRice		2505
Inbred	PhilRice		

Inbred	IRRI		
Inbred	PhilRice		
Inbred	PhilRice		
Inbred	IRRI		
Inbred	PhilRice		

Inbred	IRRI	4609	5062
Inbred	UPLB	4627	5243
Inbred	IRRI	4069	2586
Hybrid	Syngenta Phils. Inc.	7293	6223

Hybrid	Pioneer Hi-Bred	6853	6494
Inbred	IRRI		3910
Inbred	PhilRice		
Inbred	PhilRice		
Inbred	UPLB		
Inbred	IRRI		
Inbred	Philrice	5734	

Inbred	Philrice	5621	
Inbred	Philrice	2498	3559
Inbred	IRRI	2737	3161
Inbred	IRRI	2912	3107
Inbred	IRRI	2387	3205

Inbred	IRRI	3261	3337
Hybrid	Long Ping High Tech	2998-10165	
Hybrid	Long Ping High Tech	4533-13067	
Hybrid	Long Ping High Tech	4718-9636	
Hybrid	Green and Grow Teknologies, Inc	4086-9549	

Inbred	Philippine Rice Research Institute		
Inbred	Philippine Rice Research Institute		
Inbred	Philippine Rice Research Institute		
Inbred	International Rice Research Institute		
Inbred	International Rice Research Institute		
Inbred	International Rice Research Institute		
Inbred	International Rice Research Institute	2079-7277 kg/ha	
Inbred	International Rice Research Institute		
Inbred	Philippine Rice Research Institute	6070	5458
Inbred	Philippine Rice Research Institute	5063	5197
Inbred	Philippine Rice Research Institute	6199	3616

Inbred	Philippine Rice Research Institute	6198	5425
Inbred	Philippine Rice Research Institute	5047	4668
Inbred	International Rice Research Institute	4617	3510
Inbred	International Rice Research Institute		
Hybrid	Long Ping High Tech	Luzon: 4617-10674 kg/ha; Visayas: 1800-7903 kg/ha	
Hybrid	Long Ping High Tech	5117-10778 kg/ha	
Hybrid	Long Ping High Tech	4400-10859 kg/ha	
Hybrid	Long Ping High Tech	Luzon: 4817-10746 kg/ha Visayas: 2182-7633 kg/ha	
Hybrid	Long Ping High Tech	3994-9922 kg/ha	

Hybrid	Long Ping High Tech	4774-10844 kg/ha	
Hybrid	Corteva Agriscience Philippines, Inc	3867-11241 kg/ha	
Hybrid	Leads Agriventures Corporation	1945-8507 kg/ha	
Hybrid	Advanta Seeds Philippines, Inc.	6395	5561
Inbred	Philippine Rice Research Institute		
Inbred	Philippine Rice Research Institute		
Inbred	International Rice Research Institute		
Inbred	International Rice Research Institute		
Inbred	International Rice Research Institute		
Inbred	UPLB		
Inbred GM/Biotech	Philippine Rice Research Institute - International Rice Research Institute	National: 5330-5813 kg/ha	

Inbred	International Rice Research Institute	6062	4736
Inbred	Philippine Rice Research Institute	2805	3333
Hybrid	Bayer Crop Science	Central Luzon: 5918-8573 kg/ha; Western Mindanao: 7472-9853 kg/ha	
Hybrid	International Rice Research Institute	Yield Range: 5602-8465 with average yield of 7300 at Central Luzon	
Hybrid	Leads Agriventures Corporation	North West Luzon: 3000-5400 kg/ha; Central Luzon: 5628-8377 kg/ha; Eastern Mindanao: 3453-7908 kg/ha	
Hybrid	Leads Agriventures Corporation	North East Luzon: 4959-9621 kg/ha; Central Luzon: 5778-9313 kg/ha; Southern Luzon: 5865-11241 kg/ha; Western Visayas and Southern Luzon: 4418-8180 kg/ha; Western Mindanao: 5801-	
Hybrid	Long Ping High Tech	North West Luzon: 2833-5548 kg/ha; Central Luzon: 4726-9121 kg/ha; Southern Luzon: 6159-7441 kg/ha	
Hybrid	Long Ping High Tech	North East Luzon: 6023-10771 kg/ha; Central Luzon: 6011-9493 kg/ha; Southern Luzon: 5830-7307 kg/ha; Western Mindanao: 5233-8588 kg/ha	
Hybrid	Philippine-Sino Center for Agricultural Technology	North West Luzon: 3333-6080 kg/ha; Western Visayas and Southern Luzon: 4551-7013 kg/ha	
Hybrid	Philippine-Sino Center for Agricultural Technology	Central Luzon: 5050-8591 kg/ha; Western Mindanao: 7707-7938 kg/ha	

Hybrid	Philippine-Sino Center for Agricultural Technology	Central Luzon: 5452-8204 kg/ha
Hybrid	Philippine-Sino Center for Agricultural Technology	Central Luzon: 5401-8859 kg/ha
Hybrid	Bayer Crop Science	Central Luzon: 4951-8244 kg/ha; Western Mindanao: 8556-10461 kg/ha
Hybrid	Corteva Agriscience Philippines, Inc	Western Mindanao: 8575-10917 kg/ha
Hybrid	Corteva Agriscience Philippines, Inc	North East Luzon: 6055-10950 kg/ha; Central Luzon: 4343-9691 kg/ha
Hybrid	Philippine Rice Research Institute	Yield Range: 5187-9312 with average yield of 6911 at Central Luzon
Hybrid	SL-Agritech Corp.	North East Luzon: 5365-10528; North West Luzon: 3667-5132; Central Luzon: 5792-9406; Southern Luzon: 5716-7151; Western Visayas and Southern Luzon: 3190-6503; Eastern Mindanao: 4808-7383
Hybrid	SL-Agritech Corp.	Southern Luzon: 5481-8520; Western Mindanao: 4467-8549
Hybrid	SL-Agritech Corp.	Southern Luzon: 39358428
Hybrid	SL-Agritech Corp.	Central Luzon: 4860-8777; North West Luzon: 4333-5440
Hybrid	SL-Agritech Corp.	Central Luzon: 4023-9407

Hybrid	SL-Agritech Corp.	Central Luzon: 4738 - 9485	
Special Purpose	Philippine Rice Research Institute	7120	6422
Inbred	Philippine Rice Research Institute		National: 2615-6979
Inbred	University of the Philippines Los Banos		National: 2966-7744
Inbred	International Rice Research Institute		National: 2417-8348
Inbred	Philippine Rice Research Institute	5994	5874
Inbred	Philippine Rice Research Institute	6065	5763
Hybrid	Bioseed Research Phils., Inc.	8275	
Hybrid	Bioseed Research Phils., Inc.	6758	
Hybrid	Ramgo International Corporation	8683	
Hybrid	Bayer Crop Science	Zamboanga Peninsula: 9800 ; North South Eastern and North Central Mindanao: 8129	

Hybrid	Longping High-Tech Incorporated	Zamboanga Peninsula: 9213; North East Luzon: 8725; North West Luzon: 7633; North South Eastern and North Central Mindanao: 7991	
Hybrid	Longping High-Tech Incorporated	Zamboanga Peninsula: 9223; North West Luzon: 7447; North South Eastern and North Central Mindanao: 7977	
Hybrid	Corteva Agriscience Philippines, Inc	Zamboanga Peninsula: 10661; North East Luzon: 8792; North South Eastern and North Central Mindanao: 8110	
Hybrid	SL Agritech Corporation	Zamboanga Peninsula: 8869; North South Eastern and North Central Mindanao: 8145	
Hybrid	SL Agritech Corporation	North South Eastern and North Central Mindanao: 7990 and North West Luzon: 7998	
Inbred	Philippine Rice Research Institute - International Rice Research Institute	6633	5582
Inbred	Philippine Rice Research Institute - International Rice Research Institute	6990	5736
Inbred	Philippine Rice Research Institute - International Rice Research Institute	6687	5592
Inbred	UPLB		
Inbred	Philippine Rice Research Institute	4044	4046
Hybrid	International Rice Research Institute	North West Luzon: 6363 North South Eastern and North Central Mindanao: 8072	

Hybrid	Longping High-Tech Incorporated	North West Luzon: 6876; Southern Luzon: 7630; Zamboanga Peninsula: 9293; North South Eastern and North Central Mindanao: 8209
Hybrid	Longping High-Tech Incorporated	North East Luzon: 7775; North West Luzon: 6740; Central Luzon: 6312; Southern Luzon: 7494; Central Mindanao: 7491; Zamboanga Peninsula: 8885; North South Eastern and North Central Mindanao: 8708
Hybrid	SL Agritech Corporation	North East Luzon: 7653; North West Luzon: 6257; Central Mindanao: 7796; Zamboanga Peninsula: 8883; North South Eastern and North Central Mindanao: 8547
Hybrid	SL Agritech Corporation	Central Mindanao: 7645; Zamboanga Peninsula: 9293; North South Eastern and North Central Mindanao: 8452
Hybrid	Ramgo International Corporation	Zamboanga Peninsula: 8610; North South Eastern and North Central Mindanao: 8171
Hybrid	Universal Harvester Inc.	Zamboanga Peninsula: 8668; North South Eastern and North Central Mindanao: 8124
Hybrid	Enviro Scope Synergy Inc.	North Eastern Luzon: 8037 ; North South Eastern and Central Mindanao: 8226

Inbred	International Rice Research Institute, Philippine Rice Research Institute, University of the Philippines-Los Baños	6386	5879
Hybrid	Philippine Dadi Agricultural Technology Corporation	North East Luzon (NEL): 10,391; Zamboanga Peninsula and Western Mindanao (ZPWM): 10,510	
Hybrid	Philippine Dadi Agricultural Technology Corporation	Southern Luzon (SL): 8,189, Zamboanga Peninsula and Western Mindanao (ZPWM): 9,951, North South Eastern and North Central Mindanao (NSENCM): 8,833	
Hybrid	Philippine Dadi Agricultural Technology Corporation	North East Luzon (NEL): 9878, Southern Luzon (SL): 8363, Central Mindanao (CM): 7188, Zamboanga Peninsula and Western Mindanao (ZPWM): 9993, North South Eastern and North Central Mindanao (NSENCM): 8879	
Hybrid	Green and Grow Teknologies, Inc	Zamboanga Peninsula and Western Mindanao (ZPWM): 9312	
Hybrid	Bayer Crop Science	North West Luzon (NWL): 8202, Central Luzon (CL): 8255, Southern Luzon (SL): 8648, Zamboanga Peninsula and Western Mindanao (ZPWM): 10542, North South Eastern and North Central Mindanao (NSENCM): 9560	

Hybrid	Bayer Crop Science	North East Luzon (NEL): 9820, Central Luzon (CL): 8182, Central Mindanao (CM): 7959, Zamboanga Peninsula and Western Mindanao (ZPWM): 10653, North South Eastern and North Central Mindanao (NSENCM): 8827
Hybrid	Bayer Crop Science	North East Luzon (NEL): 9733, Central Luzon (CL): 7960, Southern Luzon (SL): 8235, Central Mindanao (CM): 7808
Hybrid	International Rice Research Institute	Central Luzon (CL): 7968, Central Mindanao (CM): 8436, Zamboanga Peninsula and Western Mindanao (ZPWM): 9782, North South Eastern and North Central Mindanao (NSENCM): 9065
Hybrid	Longping High-Tech Philippine R&D Center, Inc.	North West Luzon (NWL): 8505, Zamboanga Peninsula and Western Mindanao (ZPWM): 10192, North South Eastern and North Central Mindanao (NSENCM): 9274, Southern Luzon (SL): 8733
Hybrid	Longping High-Tech Philippine R&D Center, Inc.	North East Luzon (NEL): 10402, North West Luzon (NWL): 8535, Central Luzon (CL): 8288, Southern Luzon (SL): 8670, Zamboanga Peninsula and Western Mindanao (ZPWM): 10394, North South Eastern and North Central Mindanao (NSENCM): 9527

Hybrid	Longping High-Tech Philippine R&D Center, Inc.	North East Luzon (NEL): 10413, North West Luzon (NWL): 8278, Central Luzon (CL): 8052, Southern Luzon (SL): 8089, Central Mindanao (CM): 7848, Zamboanga Peninsula and Western Mindanao (ZPWM): 10656, North South Eastern and North Central Mindanao (NSENCM): 9063
Hybrid	Mahyco International Philippines Corporation	Central Luzon (CL): 8240, Southern Luzon (SL): 8459, Central Mindanao (CM): 8027, Zamboanga Peninsula and Western Mindanao (ZPWM): 10057
Hybrid	Mahyco International Philippines Corporation	North East Luzon (NEL): 9540, North West Luzon (NWL): 8249, Central Luzon (CL): 7947, Central Mindanao (CM): 7454, Zamboanga Peninsula and Western Mindanao (ZPWM): 8850, North South Eastern and North Central Mindanao (NSENCM): 8929
Hybrid	Mahyco International Philippines Corporation	Southern Luzon (SL): 8199, Zamboanga Peninsula and Western Mindanao (ZPWM): 9376, North South Eastern and North Central Mindanao (NSENCM): 8450
Hybrid	Mahyco International Philippines Corporation	North West Luzon (NWL): 7978

Hybrid	Central Luzon State University - Philippine Sino Agricultural Technology	North South Eastern and North Central Mindanao (NSENCM): 9501
Hybrid	Central Luzon State University - Philippine Sino Agricultural Technology	North South Eastern and North Central Mindanao (NSENCM): 9186
Hybrid	Philippine Rice Research Institute and University of the Philippines - Los Baños	Central Mindanao (CM): 8191
Hybrid	RiceTec, Inc.	North East Luzon (NEL): 10046 , Central Mindanao (CM): 7181, Zamboanga Peninsula and Western Mindanao (ZPWM): 8704, North South Eastern and North Central Mindanao (NSENCM): 8800
Hybrid	RiceTec, Inc.	Southern Luzon (SL): 7888 , Zamboanga Peninsula and Western Mindanao (ZPWM): 8729

Hybrid	Universal Harvester Inc.	Zamboanga Peninsula and Western Mindanao (ZPWM):8796
Hybrid	Winall Agristar Corporation	North East Luzon (NEL): 9852, Central Mindanao (CM): 8033, Zamboanga Peninsula and Western Mindanao (ZPWM): 9431, North South Eastern and North Central Mindanao (NSENCM): 8781
Hybrid	Winall Agristar Corporation	Zamboanga Peninsula and Western Mindanao(ZPWM): 10127

REGISTERED RICE VARIE

Oryza sativa L.

g/ha)		Maturity (DAS) *Based on NCT, including trials conducted under adverse environments*				Maturity (DAS) *Based on Seed Production under Favorable Conditions*	
DWSR		TPR		DWSR			TP
DS	WS	DS	WS	DS	WS	*Addendum (Pursuant to NSIC Resolution No. 01, series of 2026*)	DS

		127					97
		120-125					
		110-115					85
		132	129				
		142					105
		115	120				95
			123				107
		121					100
		0					90
		129					77
		125					104
		125					109
		120					90
		133					98-107
		120					117
		119					96
		116					96
		104					82
		166					104
		121					90
		112	104				77
		110	108				76
		108					80-87
		86					107
		100					75
		110	120				100
		113	114				78
		113-115					103
		122					
		109					83

		131	128				88
		112	113				88
		121	179				92
		129	127				90
		126	124				90
		121					104
		124	121				93
		105	102				75
		112	113				84
		108	112				82
		106	104				77
		110	105				95
		110	111				94
		125					112
		120	125				93
		110	110				81
		129	128				92
	3101				117		
		112	110				83
		110	112				83
		118	117				86
		112	112				93

		127	120				100
		127					121
		127					119
			130				
	3224				114		
	144			98			
		135					98
		126					88
		118					90
		115	115				86
		113	113				91
		114	114				88
		122	126				93
		113					100
			117				
		123	124				96
		120	125				85

			116				
			114				
		123					106
		122					104
		124	120				96
6925	3355	113	115				91
		100	111				85
		112	110				86
		112	113				87
		106	112				87
		111					77
		113					82
		116					88
		121	127				75
			131				
			133				

			136				
			116				
			118				
	2995				121		
		120	113				109
			117				
		155	148				93
		119	115				74
		128	127				78
	2851				119		
	2635				125		
4686	4525	113	112				87
4851	5211	111	111				90
		106	105				94
		106	106				93
5401	4701	109.5	104	104.5	97		87

5334	4339	111	108	105	100		86
		122	120				97
		118	118				106
		109	110				113
		118	118				93
5164	4515	108	107	106	101		87
		115	111				102
		121	119				86
		126	119				85
		118	116				85
5527	5099	106	109	105	100		88
		125	125				98
		107	108				100
5454	5512	110	111	108	104		90
5541	5459	110	109	105	102		89
5415	5774	111	112	106	104		83

5446	5484	107	107	104	101		89
5220	4466	112	109	107	103		90
		110	112				85
6629	5593	108	109	102	102		90
5905	5620	108	110	101	102		92
		109	110				90
5605	5497	110	112	102	105		87
		112	113				90
5748	5496			107	107		
		110	112				104
		107	105				100
		108	113				100
		113	114				108
		112	112				87
		118	109				66

		113	111				107
		116	111				111
		106	101				96
		111	109				107
		114	110				87
		119	121				87
		117	113				79
		114	113				82
		119	121				83
	3673				106		
		112	125				97
		104	102				97
		106	104				97
		110	112				107

		108	111				101
		110	112				107
		105	110				102
		116	115				110
		111	115				110
5059	6050	114	115	110	111		105
4867	6175	116	116	108	112		104
4922	6408	111	112	104	104		94
		122	118				101
		108	110				88
5004	6346	112	115	106	106		98
		112	110				101
5220	5651	114	110	104	103		100
		107	110				97

		107	109				99
		108	111				105
		115	116				116
		113	114				113
		108	111				99
6694	4878	115	115	109	107		103
		120	124				84
		107	111				89
		107	112				99

		106	107				98
		107	110				106
		107	111				103
		107	110				100
		118	117				115
		110	108				102
		106	106				97

		108	111				100
		113	115				109
		117	119				111
		110	110				107
		115	115				110
		106	112				107
	3003				110		

	2977				116		
	2630				119		
	2421				110		
	2516				123		
	2878				120		
	3733				114		
	3494				115		

	3562				118		
	2993				108		
		113	113				80
		111	111				85
		117	117				83
		116	117				79
5076	4966			104	103		
5630	4704	117	112	105	104		91
5331	4780	118	111	107	104		94
		112	115				115

		113	110				108
6815	4685	111	111	106	103		94
		114	111				99
		118	112				100
		119	118				106
		110	111				95
		109	111				106

		110	107				97
		115	115				104
		113	113				87
		115	116				93
		112	114				95
		115	114				95
		113	113				88
		114	116				95
		115	116				92
		115	114				96
		113	113				92
		113	115				101

		116	120				99
	3315				105		
	3027				103		
		115	117				108
	3056				107		
	2664				107	108-112	
	2272				108		
		109	112				103
		110	113				96

		114	117				104
		113	115				98
		119	117				104
		116	114				116
		106	106				95
		106	108				95
		109	108				103
		109	110				99

		109	107				113
		109	109				96
		110	111				102
		110	109				94
		109	109				99
		109	107				112
		113	108				105

		108	108				98
		107	105				100
		108	114				100
		109	116				93
5997	5070			107	104		
5866	4960			107	105		
5623	5275			107	104		
5742	5304	121	118				102

6141	5148	116	112	109	105		96
		112	112				99
		114	111				101
		109	107				100
		111	114				105
		114	112				100
		121	116				116
	3484				116		

	3758				113		
	3682				108		
	3426				112		
	3606				110		
	3671				110		
	3474				107		
	3474				111		
		110	111				101
	3369				105		

6096	4994	107	107				88
6336	4985	105	107	100	103		89
6135	4941	108	109				89
6300	5148	112	113				96
		112	108				104
		114	112				102
		117	109				102

		112	108				106
		115	112				108
		111	111				102
		113	111				103
		111	112				100
		118	114				101
		108	114				94
		112	109				96
		113	118				106

		132	109				97
		129	112				98
			117				
			116				
			113				
			113				
			107			115-118	
		119	113				100
		122	119				87

		126	123				98
		113	113				101
		112	113				106
		115	113				107
		113	114				110
		112	113				103
		107	110				104

		107	110				102
		106	110				104
		115	112				104
		109	111				95
5398	5222	112	108				96
5791	5598	112	108				93
4648	5451	111	108				101
5466	5686	115	109	106	103		89

5495	5105	114	108				91
		112	109				98
		110	110				105
		113	110				104
		113	113				105
		109	113				107
		115	112				106
		133	134				87

		134	131				96
		131	135				90
		133	127			115-117	85
		133	132				85
		113	112				102
		110	111				105
		114	113				104

		117	114				105
		111	112				105
		113	117				113
		113	111				95
		110	111				101
		135	126				94
		136	125				88

		133	125				86
		143±7					95
		143±6					97
		143±6					92
			145±7				
	2763				114±3		

	2487				115±4	117	
	2772				109±3		
	2480				113±3		
	2516				113±4		
	2498				113±4		

		114	109				103
4866	4608	109	110				99
		113	111				94
		110	114				102

		114	116				105
			144			125	
	3154				105		
	3026				109	120	
	3015				113		
	3111				118		
		110					95

		109					96
		102	107				90
		103	113				91
		105	113				97
		103	110				97

		106	111				95
		114	110				101
		112	103				101
		113	116				101
		113	112				98

5938	5085			103	108		
6026	4814			105	101		
5457	4814			105	106		
5849	4873			104	111		
5759	5109			105	107		
5720	4870			105	107		
2666-7090 kg/ha		113	111				102
3319-7732 kg/ha				106	110		
		115	114				102
		120	117				103
		118	117				119

		119	118				102
		116	118				103
		124	120				118
	1399-9480 kg/ha.				109		
		110	113				109
		112	111				98
		109	113				107
		111	114				103
		112	111				109

		113	112				102
		111	112				119
		110	111				107
		107	111				103
7545	2915			104	105		
4862	4451			107	105		
5008	4411			107	106		
5793	4573			105	105		
Visayas: 1964-7338 kg/ha Mindanao: 2255-9065 kg/ha				105	106		
5872	4485			102	97		
		113	122				97

		114	114				98
		112	118				95
		115	114				117
		114	113				115
		115	112				115
		114	110				106
		112	111				110
		110	110				102
		103	103				102
		113	114				109

		112	113				112
		115	114				115
		116	116				113
		115	115				120
		113	113				125
		108	110				103
		114	111				113
		114	114				123
		103	104				104
		116	110				111
		114	115				119

		113	114				112
		115	114				111
			110				
			104				
			105			120-122	
		112	109				97
		112	109				97
		109	114				102
		114	118				100
		111	112				100
		112	110				105

		112	113				99
		110	113				101
		113	115				102
		110	114				98
		109	113				98
		113	112				106
		111	116				104
		110	112				106
	3339				129		
		120	112				83
		110	108				104

		109	107				103
		110	107				104
		111	113				117
		106	106				106
		108	108				103
		110	110				109
		111	104				109

		113 (100-122)	115 (104-125)				103 (76-120)
		113 (108-128)	112 (104-118)				116
		112 (106-126)	111 (104-116)				110 (90-122)
		110 (103-121)	109 (106-115)				107 (87-118)
		110 (102-123)	109 (102-113)				116 (97-131)
		113 (107-122)	115 (105-122)				118 (94-132)

		112 (109-123)	115 (109-119)				120 (98-139)
		110 (100-121)	111 (103-117)				109 (89-124)
		107 (100-115)	111 (103-116)				107 (81-124)
		114 (105-126)	111 (104-116)				111 (90-126)
		113 (107-126)	110 (104-116)				112 (93-126)

		111 (102-126)	110 (101-106)				111 (91-124)
		108 (94-118)	113 (105-119)				111 (85-128)
		110 (103-122)	107 (103-116)				112 (83-124)
		104 (97-116)	105 (98-113)				107 (83-134)
		93 (87-104)	96 (90-101)				98 (75-113)

		113 (104-122)	112 (106-116)				117 (105-126)
		115 (108-127)	115 (110-119)				118 (95-134)
		111 (104-119)	112 (106-118)				117 (91-133)
		106 (103-110)	108 (100-112)				109 (79-125)
		109 (100-120)	108 (103-113)				112 (87-126)

		104 (97-117)	107 (101-115)				106 (103-134)
		108 (100-115)	108 (100-116)				114 (93-129)
		111 (101-120)	122 (105-116)				114 (94-126)

TIES

Plant Height (cm)			Productive Tillers				Culm Strength	
PR	DWSR		TPR (no./hill)		DWSR (no./li.m)			
WS	DS	WS	DS	WS	DS	WS		Green Leafhopper

								MR
								R
								R
90								
87								
94								
								R
93								

90								
88								R
107								I
94								I
94								
								I
100								MR
85								MR
93								I
92								I
89								MR
91								I
94								I
108			17	14				I
90			17	15				I
100			15	13				I
		99				84		I
92			19	17				I
102			16	15				I
89			16	15				I
95			16	15				I

102			16	15				R
			11					I
			11					I
126				10				I
		113				75		S
16								
			13					
			19					I
			18					
								R
								I
								MS
116								
								I
95								

116								
116								
101								
97			15	14				R
114			14	13				I
102			15	14				MS
101			14	14				MS
106			15	15				MS
			19					MS
			16					MS
			17					
96			15	16				MS
94				15				S
80				16				S

80				16				S
98				13				I
111				12				S
		119				75		S
109			14	13				I
101				13				I
94			18	15				MS
80			14	15				MS
92			15	16				MS
		98				79		S
		104				75		S
93			14	14				S
97			14	13				I
102			13	14				MS
103			12	13				I
99	93.5	99	14.5	13.5	229.5	206.5		MS

94	87	93	14	15	243	218		MR
110			16	14				I
109			13	13				MS
128			13	12				I
106			15	13				MS
93	85	92	14	14	232	215		I
112			11	11				I
99			18	15				MS
108			16	13				MS
96			14	14				MS
94	85	90	14	15	247	225		MS
107			19	16				MS
116			13	14				MS
97	90	94	15	15	260	324		I
99	90	94	14	14	264	318		I
92	82	88	14	14	251	319		I

99	93	93	14	14	116	346		MS
107	91	103	16	14	124	96		MS
97			14	14				I
99	87	88	14	14				MS
107	91	99	15	15	304	386		I
99			14	14				I
96	84	91	14	14				MS
98			15	15				I
	92	100			285	342		I
116			14	13				MS
110			14	13				I
113			13	13				I
121			14	14				I
92			18	15				MS
84			19	15				MS

113			12	12				I
114			12	12				MS
104			13	13				MS
119			15	14				I
84			13	13				S
87			13	14				MS
88			15	15				MS
81			14	15				MS
89			14	14				I
		109				91	Strong under normal condition	S
93			20	18				MR
103			14	13				MR
104			14	14				I
117			13	12				MS

114			13	13				MR
117			14	12				MR
117			15	13				MR
119			13	12				S
119			12	13				MS
112	100	108	14	15	297	290		MR
107	100	105	14	15	297	282		MR
98	88	96	13	14				MR
108			15	14				MS
89			13	13				S
104	94	102	13	14	292	263		MR
106			14	13			Strong	MR
104	98	105	14	14	288	302	Strong	MR
109			14	14			Strong	I

107			14	14			Strong	MR
116			13	15			Strong	I
122			13	12				MS
117			13	13			Strong	I
108			15	14			Strong	MR
110	100	108	12	12	266	252	Strong	MS
97			17	17			Strong	I
99			17	12			Strong	MS
109			15	14				I

108			14	13			Strong	I
117			14	13			Strong	I
116			14	13			Strong	MR
109			16	14			Strong	I
121			14	14			Strong	I
106			13	13			Strong	MS
109			15	14			Strong	I

114			13	13			Strong	I
118			14	13			Strong	MR
118			13	13			Strong	MS
116			12	12			Strong	MS
117			12	13			Strong	MS
128			13	12			Strong	I
		88				94	Strong	MR

		92				69	Strong	MR
		100				81	Strong	MR
		122				86	Moderate	MR
		104				81	Strong	I
		115				68	Strong	I
		98				91		I
		94				97		I

		127				60	Moderately Strong	MS
		108				64	Strong	S
82			15	14			Strong	MR
86			14	15			Strong	MS
85			15	15			Strong	MS
86			15	16			Strong	MS
	88	98			261	284	Moderate	I
105	92	99	16	13	256	281	Moderate to Strong	MR
105	90	101	16	14	256	275		MR
113			16	11			Intermediate	S

112			13	13				MS
103	90	98	15	14	302	287	Strong	I
107			14	13			Strong	I
105			14	13			Strong	I
114			13	14			Strong	MS
103			15	14			Strong	MR
112			14	13			Strong	MR

102			15	13			Strong	MR
115			13	13			Strong	I
95			13	14			Strong	MR
95			14	13			Strong	MR
94			13	14			Strong	MR
91			15	14			Strong	MR
89			13	14			Strong	MR
95			14	14			Strong	I
90			13	14			Strong	MR
93			13	13			Strong	MR
94			13	13			Strong	MR
105			16	15			Strong	MR

106			18	17			Strong	I
		97				111	Strong	I
		119				74	Strong	I
117			14	12			Strong	MS
		84				62	Strong	I
		87				67	Strong	MS
		90				65	Strong	I
104			14	13			Strong	S
94			14	14			Strong	MR

106			14	14			Strong	MS
98			14	14			Strong	MS
102			14	14			Strong	I
122			13	13			Strong	MS
103			14	14			Strong	I
105			14	14			Strong	I
113			13	13			Strong	I
109			15	13			Strong	MS

116			10	11			Strong	I
105			14	14			Strong	I
108			14	14			Strong	MS
103			14	13			Strong	I
108			14	14			Strong	I
118			13	12			Strong	I
106			13	11			Strong	I

108			14	13			Strong	MS
103			14	13			Strong	I
98			11	15			Strong	MS
89			12	13			Strong	I
	93	96			521	229	Strong	I
	89	87			499	217	Strong	I
	96	100			525	225	Strong	I
107			14	15			Strong	I

103	95	94	14	15	566	223	Strong	I
105			12	12			Strong	I
104			13	13			Strong	I
108			13	13			Strong	I
111			13	13			Strong	I
107			12	12			Strong	I
107			15	12			Strong	I
		87				110	Strong	MR

		101				99	Strong	I
		104				107	Strong	I
		89				104	Strong	I
		94				85	Strong	I
		94				107	Moderately Strong	I
		97				84	Strong	I
		92				101	Strong	I
110			14	12			Strong	I
		94				108	Strong	I

95			14	13			Strong	I
98	89	93	13	13	327	296	Strong	I
95.5			15	14			Strong	I
106			14	14			Strong	I
112			14	12			Strong	I
111			13	13			Moderately Strong	I
110			13	13			Strong	I

108			14	13			Moderately Strong	I
113			12	12			Strong	I
110			13	12			Moderately Strong	I
109			13	12			Strong	I
111			13	14			Strong	I
106			16	13			Strong	I
96			14	14			Strong	S
89			13	13			Strong	S
96			12	13			Strong	I

93			14	11			Strong	I
95			14	13			Strong	I
86				88			Moderately Strong	I
89				92			Strong	S
93				93			Moderately Strong	S
97				100			Moderately Strong	S
96				94			Moderately Strong	I
99			15	12			Strong	S
88			16	14			Strong	S

113			15	12			Strong	I
105			13	12			Strong	I
115			12	11			Strong	S
111			13	13			Strong	I
115			13	12			Strong	S
114			14	13			Strong	I
110			14	13			Strong	I

108			15	13			Strong	I
110			14	14			Strong	S
113			13	13			Strong	I
106			14	12			Strong	I
106			13	13			Strong	I
105			14	14			Strong	I
113			13	13			Strong	I
97	88	98	12	13	315	234	Strong	I

98			13	13			Strong	I
100			14	13			Strong	S
115			14	13			Strong	I
108			13	13			Strong	S
117			12	12			Strong	S
114			14	15			Strong	I
109			13	13			Strong	S
91			15	14			Strong	I

96			15	13			Strong	I
89			13	13			Strong	I
84			13	12			Strong	I
89			12	13			Strong	I
107			13	12			Strong	I
110			12	11			Strong	I
107			13	13			Strong	S

116			14	12			Strong	I
115			13	12			Strong	S
114			15	13			Strong	S
101			12	12			Strong	I
109			13	13			Strong	I
96			13	13			Strong	I
91			13	13			Strong	I

94			14	14			Strong	I
			13					MS (Screenhouse)
			12					I (Screenhouse)
			13					MS (Screenhouse)
84				14				MS (Screenhouse)
		96				72	Strong	I

		93				81	Strong	I
		99				74	Strong	I
		94				76	Strong	I
		93				95	Strong	I
		88				75	Strong	I

111			14	13			Strong	I
109			13	14			Strong	I
88			13	13			Strong	S
112			13	12			Strong	I

113			14	14			Strong	I
121				11			Strong	I
		96				81	Intermediate	R
		93				74	Strong	I
		96				69	Intermediate	I
		90				63	Strong	I
			13				Strong	I

			13				Strong	I
100			14	12			Intermediate	I
104			15	13			Strong	I
117			15	14			Intermediate	I
109			14	15			Strong	I

102			15	14			Moderately Strong	I
107			14	13				I
108			14	12				I
114			14	13				I
106			13	13				I

	96	103			299	288		I
	97	108			257	266		I
	99	103			265	282		I
	101	106			314	292		I
	102	109			275	271		I
	94	100			287	271		I
110			13	13				I
	95	102			264	247		I
116			15	15				I
123			13	11				I
127			15	13				I

119			16	13				I
120			13	12				I
115			15	13				I
		101				73		I
116			15	12				S
113			13					I
118			14	12				I
113			15	13				I
115			14	13				I

116			13	12				S
124			14	13				I
110			13	12				S
110			16	13				S
	102	110			292	304		I
	96	103			224	318		I
	103	108			212	336		I
	97	103			301	319		I
	100	106			208	321		S
	99	97			300	152		I
142.8			10	16				S

112			13	14				S
104			16	13				S
123			13	13				S
124			13	13				S
120			12	12				S
116			13	13				S
117			13	12				I
115			13	13				S
112			13	13				S
117			12	13				I

120			13	12				I
123			12	12				I
125			11	13				S
127			12	12				I
136			13	13				S
115			13	13				S
120			11	11				I
131			12	12				I
111			13	14				S
117			11	11				I
131			11	12				I

123			11	12				S
122			16	15				I
109				102			Moderately Strong	I
107				95			Moderately Strong	I
113				109			Moderately Strong	I
106			14	15				I
105			14	14				I
118			13	13				S
114			14	14				S
113			11	12				S
117			11	11				S

114			13	13				S
117			11	12				S
119			12	14				S
113			11	12				S
111			11	11				I
119			17	17				I
118			14	16				I
121			16	17				I
		78				97		I
100			14	12				I
117			13	12				I

115			12	11				I
119			12	11				I
135			12	11				I
122			11	10				MS
111			12	11				I
120			11	11				MS
115			13	11				I

112 (96-134)			16 (9-27)	14 (10-21)				I
124 (111-169)			13 (8-17)	12 (9-17)				S
119 (107-131)			13 (9-16)	13 (9-18)				I
114 (101-126)			12 (8-15)	12 (9-17)				I
125 (109-145)			13 (8-15)	12 (8-18)				I
127 (114-140)			14 (9-16)	14 (10-23)				MS

130(114-147)			14 (9-18)	13 (8-18)				I
121 (104-156)			14 (9-19)	13 (9-18)				S
120 (109-128)			14 (9-21)	14 (9-19)				MS
121 (109-131)			14 (11-18)	13 (10-19)				S
119 (104-130)			13 (9-17)	13 (9-16)				MS

121 (104-135)			13 (8-17)	13 (9-21)				MS
124 (100-134)			14 (10-20)	14 (9-20)				MS
122 (102-131)			13 (8-17)	13 (10-17)				MS
114 (102-130)			14 (10-17)	14 (8-22)				MS
111 (96-154)			12 (8-19)	12 (8-17)				MS

120 (100-129)			12 (8-17)	12 (8-17)				I
125 (107-141)			13 (10-17)	13 (9-20)				MS
129 (121-143)			13 (9-16)	13 (8-18)				I
120 (103-135)			13 (8-19)	13 (9-17)				R
124 (110-134)			13 (8-16)	12 (9-17)				R

120 (108-134)			11 (6-14)	11 (7-15)				I
123 (108-134)			12 (8-17)	12 (8-19)				I
123 (114-140)			13 (8-15)	13 (9-20)				MS

REGISTERED RICE VARIETIES

Oryza sativa L.

Reaction to Pest				
Brown Planthopper	**Stem Borrer**	**Whorl Maggot**	**Rice Black Bug**	**Whiteheads**

R				

MR				
R	MR			
R				
R				
R				
R	MR			
R				
I	I	I		
I	I	I		
MR	R			
MR	R			
MS	R			
S	R	R		
R	I			
I	I			
MR	MR			
R				
R	MR			
MR	MR			
R	MR			
R	MR			
MR	I			
MR	I			

I	I			
R				
I	I			
I				
	I		MR	
MS	R			
MR	I			
MS	MS			
MS	I			
MS	I			
MR	MS			
I	MS			
	I			
MS	MS			
I	I			MS
I	I			I
I	I			MS
I	I			MS
I	I			MS
R	I			MS
I	I			MS
R	I			MS

R	R			MS
I	I			MS
S	I			I
I	S			MS
S	S			MS
I	R			
I	R			
R				
I	I			
MS	MS			
I	I			

	R			R
	R			R
MR	MR			MR
MS	MR			I
I	S			S
I	MR			MS
I	MR			S
MR				I
MR				MS
I				I
MS				R
I				I
I				

MR				S
R	I			I
I	I			R
MS	I			S
MR	I			MR
I	I			MR
I				S
MS				R
MS				MR
MS	I			MR
S	I			MR
MS	I			I
I	I			I
MS	I			MS
I	I			MS
MS				MS

I	MS			I
I	I			I
MS	I			I
MS	I			S
MS	MS			I
MS	MS			I
MS	I			MS
MS	I			I
S	I			I
MS	I			I
MS	I			I
MS	MS			I
MS	MS			MR
I	MS			MR
I	MS			MR
I	I			MR

MS	MS			R
S	I			MR
I	MS			WSB - I YSB - S
MS	MS			WSB - I YSB - R
MS	I			WSB - I YSB - MR
I	I			WSB - I YSB - MR
I	I			WSB - I YSB - R
MS	MS			WSB - R YSB -
MS	MS			WSB - MR YSB - R
MS	MS			WSB - MR YSB - MR
MS	MS			MR
MS	I			I
I	I			WSB - I YSB - R
S	MS			I
S	I			I

MS	MS			WSB - I YSB - I
MS	MS			WSB - MS YSB - MR
MS	I			WSB - MR YSB - I
I	I			WSB - I YSB - MR
MS				MR
MS				R
MS	I			MS
MS	I			I
MS	I			MS
MS	MS			MR
I	I			R
MS	I			R
MS	I			MS

I	I			I
MR	MS			I
I	MS			MS
MS	MS			I
I	MS			MS
MR	MS			R
MR	MS			R
MR	I			MR
I	MS			MS
MS	I			MR
MR	I			MR
MR	MS			WSB - MS YSB - I
MR	MS			WSB - MS YSB - I
I	MS			WSB - MR YSB - MS

I	MS			WSB - R YSB - I
MS	MS			WSB - R YSB - MS
I	MS			WSB - I YSB - I
MS	MS			WSB - MR YSB - I
I	I			PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MS UPLB (YSB) - R
I	MS			PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - I UPLB (YSB) - I
I	I			PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - R VSU (YSB) - MR
MS	MS			PhilRice Midsayap (WSB) - MR PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - S VSU (YSB) - MR
I	I			PhilRice Agusan (WSB) - MR PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - S PhilRice CES (YSB) - I UPLB (YSB) - R

I	MS			PhilRice Agusan (WSB) - MR PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - S PhilRice CES (YSB) - I UPLB (YSB) - MR
I	I			PhilRice Agusan (WSB) - I PhilRice Midsayap (WSB) - MR PhilRice CES (YSB) - I UPLB (YSB) - R
I	I			PhilRice Agusan (WSB) - MR PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - MS PhilRice CES (YSB) - I UPLB (YSB) - R
I	I			PhilRice Agusan (WSB) - MR PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - MS PhilRice CES (YSB) - MR
I	I			PhilRice Agusan (WSB) - I PhilRice Midsayap (WSB) - I PhilRice Isabela (YSB) - MS PhilRice CES (YSB) - MR UPLB (YSB) - R
MS	I			PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I UPLB (YSB) - I
I	I			PhilRice Agusan (WSB) - I PhilRice Midsayap (WSB) - I PhilRice Isabela (YSB) - MR PhilRice CES (YSB) - MR UPLB (YSB) - R

I	I			PhilRice Agusan (WSB) - MR PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - S PhilRice CES (YSB) - MR UPLB (YSB) - R
I	MS			PhilRice Agusan (WSB) - I PhilRice Midsayap (WSB) - I PhilRice Isabela (YSB) - MR PhilRice CES (YSB) - I UPLB (YSB) - R VSU (YSB) - MR
I	I			PhilRice Agusan (WSB) - I PhilRice Midsayap (WSB) - I PhilRice Isabela (YSB) - S PhilRice CES (YSB) - MR UPLB (YSB) - MR VSU (YSB) - MR
I	MS			PhilRice Agusan (WSB) - MR PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - MR PhilRice CES (YSB) - I VSU (YSB) - MR
I	I			PhilRice Agusan (WSB) - MS
I	MS			PhilRice Agusan (WSB) - I PhilRice Midsayap (WSB) - I PhilRice Isabela (YSB) - R PhilRice CES (YSB) - I UPLB (YSB) - I
I	MS			PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MR PhilRice Isabela (YSB) - R VSU (YSB) - MR CVIARC (YSB) - MR

MS	MS			VSU (YSB) - RPhilRice Midsayap (WSB) - MS PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MR PhilRice Isabela (YSB) - R CVIARC (YSB) - R
I	MS			PhilRice Midsayap (WSB) - MS PhilRice Agusan (WSB) - I PhilRice CES (YSB) - MR VSU (YSB) - MR CVIARC (YSB) - I
I	MS			PhilRice Midsayap (WSB) - MR PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R VSU (YSB) - MR CVIARC (YSB) - MR
I	MS			PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice Isabela (YSB) - MR VSU (YSB) - R CVIARC (YSB) - R
I	MS			PhilRice Midsayap (WSB) - MR PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I VSU (YSB) - MR CVIARC (YSB) - MS
S	I			PhilRice Midsayap (WSB) - MS PhilRice Agusan (WSB) - MR VSU (YSB) - R BIARC (YSB) - R CVIARC (YSB) - MR
I	I			PhilRice CES (YSB) - MR BIARC (YSB) - R CVIARC (YSB) - MR

MS	I			PhilRice Midsayap (WSB) - MR PhilRice Agusan (WSB) - MR PhilRice Isabela (YSB) - I VSU (YSB) - I WESVIARC (YSB) - R BIARC (YSB) - R
I	S			PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice Isabela (YSB) - R CVIARC (YSB) - MR
MR	MR			I
I	I			S
MS	I			I
MS	I			S
MR	MS			PhilRice CES (YSB) - MR
MR	I			PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I
I	MS			PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I
MS	MS			PhilRice Agusan (WSB) - MS PhilRice Isabela (YSB) - I VSU (YSB) - R

I	MR			PhilRice Agusan (WSB) - MS PhilRice CES (YSB) - MS UPLB (YSB) - R
MR	I			PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - I UPLB (YSB) - R
I	I			PhilRice Agusan (WSB) - MS PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - MS VSU (YSB) - R BIARC (YSB) - R
MR	MS			PhilRice Agusan (WSB) - MS PhilRice CES (YSB) - I PhilRice Isabela (YSB) - MS VSU (YSB) - MR WESVIARC (YSB) - I BIARC (YSB) - R
I	I			PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R VSU (YSB) - R WESVIARC (YSB) - I BIARC (YSB) - R
MR	MS			PhilRice Agusan (WSB) - S PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - I VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R
I	I			PhilRice Agusan (WSB) - S PhilRice CES (YSB) - MS UPLB (YSB) - MR WESVIARC (YSB) - MS BIARC (YSB) - R

MS	I			PhilRice Agusan (WSB) - R PhilRice CES (YSB) - S PhilRice Isabela (YSB) - S VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R
I	MS			PhilRice Agusan (WSB) - S PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - MS VSU (YSB) - MR WESVIARC (YSB) - MS BIARC (YSB) - MR
I	MS			MS
I	MS			S
I	MS			MS
I	MS			I
I	MS			MS
I	I			S
I	MS			S
I	MS			MS
I	MS			S
I	I			PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I VSU (YSB) - MR WESVIARC (YSB) - I

MR	I			PhilRice Agusan (WSB) - MS PhilRice CES (YSB) - MS VSU (YSB) - I WESVIARC (YSB) - MR
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - MR
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - MR
MS	MR			I
MS				PhilRice Agusan (WSB) - MR
I				PhilRice Agusan (WSB) - MR
MS				PhilRice Agusan (WSB) - MR
S				PhilRice Agusan (WSB) - MR UPLB (YSB) - MR WESVIARC (YSB) - MS
MS				PhilRice Agusan (WSB) - MR UPLB (YSB) - MR WESVIARC (YSB) - MS

I				PhilRice Agusan (WSB) - MS PhilRice CES (YSB) - R UPLB (YSB) - R PhilRice Isabela (YSB) - MR WESVIARC (YSB) - MR
I				PhilRice Agusan (WSB) - I PhilRice CES (YSB) - MS UPLB (YSB) - R PhilRice Isabela (YSB) - MR WESVIARC (YSB) - MR
MS				PhilRice Agusan (WSB) - I PhilRice CES (YSB) - R UPLB (YSB) - R PhilRice Isabela (YSB) - R
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - MR VSU - MR WESVIARC (YSB) - I BIARC (YSB) - R
I				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - I VSU (YSB) - MR WESVIARC (YSB) - MS BIARC (YSB) - R
I				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - S PhilRice Isabela (YSB) - MS WESVIARC (YSB) - I BIARC (YSB) - R
I				PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I BIARC (YSB) - R
MS				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - S PhilRice Isabela (YSB) - I VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R

I				PhilRice Agusan (WSB) - S PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - MR WESVIARC (YSB) - MS BIARC (YSB) - R
I				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - I VSU (YSB) - R
MS				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - I VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R
I	MR			PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - MR WESVIARC (YSB) - MS BIARC (YSB) - R
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - S VSU (YSB) - R WESVIARC (YSB) - MR BIARC (YSB) - R
MS				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - MS VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - MS VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R

I				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - S PhilRice Isabela (YSB) - MS VSU (YSB) - MR WESVIARC (YSB) - S BIARC (YSB) - R
I				PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MS PhilRice Isabela (YSB) - I VSU (YSB) - R WESVIARC (YSB) - MS BIARC (YSB) - R
I				MS
I				MS
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - S PhilRice Isabela (YSB) - I
I				PhilRice Agusan (WSB) - R PhilRice Isabela (YSB) - I
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - S UPLB (YSB) - R PhilRice Isabela (YSB) - S VSU (YSB) - R
I				PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R

I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - S PhilRice Isabela (YSB) - I
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - I BIARC (YSB) - R
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - S PhilRice Isabela (YSB) - R WESVIARC (YSB) - I BIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - S PhilRice Isabela (YSB) - I VSU (YSB) - R WESVIARC (YSB) - I BIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
I				PhilRice Agusan (WSB) - I PhilRice CES (YSB) - S PhilRice Isabela (YSB) - R WESVIARC (YSB) - I
MR	I			PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - MR PhilRice CES (YSB) - MR PhilRice Isabela (YSB) - I VSU (YSB) - MR

I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R
I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I
I				PhilRice Midsayap (WSB) - S PhilRice Agusan (WSB) - I PhilRice CES (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - S
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I
I				R
I				R

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - R

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R
I				S
S				R
I				I

I				I
I				I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I
I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I
S				PhilRice Midsayap (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I WESVIARC (YSB) - R

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB)- R
I				PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - I BIARC (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - R BIARC (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R BIARC (YSB) - R
I				PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R WESVIARC (YSB) - I BIARC (YSB) - I

I				PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - I BIARC (YSB) - I
S				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R BIARC (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I WESVIARC (YSB) - R BIARC (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - I BIARC (YSB) - S
I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R BIARC (YSB) - I
S				PhilRice Midsayap (WSB) - S PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I
S				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R BIARC (YSB) - I
I				I

I				I
I				I
S				R
I				R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - I CBES (YSB) - I UPLB (YSB) - R
I				PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R CBES (YSB) - I UPLB (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I PhilRice Isabela (YSB) - I PhilRice Isabela (YSB) - I CBES (YSB) - I UPLB (YSB) - R

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - I CBES (YSB) - I UPLB (YSB) - R
S				PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R CBES (YSB) - I
S				PhilRice Midsayap (WSB) - S PhilRice CES (YSB) - R
I				PhilRice Midsayap (WSB) - S PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - I WESVIARC (YSB) - R CBES (YSB) - I UPLB (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - S PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I UPLB (YSB) - R
I				
I				

I				
S (Screenhouse)	Yellow SB - MS (Screenhouse)			S (Field)
I (Screenhouse)	Yellow stemborer - MS (Screenhouse)			S (Field)
MS (Screenhouse)	Yellow stemborer - MS (Screenhouse)			S (Field)
MS (Screenhouse)	Yellow stemborer - I (Screenhouse)			R (Field)
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I UPLB (YSB) - R WESVIARC (YSB) - R CBES (YSB) - I

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R UPLB (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R UPLB (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - I UPLB (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R UPLB (YSB) - R WESVIARC (YSB) - I CBES (YSB) - R
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R UPLB (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - S CBES (YSB) - I
I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I
I				PhilRice Midsayap (WSB) - I R PhilRice Isabela (YSB) - R WESVIARC (YSB) - I CBES (YSB) - I PhilRice CES (YSB) - I
I				PhilRice CES (YSB) - IPhilRice Isabela (YSB) - R WESVIARC (YSB) - I UPLB (YSB) - R

I				PhilRice Midsayap (WSB) - I PhilRice Agusan (WSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - S UPLB (YSB) - R
I				R
I				I
I				PhilRice(WSB) - I PhilRice (YSB)-R
I				UPLB(WSB) - I UPLB (YSB)-R
I				IRRI(WSB) - I IRRI (YSB)-R
I				R

I				R
I				R
I				I
I				I
I				R

I				I
I				I
I				R
I				WSB- I YSB-R
S				WSB - R YSB - I

I				I
I				WSB - I YSB - R
I				I
I				WSB - I YSB - R
I				R
I				WSB - I YSB - R
I				WSB - I YSB - R
I				I
I				I
I				I
I				WSB - I YSB - R

I				WSB - I YSB - R
I				WSB - I YSB - R
I				WSB - I YSB - R
I				I
I				WSB - I YSB - R
I				WSB - I YSB - R
I				I
I				I
I				WSB - S YSB - R

S				WSB - I YSB - R
I				R
S				WSB - I YSB - R
I				I
I				WSB- I YSB-R
I				WSB- I YSB-I
I				WSB- I YSB-R
I				WSB- I YSB-R
I				WSB- I YSB-R
I				WSB- R YSB-R
I				YSB-I

S				WSB- R YSB-R
S				YSB-R
S				R
S				R
S				R
S				R
I				I
S				I
S				I
S				I

S				R
S				R
I				R
S				I
S				R
S				R
S				I
I				I
S				I
I				I
I				R

I				R
I				WSB - I YSB - R
I				WSB - R YSB - R
I				WSB - R YSB - R
I				WSB - R YSB - R
I				WSB - R YSB- I
I				WSB - R YSB - R
S				WSB - I YSB - R
S				WSB - I YSB - I
S				WSB - I YSB - R
S				WSB - I YSB - I

S				WSB - I YSB - I
S				WSB - I YSB - I
S				WSB - I YSB - R
S				WSB - I YSB - R
I				WSB - I YSB - I
I				WSB - R YSB - I
S				WSB - I YSB - I
I				WSB - I YSB - I
I				WSB - R YSB - R
I				YSB - MS
I				WSB - MR YSB - MS

I				WSB - MR YSB - MS
I				WSB - I YSB - MS
I				WSB - I YSB - MS
I				WSB - I YSB - MS
I				WSB - MS YSB - MS
I				YSB - I
I				YSB - R

I				WSB - MR; YSB - MS
S				WSB - I; YSB - MR
MS				WSB - MS; YSB - R
I				WSB - M; YSB - MR
MS				WSB - MS; YSB - MR
S				WSB - I; YSB - MR

MS				WSB - I;YSB - MR
MS				WSB - I; YSB - MR
S				WSB - MR; YSB - MR
S				WSB - MR; YSB - MR
I				WSB - MR; YSB - MR

MS				WSB - I; YSB - MR
MS				WSB - MS; YSB - MR
MS				WSB - MS; YSB - MR
MS				WSB - I; YSB - I
MS				WSB - MS; YSB - I

MS				WSB - I; YSB - I
MS				WSB - MR; YSB - I
I				WSB - I; YSB - MR
MR				WSB - MS; YSB - I
I				WSB - I; YSB - MR

MS				WSB - MS; YSB - MR
MS				WSB - I; YSB - MR
I				WSB - I; YSB - MR

	Reaction to I		
Deadhearts	Bacterial Leaf Blast	Bacterial Leaf Blight	Sheath Blight

	R		
	R		
	R	R	
	R	R	
	R	R	
	R	R	
	MR	MR	R
	R	MR	
	R	R	R
	R		
	I	S	
	I	I	S
	I	R	S
	R	R	
	MR	MS	
	MR	MR	
	R	S	S
	I		S
	I	MS	
	MR	I	
	MR	MR	
	MR	R	
	MR	MR	
	S	I	I
		I	MS

		I	
	I	I	
	R	I	
	I	I	
	R	R	R
	R	I	
	I	I	
	R	I	
	I	I	I
	I	I	I
	R	I	I
	I	I	S
	I	I	I
	I	I	I
MS	I	I	I
I	I	I	I
S	I	I	I
S	I	I	I
I	R	I	I
MS	R	I	I
S	I	I	I
S	I	R	I

R	R	R	R
MS	I	I	I
MS	I	I	I
MS	R	I	I
MS	R	I	I
	I		
	S	S	R
	S		R
	R		
	R	R	
		R	
	MS	MS	MS
	R		
	R	I	

	R		
	R		
	R		
MR	S	I	I
R	R	I	I
MR	I	R	I
MR	I	I	I
R	R	I	I
	S	I	S
MS	S	I	S
I	I	S	S
	I	S	I
	Lowland conditon - S Cool Elevated - R	S	S
	Lowland conditon - S Cool Elevated - R	S	S

	Lowland conditon - S Cool Elevated - R	S	S
S	I	I	I
	I	I	I
S	R	I	I
	I	I	I
	I	I	I
	I	I	I
	I	S	I
	I	S	I
MR	I	I	I
R	I	I	I
I	I	I	I
I	I	I	I
MR	I	I	I
MR	I	I	I
MR	I	I	S

MR	I	I	S
R	R	I	I
	R	I	I
	R	I	I
R	S	I	I
MR	I	I	I
	R	I	I
	S	I	S
	I	I	I
	I	I	I
MR	I	I	I
MR	I	I	I
MR	I	I	S
MR	S	I	I
MR	I	S	I
MR	S	I	I

MR	I	S	I
	S	I	S
	I	I	I
	I	I	I
	I	I	S
	S	I	S
	S	S	S
WSB - MR YSB -	I	I	I
	I	I	S
WSB - MR YSB - MR	I	S	I
MR	I	S	S
MR	I	S	S
WSB - MR YSB - R	I	S	S
	R	I	S
	I	I	S

WSB - R	YSB - R	I	S	I
WSB - MR	YSB - R	I	I	S
WSB - MR	YSB - R	I	S	S
WSB - MR	YSB - R	I	S	I
		S	I	S
		R	I	S
		S	S	S
		I	S	S
		I	I	I
MR		I	S	S
			MR	
MR		I	S	S
		I	S	S
MR		I	S	S

	I	S	S
	I	S	I
	I	S	S
	I	I	I
	I	S	S
	I	I	S
	I	I	I
	S	I	S
	S	I	I
	S	I	S
	I	I	S
	I	S	S
	S	S	S
	I	S	S

	I	S	S
	I	S	S
	I	I	S
	I	I	I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - S PhilRice Isabela - R UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Isabela - R UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Isabela - I UPLB - S VSU - S	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Isabela - R UPLB - I VSU - I PhilRice Isabela - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R VSU - S	PhilRice CES - S PhilRice Midsayap - S UPLB - S	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - S VSU - S WESVIARC - I

PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - R VSU - S	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I VSU - S UPLB - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - S	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - S UPLB - R VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I

	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - R UPLB - R VSU - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - S VSU - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - S WESVIARC - I
PhilRice Midsayap (WSB) -	PhilRice CES - S	PhilRice CES - I	PhilRice CES - S
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - S WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I CVIARC - R WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I

	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I CVIARC - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - S VSU - S WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I CVIARC - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I CVIARC - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - S VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I
	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - S CVIARC - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I CVIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - R VSU - R BIARC - I CVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I BIARC - I CVIARC - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I BIARC - R CVIARC - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I BIARC - R CVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - S WESVIARC - I

PhilRice Midsayap (WSB) - MR	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - R BIARC - R CVIARC - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I BIARC - R CVIARC - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I
	I	I	S
	I	S	S
	S	I	S
	S	I	I
	PhilRice CES - S PhilRice Isabela - S UPLB - S VSU - I PhilRice Midsayap - S BIARC - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - I WESVIARC - S
	PhilRice CES - S PhilRice Isabela - R UPLB - I VSU - S PhilRice Midsayap - R BIARC - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I PhilRice Midsayap - R BIARC - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Isabela - R UPLB - R VSU - S PhilRice Midsayap - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I PhilRice Isabela - R

	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Isabela - I UPLB - S VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - I WESVIARC - S
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES -S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - S
	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - S	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - S
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - R
PhilRice Midsayap (WSB) - MR PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - S

	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - S WESVIARC - S PhilRice Isabela - S
	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - S	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - R
	S	S	S
	S	I	S
	S	I	S
	S	I	I
	S	I	S
	I	I	S
	S	I	S
	S	I	S
	S	S	S
	PhilRice CES - S PhilRice Isabela - I UPLB - S VSU - S PhilRice Midsayap - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - I WESVIARC - S PhilRice Isabela - I

	PhilRice CES - S PhilRice Isabela - I UPLB - S VSU - I PhilRice Midsayap - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I PhilRice Isabela - I
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I PhilRice Midsayap - I CVIARC - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - S PhilRice Isabela - R
	PhilRice CES - S PhilRice Isabela - R UPLB - I VSU - S PhilRice Midsayap - I CVIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - I WESVIARC - I PhilRice Isabela - R
	I	I	I
	PhilRice CES - I PhilRice Isabela - I UPLB - I VSU - I PhilRice Midsayap - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - R UPLB - I VSU - I WESVIARC - I PhilRice Isabela - I
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I PhilRice Midsayap - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - S PhilRice Midsayap - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - S WESVIARC - I PhilRice Isabela - S
	PhilRice CES - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - S VSU - S WESVIARC - I	PhilRice CES - S PhilRice Isabela -R UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - R UPLB - S VSU - S WESVIARC - I	PhilRice CES - S PhilRice Isabela - R UPLB - S VSU - S WESVIARC - I

	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I
	PhilRice CES - S PhilRice Isabela - I UPLB - S VSU - S WESVIARC - S	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - S WESVIARC - S	PhilRice CES - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Isabela - R UPLB - R VSU - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I PhilRice Isabela - R
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - I WESVIARC - I PhilRice Isabela - I
	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - I WESVIARC - I PhilRice Isabela - I
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES- S PhilRice Midsayap - R PhilRice Isabela - I UPLB - S VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - I WESVIARC - S PhilRice Isabela - I

	PhilRice CES- S PhilRice Midsayap - R PhilRice Isabela - R UPLB - S VSU - I	PhilRice CES- S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - S PhilRice Isabela - I
	PhilRice CES- S PhilRice Midsayap - R PhilRice Isabela - I UPLB - S VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - S WESVIARC - I PhilRice Isabela - R
	PhilRice CES- S PhilRice Midsayap - R PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I VSU - S WESVIARC - I PhilRice Isabela - I
	PhilRice CES- I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - S PhilRice Isabela - R
	PhilRice CES- I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I PhilRice Isabela - I
	PhilRice CES- I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES- S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - S
	PhilRice CES- I PhilRice Midsayap - S PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES- I PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - I PhilRice Isabela - I

	PhilRice CES- I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES- I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - S PhilRice Isabela - S
	PhilRice CES- I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - S WESVIARC - I PhilRice Isabela - I
	S	S	S
	S	S	S
	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Isabela - R UPLB - I VSU - S WESVIARC - S PhilRice Midsayap - I
	PhilRice CES - S PhilRice Isabela - S UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - R VSU - I WESVIARC - S	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S PhilRice Midsayap - I
	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S PhilRice Midsayap - S
	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I

	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - S PhilRice Midsayap - R PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Isabela - R UPLB - I VSU - R WESVIARC - I PhilRice Midsayap - I
	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - S UPLB - I VSU - I	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - R
	PhilRice CES - I PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I PhilRice Isabela - S
	PhilRice CES - S PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - I
	PhilRice CES - I PhilRice Isabela - I UPLB - S VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I PhilRice Isabela - R
	PhilRice CES - I PhilRice Isabela - I UPLB - R VSU - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Isabela - I UPLB - R VSU - I PhilRice Midsayap - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I PhilRice Midsayap - I
	PhilRice CES - S PhilRice Isabela - I UPLB - S VSU - I PhilRice Midsayap - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - I UPLB - S VSU - I WESVIARC - I PhilRice Midsayap - S

	PhilRice CES - I PhilRice Isabela - I UPLB - R VSU - I PhilRice Midsayap - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - S WESVIARC - I PhilRice Isabela - S
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - I PhilRice Midsayap - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - R
	PhilRice CES - S PhilRice Isabela - I UPLB - S VSU - S PhilRice Midsayap - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - S WESVIARC - I PhilRice Isabela - S
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - S PhilRice Midsayap - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S VSU - S WESVIARC - I PhilRice Isabela - I
	PhilRice CES - S PhilRice Isabela - S UPLB - S VSU - S PhilRice Midsayap - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - R
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - S PhilRice Midsayap - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - I WESVIARC - I PhilRice Isabela - R
	PhilRice CES - S PhilRice Isabela - I UPLB - I VSU - S PhilRice Midsayap - S	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - I PhilRice Midsayap - I UPLB - S VSU - S WESVIARC - I PhilRice Isabela - I
	I	S	I
	I	S	S

PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I BIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - S
PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I BIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - R WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I
PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I BIARC - R	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - S PhilRice Midsayap - R PhilRice Isabela - S UPLB - R BIARC - I	PhilRice CES - I PhilRice Midsayap - I UPLB - I VSU - R WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Isabela - S UPLB - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - R WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - S VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I

PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Isabela - I UPLB - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - S PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Isabela - I UPLB - R VSU - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - S UPLB - S VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I PhilRice Isabela - I
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - R BIARC - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - I
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - I BIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I
	I	I	S
	I	S	S
	S	I	S

	S	S	S
	I	I	S
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I BIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I
	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - S VSU - I BIARC - R	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S VSU - S WESVIARC - I PhilRice Isabela - R
	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - S UPLB - S BIARC - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - SWESVIARC - I
	PhilRice CES - I PhilRice Isabela - I UPLB - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - S WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - I BIARC - R	PhilRice CES - I PhilRice Midsayap - I UPLB - I VSU - S WESVIARC - I PhilRice Isabela - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I VSU - I WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S BIARC - I	PhilRice CES - I PhilRice Midsayap - I UPLB - I WESVIARC - S PhilRice Isabela - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I PhilRice Isabela - R

PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I BIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - S PhilRice Isabela - R	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I PhilRice Isabela - R
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - I PhilRice Isabela - I VSU - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I VSU - I
PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I BIARC - R	PhilRice CES - S PhilRice Midsayap - I UPLB - S WESVIARC - S PhilRice Isabela - I VSU - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I VSU - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - S PhilRice Midsayap - R PhilRice Isabela - I UPLB - R BIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S WESVIARC - S PhilRice Isabela - I VSU - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I VSU - S
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I PhilRice Isabela - I VSU - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I VSU - S
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - R BIARC - I	PhilRice CES - I PhilRice Midsayap - I UPLB - I PhilRice Isabela - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I VSU - S
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - I UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - S PhilRice Isabela - I	PhilRice CES - S PhilRice Midsayap - I UPLB - S WESVIARC - I VSU - I

PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I	PhilRice CES - R PhilRice Midsayap - R PhilRice Isabela - I UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - S PhilRice Isabela - I	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - I VSU - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I PhilRice Isabela - I VSU - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I VSU - S
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - R UPLB - R BIARC - R	PhilRice CES - I PhilRice Midsayap - I UPLB - I WESVIARC - S PhilRice Isabela - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - R
PhilRice CES (YSB) - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I BIARC - R	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - I PhilRice Isabela - I VSU - R	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - S VSU - I
PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I BIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - R
PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I BIARC - S	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - R UPLB - R WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - S PhilRice Isabela - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I BIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - I
PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I BIARC - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I PhilRice Isabela - I

PhilRice Midsayap (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I BIARC - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - R UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I	PhilRice CES - R PhilRice Midsayap - R PhilRice Isabela - R UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - R UPLB - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - S PhilRice Isabela - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I BIARC - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - S PhilRice Isabela - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - R BIARC - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - S WESVIARC - S PhilRice Isabela - I
PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I BIARC - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S VSU - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I VSU - I WESVIARC - I PhilRice Isabela - I
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - R UPLB - R WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - S
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - R BIARC - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - S
R	S	S	S

R	I	S	S
R	I	S	S
I	I	S	S
R	S	S	S
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R UPLB (YSB) - R	PhilRice CES - R PhilRice Midsayap - I PhilRice Isabela - I UPLB - R CBES - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I PhilRice Isabela - I
PhilRice Midsayap (WSB) - S PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R CBES (YSB) - R UPLB (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I CBES - R	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - S PhilRice Isabela - R
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R UPLB (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I CBES - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - S	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - S PhilRice Isabela - I

PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R UPLB (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I CBES - R	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - I UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - R
PhilRice Midsayap (WSB) - I PhilRice CES (YSB) - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - R CBES - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - R UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S UPLB - I WESVIARC - I PhilRice Isabela - R
PhilRice Midsayap (WSB) - S PhilRice CES (YSB) - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I CBES - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I UPLB - I WESVIARC - I PhilRice Isabela - R
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - I PhilRice Isabela (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R UPLB (YSB) - R	PhilRice Midsayap - I PhilRice CES - I PhilRice Isabela - I CBES - R UPLB - I	PhilRice Midsayap - I PhilRice CES - I PhilRice Isabela - I WESVIARC - I VSU - I UPLB - S	PhilRice Midsayap - S PhilRice CES - S PhilRice Isabela - S WESVIARC - I VSU - S UPLB - I
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R PhilRice CES (YSB) - R PhilRice Isabela (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R UPLB (YSB) - R	PhilRice Midsayap - I PhilRice CES - R PhilRice Isabela - I UPLB - I	PhilRice Midsayap - S PhilRice CES - I PhilRice Isabela - I WESVIARC - S VSU - I UPLB - S	PhilRice Midsayap - S PhilRice CES - S WESVIARC - I VSU - R UPLB - I
	PhilRice CES - I	PhilRice CES - S	PhilRice CES - S
	PhilRice CES - I	PhilRice CES - I	PhilRice CES - S

	PhilRice CES - S	PhilRice CES - S	PhilRice CES - S
	Induced - I	Induced - S	Induced - S
	Induced - I	Induced - S	Induced - S
	Induced - I	Induced - S	Induced - S
	Induced - I	Induced - S	Induced - S
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R UPLB (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - R UPLB - I CBES - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - S UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I

PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R UPLB (YSB) - R WESVIARC (YSB) - R CBES (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I CBES - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R UPLB (YSB) - R CBES (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - R UPLB - I CBES - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R WESVIARC (YSB) - R CBES (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - R UPLB - I CBES - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R UPLB (YSB) - R CBES (YSB) - R WESVIARC (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I CBES - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - I WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I WESVIARC - I
PhilRice Midsayap (WSB) - R PhilRice Agusan (WSB) - R UPLB (YSB) - R CBES (YSB) - R WESVIARC (YSB) - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - R UPLB - I CBES - R	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R WESVIARC - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - S WESVIARC - I

PhilRice Isabela (YSB) - R CBES (YSB) - R VSU (YSB) - R	PhilRice CES - R PhilRice Midsayap - I PhilRice Isabela - I UPLB - S	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I VSU - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - I UPLB - I WESVIARC - I VSU - I
PhilRice Isabela (YSB) - R CBES (YSB) - R	PhilRice CES - I PhilRice Midsayap - R PhilRice Isabela - R UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - S VSU - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I WESVIARC - I VSU - I
PhilRice CES (WSB) - I CBES (WSB) - R PhilRice Isabela (WSB) - R	PhilRice CES - R PhilRice Midsayap - I PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - R UPLB - I WESVIARC - I VSU - R	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I VSU - S
PhilRice CES (YSB) - R CBES (YSB) - R PhilRice Isabela (YSB) - R	PhilRice CES - R PhilRice Midsayap - R PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - I PhilRice Isabela - I UPLB - S WESVIARC - I	PhilRice CES - S PhilRice Midsayap - I PhilRice Isabela - I UPLB - I WESVIARC - I VSU - I

PhilRice Isabela (YSB) - R CBES (YSB) - R	PhilRice CES - R PhilRice Midsayap - R PhilRice Isabela - I UPLB - I	PhilRice CES - I PhilRice Midsayap - S PhilRice Isabela - R UPLB - S WESVIARC - I VSU - I	PhilRice CES - S PhilRice Midsayap - S PhilRice Isabela - S UPLB - I WESVIARC - I VSU - I
R	I	I	I
R	I	I	S
R	I	I	S
R	I	I	I
R	I	I	I
R	I	I	I

I	I	I	I
	R	I	I
R	R	I	I
R	I	I	S
R	I	I	S

R	R	I	S
R	R	I	I
R	R	I	I
R	R	I	I
I	I	I	I

	I	I	I
	R	I	I
	I	I	I
	R	I	I
	I	I	I
	I	I	I
	R	S	I
	R	I	I
R	I	I	I
R	I	I	I
R	R	I	I

R	R	I	I
R	I	I	I
R	I	I	I
I	R	I	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I

	R	I	I
	R	I	I
	R	S	I
	R	I	I
	I	I	I
	I	I	I
	R	I	I
	I	I	I
	I	I	I
	I	I	S
	R	I	S

	R	I	I
	R	I	S
R	R	I	I
R	R	S	I
R	R	I	I
R	R	I	I
R	R	I	S
R	R	S	I
R	R	S	S
R	R	I	I

R	R	I	I
R	R	I	I
R	R	I	I
R	R	I	I
R	R	I	S
R	R	I	S
R	R	S	I
R	R	S	I
R	R	S	S
R	R	I	S
R	R	I	I

R	R	S	I
	R	I	I
	I	I	I
	I	I	I
	R	S	I
	I	I	I
	I	I	I
	R	I	I
	R	S	I
	R	I	I
	R	I	I

	R	I	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I
WSB - I YSB - I		I	I
WSB - I YSB - I		I	I
YSB - 1	R	I	S
WSB - R	R	I	I
	S	I	S
	R	I	I

	R	I	S
	R	I	I
	R	I	S
	I	I	I
	R	I	S
WSB - R	R	I	S
WSB - R	R	I	S

	I	R	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I

	R	I	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I

	R	I	I
	R	I	I
	R	I	I
	R	I	I
	I	I	S

	R	I	I
	R	I	I
	R	I	I
	R	I	I
	R	I	I

	R	I	S
	R	I	I
	R	I	I

Diseases				Amylose Content %
Tungro	Helminthosporium leaf spot	Grassy Stunt	Ragged Stunt	

R		R		

MR		MR		
R		R		
R		R		
		R		
I				
I				
I				
I				
	MR			
	MR			
R	R	R		
R		R		
I		R	R	
I		I	I	
I		I	S	
			MR	
MR		R	R	
R		R	MR	
		R	R	
I		R	I	
I		R	MR	
R		R	R	
I		I	R	
MR		MR		
R				
R				
R		S	S	
R		MS	MS	

R		R		
R		I		
R		S		
I		I		
I				
		I	I	
		R	R	
		I	I	
I				
S				
I				
S				
S				
S				
Modified (IRRI) - I Induced - S				
I				
Modified (PhilRice) - I Induced - S				
Modified (IRRI) - I Induced - S				
Modified (IRRI) - I Induced - S				
Modified (IRRI) - R Induced - S				
Modified (UPLB) - R Induced - I				

Modified (PhilRice) - I Induced - S				
Modified (PhilRice) - I Induced - S				
Modified (PhilRice) - I Induced - S				
Modified (PhilRice) - I Induced - S				
Modified (IRRI) - I Induced - S				
S				
S				
				19.70%
R				
R				
MS				
R				
R				

Modified (UPLB) - I Induced - S				
Modified (CARGIL Phils.) - I Induced - S				
Modified (PhilRice) - S Induced - S				23.50%
S				22.80%
S				21.50%
S				24.40%
S				25.40%
S				22.00%
Modified (PhilRice) - S Induced - S				24.80%
Modified (IRRI) - I Induced - S				25.40%
Modified (IRRI) - S Induced - S				26.20%

Modified (IRRI) - S Induced - S				26.20%
Modified (UPLB) - I Induced - S				26.50%
Modified (PhilRice) - S Induced - S				24.10%
Modified (PhilRice) - S Induced - S				23.80%
Modified (IRRI) - R Induced - S				25.60%
Modified (IRRI) - S Induced - S				28.70%
Modified (PhilRice) - S Induced - S				17.60%
Modified (IRRI) - S Induced - S				24.70%
Modified (PhilRice) - S Induced - S				18.70%
Modified (IRRI) - S Induced - S				25.80%
Modified (PhilRice) - S Induced - S				20.90%
Modified (IRRI) - I Induced - S				20.90%
Modified (IRRI) - S Induced - S				20.60%
Modified (IRRI) - S Induced - S				21.70%
Modified (IRRI) - S Induced - S				22.20%
Modified (IRRI) - I Induced - S				20.90%

Modified (PhilRice) - R Induced - S				20.10%
Modified (IRRI) - S Induced - I				22.50%
Modified (Bayer) - S Induced - S				20.90%
Modified (MONSANTO) - I Induced - S				23.00%
Modified (PhilRice) - S Induced - S				17.50%
Modified (PhilRice) - R Induced - S				18.30%
Modified (SL Agritech) - S Induced - S				23.40%
Modified (UPLB) - S Induced - S				4.50%
Modified (PhilRice) - S Induced - S				5.10%
Modified (PhilRice) - I Induced - S				2.60%
Modified (PhilRice) - S Induced - S				22.10%
Modified (PhilRice) - S Induced - S				1.80%
S				22.80%
S				19.80%
S				18.40%
S				17.00%

S				19.10%
S				22.30%
Modified (IRRI) - S Induced - S				22.10%
Modified (PhilRice) - S Induced - S				23.20%
Modified (PhilRice) - S Induced - S				28.90%
Modified (PhilRice) - S Induced - S				22.80%
Modified (IRRI) - S Induced - S				22.90%
Modified (IRRI) - I Induced - S				21.60%
Modified (PhilRice) - S Induced - S				15.70%
Modified (BIOSEED) - S Induced - S				24.80%
Modified (HyRice Seed) - S Induced - S				25.60%
Modified (Syngenta) - S Induced - S				22.30%
Modified (Bayer) - S Induced - S				23.20%
Modified (IRRI) - I Induced - I				15.30%
Modified (IRRI) - S Induced - S				16.40%

Modified (PhilSCat) - S Induced - S				23.60%
Modified (PhilSCat) - S Induced - S				14.00%
Modified (PhilSCat) - S Induced - S				19.70%
Modified (BIOSEED) - S Induced - S				23.70%
Modified (IRRI) - S Induced - S				21.50%
Modified (PhilRice) - S Induced - S				19.10%
Modified (PhilRice) - S Induced - S				21.20%
Modified (PhilRice) - S Induced - S				18.60%
Modified (PhilRice) - S Induced - S				18.20%
Modified (IRRI) - S Induced - S				23.50%
S				22.60%
Modified (PhilRice) - S Induced - S				18.90%
Modified (PhilRice) - S Induced - S				23.10%
Modified (Bayer) - S Induced - S				22.90%

Modified (PhilRice - UPLB) - I Induced - S				20.20%
Modified (PhilRice - UPLB) - S Induced - S				20.20%
Modified (IRRI) - S Induced - S				24.10%
Modified (Syngenta) - S Induced - S				16.80%
Modified (Pioneer) - S Induced - S				19.70%
Modified (IRRI) - S Induced - S				21.70%
Modified (IRRI) - S Induced - S				22.40%
Modified (PhilRice) - S Induced - S				20.30%
Modified (PhilRice) - S Induced - S				12.23%
Modified (IRRI) - S Induced - S				18.10%
Modified (IRRI) - I Induced - S				23.90%
Modified (PhilRice) - S Induced - S				19.00%
Modified (PhilRice) - S Induced - S				19.40%
Modified (Hyrice) - S Induced - S				21.40%

Modified (IRRI) - S Induced - S				21.60%
Modified (IRRI) - S Induced - S				20.60%
Modified (Syngenta) - S Induced - S				12.70%
Modified (Seedworks) - S Induced - S				16.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				23.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				2.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				13.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				22.60%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				17.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				23.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.20%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				17.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.30%
Induced (PhilRice CES) - S				18.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				25.00%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.70%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				26.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				22.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				27.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice CES) - R Modified (PhilRice Midsayap) - S Modified (PhilRice Isabela) - I Modified (IRRI) - S				22.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice CES) - S Modified (PhilRice Isabela) - I Modified (IRRI) - S				26.90%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S Modified (VSU) - S Induced (PhilRice CES) - S				19.67%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				21.20%
Modified (PhilRice) - S Induced - S				19.10%
Modified (PhilRice) - I Induced - S				18.00%
Modified (PhilRice) - S Induced - S				21.00%
Induced (IRRI) - S				23.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				19.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				20.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				24.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				15.60%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				20.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				22.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				19.70%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				20.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				19.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				22.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				21.80%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				17.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) -S Modified (IRRI) - S				19.60%
Induced (PhilRice) - S				20.00%
Induced (IRRI) - S				21.30%
Induced (IRRI) - S				20.70%
Induced (PhilRice) - S				6.50%
Induced (PhilRice) - S				21.60%
Induced (IRRI) - S				24.00%
Induced (IRRI) - S				21.40%
Induced (PhilRice) - S				21.60%
Induced (IRRI) - S				25.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.70%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - I				21.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - I Modified (IRRI) - R				22.10%
S				19.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - I Modified (IRRI) - R				19.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.00%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.70%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				24.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				22.40%
Induced (PhilRice CES) - SInduced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.00%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - I				23.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				22.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.20%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				23.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				23.20%
Induced (IRRI) - S				21.00%
Induced (IRRI) - S				20.00%
Induced (PhilRice CES) - S Induced (UPLB) - S				20.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (IRRI) - S				12.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (IRRI) - S				20.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (IRRI) - S				16.70%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (IRRI) - S				19.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				13.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				13.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.50%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				21.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				20.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - I				19.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				18.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.00%
S				
S				

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S				12.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S Modified (BIARC) - I				19.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				19.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - R				17.80%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				24.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				22.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.10%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				24.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				19.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				22.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				14.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S Modified (BIARC) - S				20.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - R				18.30%
Induced (PhilRice CES) - S				18.30%
Induced (PhilRice CES) - S				20.20%
Induced (PhilRice CES) - S				19.10%

Induced (PhilRice CES) - S				14.20%
Induced (PhilRice CES) - S				22.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S				22.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S				2.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - R Modified (BIARC) - I				19.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				20.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S				15.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S Modified (IRRI) - S				13.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - R				13.00%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S				0.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S				14.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				15.30%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				14.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				20.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - R Modified (IRRI) - S				20.90%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				22.30%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				23.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S				23.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				14.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S Modified (IRRI) - S				21.40%
Modified (PhilRice Midsayap) - S Modified (BIARC) - R				17.40%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - R				19.10%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				20.00%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - R				14.80%

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - I Modified (BIARC) - R				23.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - S				20.70%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				21.50%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				20.20%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (IRRI) - S				23.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				20.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (BIARC) - I				21.00%
S				18.60%

S				21.40%
I				22.20%
S				11.60%
S				18.60%
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (CBES) - R				14.4% Low
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (CBES) - I				22.1% High
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (CBES) - I				18.6% Intermediate

Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - I Modified (CBES) - R				17.5% Intermediate
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - I Modified (CBES) - I				21.5% Intermediate
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (CBES) - R				20.5% Intermediate
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - S Modified (CBES) - S				13.9% Low
Induced (PhilRice CES) - S Induced (UPLB) - S Modified (PhilRice Midsayap) - I Modified (CBES) - I				18.6% Intermediate
PhilRice CES - S				13.9% Low
PhilRice CES - S				14.7% Low

PhilRice CES - S				19.2% Intermediate
Induced - S				17.8% Low
Induced - S				16.9% Low
Induced - S				15.4% Low
Induced - S				15.4% Low
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - I CBES (modified) - S				22.2% High

PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - S CBES (modified) - I				21.2% Intermediate
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - S CBES (modified) - R				20.9% Intermediate
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - S CBES (modified) - S				19.2% Intermediate
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - S CBES (modified) - S				19.5% Intermediate
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - S CBES (modified) - S				19.7% Intermediate

PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - S CBES (modified) - R				18.2% Intermediate
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - I CBES (modified) - R				17.2% Low
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - I CBES (modified) - S				12.1% Low
PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - I CBES (modified) - I				14.4% Low

PhilRice CES (induced) - S UPLB (induced) - S PhilRice Midsayap (modified) - I CBES (modified) - S				16.5% Intermediate
IRRI (induced) - S IRRI(modified) - I				16.2% L
IRRI (induced) - S IRRI(modified) - I				18.1% I
PhilRice (induced) - S PhilRice (modified) - I				17.2% I
UPLB (induced) - S UPLB (modified) - I				22.2% H
S				20.3% I
S				20.1% I

S				19.8% I
S				19.6% I
S				13.8% L
S				16.6% L
S				17.8% I

S				17.2% I
Induced - S Modified- R				20.9% I
S				21.9% I
Induced- S Modified-I				12.2% L
S				19.1% I

Induced- S Modified-I				19.7% I
S				22.2% H
S				18.6% I
Induced- S Modified-R				18.0% I
Induced- S Modified-I				17.0% L
Induced- S Modified-I				16.7% L
Induced- S Modified-I				19.4% I
S				12.4% L
Induced- S Modified-I				12.9% L
Induced- S Modified-R				13.4% L
Induced- S Modified-R				12.9% L

Induced- S	Modified-R				23.0% H
Induced- S	Modified-I				11.9% L
Induced- S	Modified-R				19.5% I
Induced- S	Modified-I				20.4% I
Induced- S	Modified-I				24.2% H
Induced- S	Modified-R				23.9% H
Induced- S	Modified-R				20.1% I
Induced- S	Modified-R				21.1% I
S					19.0% I

Induced- S Modified-I				18.6% I
Induced- S Modified-R				20.7% I
Induced- S Modified-R				14.8% L
Induced- S Modified-I				20.1% I
Induced-I				20.5% I
Induced-S				23.1% H
Induced-S				18.5% I
Induced-S				22.1% H
Induced-S				19.0% I
Induced-S				18.4% I
Induced-S				18.5% I

S				17.5% I
Induced-S Modified-S				24.1% H
Induced-S Modified-R				22.0% I
Induced-S Modified-S				20.1 % I
Induced-S Modified-S				15.2% L
Induced-S Modified-S				13.7% L
Induced-S Modified-S				13.5% L
Induced-S Modified-S				14.3% L
Induced-S Modified-S				19.3% I
Induced-S Modified-S				20.7% I

Induced-S Modified-S				18.4% I
Induced-S Modified-S				20.4% I
Induced-S Modified-R				20.4% I
Induced-S Modified-S				17.0% L
Induced-S Modified-S				20.9% I
Induced-S Modified-R				23.5% H
Induced-S Modified-S				23.5% H
Induced-S Modified-S				20.8% I
Induced-S Modified-S				21.9% I
Induced-S Modified-S				19.0% I
Induced-S Modified-R				13.7% L

Induced-S Modified-S				13.3% L
Induced-S Modified-R				18.5% I
Induced-S Modified-R				23.2% H
Induced-S Modified-S				23.8% H
Induced-S Modified-R				21.3% I
Induced - S Modified - I				20.8% I
Induced - S Modified - R				20.30%
Induced - S Modified - R				23.30%
Induced - S Modified- R				20.50%
Induced - S Modified- R				21.30%
Induced - S Modified- R				20.70%

Induced - S Modified - I				19.90%
Induced - S Modified - I				18.70%
Induced - S Modified - I				23.80%
Induced - S Modified - R				23.20%
Induced - S Modified - I				16.80%
Induced - S Modified - R				17.80%
Induced - S Modified - R				16.50%
Induced - S Modified - R				16.40%
Induced - S Modified - R				18.60%
Induced - S				20% - I
Induced - S Modified - S				20.4% - I

Induced - S Modified - S				14.10%
Induced - S Modified - S				14.10%
Induced - S Modified - S				14.00%
Induced - S Modified - S				12.20%
Induced - S Modified - S				12.20%
Induced - S Modified - S				12.50%
Induced - S Modified - I				20.00%

Induced - S Modified - R				19.60%
Induced - S Modified - R				12.70%
Induced - S Modified - I				12.40%
Induced - S Modified - I				15.80%
Induced - S Modified - R				18.70%
Induced - S Modified - R				21.80%

Induced - S Modified - R				19.80%
Induced - S Modified - I				23.10%
Induced - S Modified - R				19.80%
Induced - S; Modified - R				14.50%
Induced - S; Modified - I				13.80%

Induced - S; Modified - R				13.20%
Induced - S Modified - R				23.40%
Induced - S; Modified - I				19.80%
Induced - S; Modified - R				24.40%
Induced - S; Modified - I				18.30%

Induced - S; Modified - R				14.30%
Induced - S; Modified - R				18.20%
Induced - S;Modified - R				22.00%
Induced - S; Modified - I				17.70%
Induced - S; Modified - R				13.10%

Induced - S; Modified - I				19.10%
Induced - S; Modified - R				19.80%
Induced - S; Modified - R				21.90%

REGISTERED RICE VA

Oryza sativa

Agro-climatic, Seasonal and Cultural Adaptations
Early Lowland
Med Late Lowland
Early Lowland
Med Late Lowland
Early Lowland
Early Upland
Late Upland
Late Upland
Early Upland
Early Upland
Late Upland
Med Late Lowland
Late Upland

Lowland
Upland
Upland
Lowland
Upland
Upland
Lowland
Lowland
Lowland
Lowland
Lowland
Upland
Lowland
Lowland
Lowland

Irrigated lowland
Irrigated lowland, medium tall
Irrigated lowland
Irrigated lowland
Irrigated lowland
Rainfed
Upland culture
Upland culture
Upland culture
Upland culture
Irrigated lowland
Upland culture
Rainfed lowland
Irrigated lowland
Irrigated lowland
Upland culture
Irrigated lowland
Irrigated lowland, both dry and wet season
Irrigated lowland, both dry and wet season
Irrigated lowland
Irrigated lowland, Luzon only
Adapted to all regions of Philippines, dry or wet season
General countrywide adaptation
General countrywide adaptation
Irrigated lowland
Irrigated lowland

Grown only in Palawan for both wet and dry season, irrigated lowland.
For rainfed lowland drought prone areas.
For rainfed lowland drought prone areas.
Dry seeded rainfed lowland environment from May to June.
Adaptability to both dry and wet season.
For general planting in irrigated lowland areas.
For general planting is irrigated lowland but suitable planting for wet season.
For rainfed lowland areas, dry seeding culture.
Recommended for planting of this hybrid material but in the dry season only.
Recommended to both dry and wet seasons & majority of irrigated lowland area.
Adpatable to irrigated lowland conditions both dry and wet seasons.
Overall performance indicate the adaptability to irrigated lowland both dry and wet season.

Irrigated lowland drought prone areas transplanting culture.
Rainfed lowland drought prone areas, transplanting culture.
Rainfed lowland drought prone environment transplanting culture.
Rainfed lowland drought prone areas, transplanting culture.
Rainfed lowland drought prone dry seeding culture.
Adaptable to low-medium elevation areas and for dry season.
Irrigated lowland areas for wet season.
Saline prone irrigated lowland dry and wet season.
Saline prone irrigated lowland.
Irigated lowland dry and wet season.
Irrigated lowland dry and wet better w/ wet season.
Irrigated lowland dry and wet seasons.
Highly favorable irrigated lowland areas dry and wet season with caution to hot spot areas.
For rainfed lowland areas dry seeding culture, wet season.
For rainfed lowland dry seeding culture.
Irrigated lowland.
Irrigated lowland

Rainfed lowland dry seeding culture.
Rainfed lowland dry seeding culture.
Irrigated lowland
For irrigated lowland areas suitable both dry and wet season.
Irrigated lowland wet and dry seasons.
For irigated lowland, transplanting culture.
For irrigated lowland dry and wet season planting.
For irrigated lowland dry and wet season planting.
Recommendation for saline prone irrigated lowland area.
Recommendation for saline prone irrigated lowland area.
Saline prone irrigated lowland.
Irrigated lowland medium salinity, wet dry season.
Cool elevated areas wet season low to medium elevation.
Cool elevated areas wet season cultivation and low to medium elevation.

Cool elevated areas wet cultivation medium elevation.
Rainfed lowland transplanted both under favorable and drought condition.
Ranfed lowland transplanted.
Upland
Irrigated lowland.
Rainfed transplanted, drought and favorable environment.
Suitable to medium elevation less than 1000 above sea level in both DS and WS.
Suitable for saline prone areas with moderate salinity.
Suitable for saline prone irrigated lowland areas.
Suitable for upland areas.
Suitable for upland areas specifically in areas similar to the agro-climatic condition of Nueva Ecija, Isabela, Bukidnon, Zamboanga del Surand South Cotabato.
For irrigated lowland dry and wet season planting.
For irrigated lowland dry and wet season planting.
Irrigated lowland good in both DS & WS transplanting culture.
Moderate resistant to stamborer(DH) and moderate susceptible (WH) green leafhopper, yellow stemborer & brown planthopper.
Stop gap variety for tungro hot spot in region 6.10.11.

Stop gap varieties for tungro hot spot in Region 6, 10, 11 and ARMM.
Irrigated lowland areas under both favorable condition
Recommended in irrigated lowland particularly in Nueva Ecija, Cagayan Bohol & Isabela specially during wet season.
Recommended in irrigated lowland for dry and wet season.
Irrigated lowland, dry and wet season.
Irrigated lowland, dry and wet season both for transplanting and direct seeding condition.
Recommended in irrigated lowland particularly Bohol. Cagayan during wet and dry season.
Irrigated lowland, dry and wet seasons.
Irrigated lowland, dry and wet seasons.
Irrigated lowland, wet and dry seasons.
Irrigated lowland dry and wet seasons.
Irrigated lowland dry and wet seasons.
Irrigated lowland transplanted culture in both wet and dry season.
Irrigated lowland better adaptability to direct seeding wet and dry season.
Irrigated lowland, transplanting and direct seeding culture wet and dry season.
Irrigated lowland, transplanting and direct seeding culture but could also be used in transplanting system.

Irrigated lowland better adaptability to direct seeding wet and dry season.
Adaptable to favorable irrigated lowlands in 1,2,3 and also to saline prone areas in cagayan and solar radiation.
Favorable irrigated lowland dry and wet seasons.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Favorable irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Irrigated lowland, dry and wet season, transplanting and direct seeding culture.
Suitable to irrigated lowland areas in Cagayan, Isabela, Nueva Ecija and Davao provinces preferably during the dry season under transplanting culture.
Suitable to irrigated lowland areas in Nueva Ecija, Isabela, Davao provinces, Bohol and Bukidnon and in similar areas where IR64 is adapted.
Japonica type adapted to tropical conditions in irrigated lowland areas.
Irrigated lowland, transplanting culture suitable to both DS and WS with caution on the susceptability to bacterial diseases.

Irrigated lowland, can be planted in both dry and wet seasons, transplanting culture.
Irrigated lowland. Can be planted in both dry and wet seasons. Transplanting culture.
Irrigated lowland, can be planted in both seasons but preferable wet season, transplanting culture.
Irrigated lowland, transplanting culture, dry season adaptation.
Irrigated saline prone areas.
Irrigated saline prone areas.
Irrigated saline prone areas, dry season only, transplanting culture, weak resistance to insect pests and diseases, not to be planted in wet season.
Irrigated saline prone areas, dry & wet seasons transplanting culture.
Irrigated saline prone areas, dry & wet seasons, transplanting culture not to be planted in areas hot spot to tungro virus.
Rainfed lowland areas, good for dry seeding and transplanting culture.
Flood prone areas with complete submergence up to 14 days at seedling to vegetative stage wet season adaptation, transplanted culture.
Dry season only transplanting culture.
Irrigated lowland areas in N.Ecija, Isabela, Cagayan, Davao Del sur, Bukidnon & Bohol & similar areas during dry season only transplanting culture.
Irrigated lowland areas in N.Ecija, Isabela, Cagayan, Davao Del sur, Bukidnon & Bohol & similar areas during dry season only transplanting culture adaptation.

Irrigated lowland areas in N.Ecija, Isabela, Cagayan, Davao Del sur & del norte, Gen. Santos, Bukidnon & Bohol & similar areas
Irrigated lowland areas in N.Ecija, Isabela, Cagayan, Davao Del sur & del norte, Gen. Santos, Bukidnon & Bohol & similar areas
National recommendation in irrigated lowland areas wet and dry seasons under tansplanting culture.
Recommended in Isabela, Gen. Santos and Bukidnon similar growing conditions preferable during the dry season under transplanting culture.
Found to excel in Nueva Ecija, Cagayan & Laguna, Bohol, Dav. Del sur, Bukidnon. For national release but only for dry season under transplanting culture
Irrigated lowland dry & wet seasons, transplanting & direct wet seeding culture.
Irrigated lowland dry & wet but preferably dry seasons, transplanting & direct wet seeding culture.
Irrigated lowland areas, dry and wet seasons under transplanting or direct wet seeding culture.
Favorable irrigated lowland areas, preferably dry seasons, transplanting culture.
Irrigated lowland preferably during the DS (to be sown before Dec 15) under transplanting culture.
For irrigated lowland adaptation, dry and wet seasons with very good potential during the WS. It is suitable under transplanting or direct wet seeding culture.
Generally adapted to irrigated lowland transplanted culture preferably in the DS.
Wide adaptation in majority of test locations in Luzon, Visayas and Mindanao, can be planted in both DS and WS but preferably DS; performed better as trasnplanted crop.
National recommendation in irrigated lowland areas, preferably WS under transplanting culture.

National recommendation , irrigated lowland areas during the DS under transplanting culture.
Wide adaptation irrigated lowland areas in Luzon, Visayas and Mindanao test sites except region 2; with good performance in both DS and WS under transplanting culture.
Location specific adaptation during DS, transplanting culture in irrigated areas of Davao del Sur, Cagayan, Bohol, Davao Oriental and Bukidnon.
Recommended in irrigated lowland areas in Davao del Sur and Bohol during the dry season under transplanting culture.
National recommendation in irrigated lowland areas in both dry season and wet season.
Irrigated lowland, dry and wet seasons but preferably DS can be direct seeding or transplanting culture.
Irrigated lowland, better during DS, transplanted culture.
Distinctly recommended in DS ut can also be planted in WS in irrigated lowland areas.
National recommendation, dry and wet seasons, irrigated lowland, transplanting culture.

National recommendation, dry and wet seasons, irrigated lowland.
For national recommendation in irrigated lowland areas, wet and dry seasons preferably in Nueva Ecija, Laguna, Bohol, General Santos and Bukidnon and similar areas of same climatic condition.
Location season specific recommendation in irrigated lowland areas in Nueva Ecija, Cagayan, and Bohol.
Location specific recommendation dry and wet but distinct WS performer, irrigated lowland areas, such as Laguna Nueva Ecija, Cagayan in Luzon. Bohol in the Visayas and Davao del Sur, General Santos City.
Distinctly WS performer (July-Dec) in irrigated lowland areas across Luzon, Bohol in the Visayas and in Davao del Sur and Bukidnon in Mindanao.
Irrigated lowland. Location season specific adaptation, preferably in Nueva Ecija, Isabela,Cagayan, Bohol, Davao del Sur and Bukidnon.
Distinctly wet season recommendation in irrigated lowland areas in Nueva Ecija, Cagayan, Bohol, General Santos and Bukidnon.

Location specific recommendation, dry and wet seasons in irrigated lowland areas in Laguna, Nueva Ecija, Bohol and Bukidnon.
For dry season recommendation in irrigated lowland areas particularly in Cagayan, Bohol, Bukidnon and Nueva Ecija.
Wet season recommendation in irrigated lowland areas in Bohol and Bukidnon. Can also be planted in Nueva Ecija Isabela, Cagayan and Davao del Sur which produced positive yield of at least 6 t/ha.
Location specific recommendation in irrigated lowland areas particularly in Cagayan, Bohol and Bukidnon and in WS in General Santos City.
Location specific recommendation in irrigated lowland areas of Isabela and Davao del Sur and preferably in WS in Bukidnon.
Location specific recommendation in irrigated lowland areas in Cagayan and Bohol.
Rainfed lowland favorable and drought prone.

Rainfed lowland favorable and drought prone.
Rainfed lowland areas with prolonged rainfall.
Rainfed lowland favorable and drought prone.
Rainfed lowland, dry seeding and in areas with prolonged rainfall.
Rainfed lowland areas with prolonged rainfall.
Rainfed lowland favorable and drought prone, seeding.
Rainfed lowland favorable and drought prone areas in Luzon, some areas in Visayas and Mindanao.

Rainfed lowland favorable and drought prone, wet season dry seeding culture.
National recommendation in upland areas, wet season, dry seeding culture.
Saline-prone lowland (coastal areas, transplanted culture, dry and wet seasons.
Saline-prone coastal lowland areas with moderate salinity level, dry and wet seasons.
Saline-prone, coastal lowland areas, dry and wet season, transplanting culture.
Saline-prone, coastal lowland areas with slight to moderate salinity level, dry and wet seasons.
Irrigated lowlands, dry and wet seasons under direct seeding culture.
Irrigated lowlands, dry and wet seasons, both for direct seeding and transplanting culture.
Irrigated lowlands, dry and wet seasons, both for direct seeding and transplanting culture.
Dry and wet seasons, irrigated lowland areas.

Location-season specific recommendation in irrigated lowlands in Bohol (DS/WS) and Iloilo and Leyte in the DS.
(DS/WS) transplanting and direct seeding culture.
Location season specific recommendation in irrigated lowland in Bohol (Ds/WS) and VSU (DS). These are the only sites which performed in at least 50% of the valid trials in the Visayas.
Location specific recommendation in the provinces of Visayas particularly in Bohol and VSU (DS/WS).
Location-season specific recommendation in irrigated lowland in USM & WESVIARC (WS), CMU and PhiolRice Agusan (DS/WS).
Location-soeacon specific recommendation in irrigated lowland Mindanao particularly in the province of Davao de Sur (USM), General Santos, Zamboanga Sibugay(WESMIARC) and Bukidnon (CMU) (DS/WS) and Visayas particularly in Bohol (DS) and Leyte (VSU) (DS/WS).
National recommendation in irrigated lowland areas, dry and wet seasons, transplanting culture.

Location specific recommendation in irrigated lowland in Bohol and WESVIARC(DS/WS) and VSU (WS).
location-season specific recommendation in irrigated lowland in Bohol and VSU (DS/WS).
Saline prone lowland coastal areas, transplanted, dry and wet seasons.
Saline prone lowland coastal areas, dry and wet seasons, transplanting culture.
Saline prone lowland coastal areas, transpalnted, dry and wet seasons.
Saline prone lowland coastal areas, transpalnted, dry and wet seasons.
Saline prone lowland coastal areas, transpalnted dry and wet seasons.
Saline prone lowland coastal areas, transpalnted , dry and wet seasons.
Saline prone lowland coastal areas, transpalnted dry and wet seasons.
Saline prone lowland coastal areas, transplanted, dry and wet seasons.
Saline prone lowland coastal areas, transplanted dry and wet seasons.
Irrigated lowland, dry and wet season planting.

Irrigated lowland, Dry and wet seasons planting, prefereably during DS.
Rainfed lowland, wet seasons, dry seeding culture
Rainfed lowland, wet seasons, dry seeding culture.
Location season specific recommendation in irrigated lowland in Bohol and Hagonoy (USM) both in DS & WS.
Upland areas, wet season planting.
National recommendation in upland areas, wet season planting.
Upland areas, wet season planting.
Adapted to both dry and wet seasons, transplanting culture.
National recommendation in irrigated lowland areas, dry and wet seasons, transplanting culture.

Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Samar, Aklan and Bohol , dry and wet seasons, transplanting culture.
Dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Aklan, and Bohol, dry and wet seasons, transplanting culture.
Dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol, Iloilo and Leyte, dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol, Iloilo and Leyte, dry and wet seasons, transplanting culture.
National recommendation in irrigated lowland areas, dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol, Iloilo, Leyte and Negros, dry and wet seasons, transplanting culture.

Location-specific recommendation in irrigated lowland in Visayas particularly in the provinces of Bohol, Iloilo, Leyte and Negros, dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol, Iloilo, Negros and Leyte, dry and wet seasons, transplanting culture.
National recommendation in irrigated lowland areas, dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol, Iloilo, Leyte, and Negros, dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol, Iloilo and Leyte, dry and wet seasons, transplanting culture.
National recommendation in irrigated lowland areas, dry and wet seasons, transplanting culture.
Location-specific recommendation in irrigated lowland areas in Visayas particularly in the provinces of Bohol and Leyte, dry and wet seasons, transplanting culture.

National recommendation in irrigated lowland areas, dry and wet seasons, transplanting culture.
For dry season recommendation in irrigated lowland areas, transplanting culture.
Saline prone lowland coastal areas, transplanted, dry and wet seasons.
Saline prone lowland coastal areas, transplanted, dry and wet seasons.
Irrigated lowland areas in Luzon, particularly in the province of Isabela, Pangasinan, Cagayan and Ilocos Norte, dry and wet seasons, direct wet seeding culture (DWSR).
Regional recommendation in irrigated lowland areas in Visayas particularly in the provinces of Samar and Bohol, dry and wet seasons, direct wet seeding culture.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Pangasinan, Cagayan and Oriental Mindoro, dry and wet seasons, direct wet seeding culture.
National recommendation, irrigated lowland areas, transplanting culture, dry and wet season.

Regional recommendation in irrigated lowland areas in the province of Samar, dry and wet seasons, transplanting culture and in the provinces of Luzon particularly in Pangasinan, Cagayan, Ilocos Norte and Tabuk, Kalinga, dry and wet seasons, direct seeding culture.
Regional recommendation in irrigated lowland areas in Luzon provinces such as Nueva Ecija, Isabela, Laguna, Pangasinan, Kalinga and Ilocos Norte.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Laguna, Nueva Ecija, Isabela, Kalinga, Cagayan, Oriental Mindoro, Pangasinan and Ilocos Norte.
Regional recommendation in irrigated lowland areas in Luzon in the provinces of Isabela, Cagayan, Pangasinan, Kalinga and Ilocos Norte.
Regional recommendation in irrigated lowland areas in Mindanao particularly in the provinces of Davao del Sur, Zamboanga Sibugay, North Cotabato and Agusan del Norte.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Cagayan, Kalinga and Ilocos Norte.
National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in rainfed lowland areas, wet season planting.

National recommendation in rainfed lowland areas, wet season planting.
National recommendation in rainfed lowland areas, wet season planting.
National recommendation in rainfed lowland areas, wet season planting.
National recommendation in rainfed lowland areas, wet season planting.
National recommendation in rainfed lowland areas, wet season planting.
National recommendation in rainfed lowland areas, wet season planting.
National recommendation in rainfed lowland areas, wet season planting.
National recommendation in irrigated lowland areas during the dry season.
National recommendation in rainfed lowland areas, wet season planting.

Regional Recommendation in Luzon paticularly in the provinces of Isabela, Pangasinan, Tarlac, Cagayan, Tabuk, Kalinga, Palawan and Ilocos Norte and Visayas in the provinces of Samar and Bohol, dry and wet seasons,transplanting and direct seeding culture.
Regional recommendation in irrigated lowland areas in Luzon particulary in the provinces of Isabela,Pangasinan, Tarlac, Cagayan, Tabuk, Kalinga, Bataan and Palawan and Visayas in the provinces of Samar and Bohol, dry and wet seasons, transplanting culture.
National recommendation in irrigated lowland areas, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particulary in the provinces of Isabela, Pangasinan, Cagayan, Ilocos Norte and Bataan and Mindanao particularly in Bukidnon, Agusan del Norte and Zamboanga, Sibugay.
Recommended for dry season planting in irrigated lowland areas.
National recommendation in irrigated lowland areas, dry and wet seasons.
Recommended for dry season planting in irrigated lowland areas.

Recommended for dry season planting in irrigated lowland areas.
Recommended for dry season planting in irrigated lowland areas.
Regional recommendation in Luzon paticularly in the provinces of Nueva Ecija, Isabela, Cagayan, Oriental Mindoro, Los Baños,Laguna,Pangasinan and Ilocos Norte,dry and wet seasons.
Regional recommendation in Luzon, Visayas and Mindanao.
Regional recommendation in irrigated lowland areas in Luzon, Visayas and Mindanao, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
Recommended for saline prone lowland coastal areas, dry and wet seasons.
Recommended for saline prone lowland coastal areas, dry and wet seasons.
Recommended for saline prone lowland coastal areas, dry and wet seasons.

Recommended for saline prone lowland coastal areas, dry and wet seasons.
Recommended for saline prone lowland coastal areas, dry and wet seasons.
National recommendation in rainfed lowland areas.
National recommendation in rainfed lowland areas.
National recommendation in rainfed lowland areas.
National recommendation in rainfed lowland areas.
National recommendation in rainfed lowland and saline prone areas (dual adaptation).
National recommendation in irrigated lowland, dry and wet seasons planting.
National recommendation in irrigated lowland, dry and wet seasons planting.

National recommendation in irrigated lowland, dry and wet seasons planting.
Regional recommendation in irrigated lowland areas in Luzon in the provinces of Nueva Ecija, Isabela, Cagayan, Albay and Kalinga.
Regional recommendation in irrigated lowland areas in Bohol in the provinces of North Cotabato and Agusan del Norte in Mindanao.
Regional recommendation in irrigated lowland areas in Luzon in the provinces of Nueva Ecija, Isabela, Cagayan, and Ilocos Norte and in the provinces of Leyte and Bohol in Visayas.
Regional recommendation in irrigated lowland areas in Visayas particularly in the provinces of Leyte, Bohol and Iloilo.
Regional recommendation in irrigated lowland areas in Mindanao in the provinces of Hagonoy, Bukidnon and Agusan del Norte.
Regional recommendation in irrigated lowland areas in Luzon in the provinces of Nueva Ecija, Isabela, Cagayan and Ilocos Norte.

Regional recommendation in irrigated lowland areas in Bohol and Zamboanga Sibugay.
Regional recommendation in irrigated lowland areas in Bohol and Iloilo.
Regional recommendation in irrigated lowland areas in Nueva Ecija, Cagayan, Albay, Kalinga and Ilocos Norte.
Regional recommendation in irrigated lowland areas in Leyte and Iloilo.
Regional recommendation in irrigated lowland areas in the provinces of Iloilo and Samar and other similar growing conditions, dry and wet season.
Regional recommendation in irrigated lowland areas in Luzon in the provinces of Isabela, Cagayan, Kalinga, Or. Mindoro, Camarines Sur and Palawan, and other areas with similar growing conditions, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Iloilo in Visayas and Bukidnon in Mindanao and in other areas with similar growing conditions.
National recommendation in irrigated lowland areas (dwrs) and other similar areas with similar growing conditoins.

Regional recommendation in irrigated lowland areas in Luzon in the provinces of Isabela, Pangasinan, Ilocos Norte, Kalinga, Oriental Mindoro and Palawan and in Samar and Iloilo in Visayas and other areas with similar growing conditions.
National recommendation in irrigated lowland areas during the dry season.
National recommendation in irrigated lowland areas during the dry season.
Regional recommendation in irrigated lowland areas in Bukidnon and Agusan del Norte in Mindanao and other areas with similar growing conditions.
Regional recommendation in irrigated lowland areas in Luzon in the provinces of Nueva Ecija, Isabela, Cagayan and Kalinga and Mindanao areas in North Cotabato, Zamboanga, Sibugay, Bukidnon, Agusan del Norte and in other areas with similar growing conditions.
National recommendation in irrigated lowland areas, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Agusan del Norte in Mindanao, and other areas with similar growing conditions.
Recommended in saline prone irrigated lowland areas, dry and wet seasons.

Recommended in saline prone irrigated lowland areas, dry and wet seasons.
Recommended in saline prone irrigated lowland areas, dry and wet seasons.
Recommended in saline prone irrigated lowland areas, dry and wet seasons.
Recommended in saline prone irrigated lowland areas, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Mindanao particularly in the provinces of Agusan del Norte and Bukidnon and other areas with similar growing conditoin.
Regional recommendation in irrigated lowland areas in Visayas in the provinces of Iloilo, Leyte and Bohol and other areas with similar growing condition.
Regional recommendation in irrigated lowland areas in Mindanao particularly in the provinces of Agusan del Norte and Bukidnon and other areas with similar growing conditoin.

National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Visayas particularly in the provinces of Leyte and Iloilo and other areas with similar growing condition
National recommendation in irrigated lowland areas, dry and wet seasons.
Recommended in saline prone irrigated lowland areas, dry and wet seasons.
Recommended in saline prone irrigated lowland areas, dry and wet seasons.

Recommended in saline prone irrigated lowland areas, dry and wet seasons .
Cool elevated areas, low to medium elevation, especially during the dry season.
Cool elevated areas, low to medium elevation, dry season cultivation.
Cool elevated areas, dry season cultivation, low to medium elevation.
Cool elevated areas, wet season cultivation, low to medium elevation.
Rainfed lowland, wet season, dry seeding culture

Rainfed lowland, wet season, dry seeding culture
Rainfed lowland, wet season, dry seeding culture
Rainfed lowland, wet season, dry seeding culture
Rainfed lowland, wet season, dry seeding culture
Rainfed lowland, wet season, dry seeding culture

Regional recommendation in irrigated lowland areas in Bohol and Iloilo in Visayas; Bukidnon, Agusan del Norte and Zamboanga Sibugay in Mindanao and in areas with similar growing condition, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Samar and Bohol in Visayas and in areas with similar growing condition, dry and wet seasons.
Irrigated lowland, dry and wet seasons
Regional recommendation in irrigated lowland areas in Nueva Ecija, Cagayan, Pangasinan, Kalinga, Ilocos Norte, Albay and Isabela in Luzon; Agusan, Davao, Bukidnon and North Cotabato in Mindanao and in areas with similar growing condition, dry and wet seasons.

National recommendation in irrigated lowland areas, dry and wet seasons.
Flood prone areas , wet season, transplanted culture.
National recommendation in rainfed lowland areas, wet season, dry seeding culture.
National recommendation in rainfed lowland areas, wet season, dry seeding culture.
National recommendation in rainfed lowland areas, wet season, dry seeding culture.
National recommendation in rainfed lowland areas, wet season, dry seeding culture.
National recommendation in high temperature-prone in irrigated lowland areas.

National recommendation in high temperature-prone in irrigated lowland areas.
National recommendation in saline prone irrigated lowland areas, dry and wet seasons.
National recommendation in saline prone irrigated lowland areas, dry and wet seasons.
National recommendation in saline prone irrigated lowland areas, dry and wet seasons.
National recommendation in saline prone irrigated lowland areas, dry and wet seasons.

National recommendation in saline prone irrigated lowland areas, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija,Isabela,Ilocos Norte, Cagayan, Albay and Pampanga during dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Ilocos Norte, Cagayan, Tabuk and Pampanga during dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela,Ilocos Norte, Or. Mindoro, Cagayan, Pampanga, Albay, and Tabuk during dry and wet seasons.
Regional recommendation in Luzon in irrigated lowland areas particularly in the provinces of Isabela, Nueva Ecija, Cagayan, Kalinga and Ilocos Norte, dry and wet seasons. With limitations on tungro-infected areas.

National recommendation in irrigated lowland areas, direct wet-seeding culture, dry and wet seasons
National recommendation in irrigated lowland areas, direct wet-seeding culture, dry and wet seasons
National recommendation in irrigated lowland areas, direct wet-seeding culture, dry and wet seasons
National recommendation in irrigated lowland areas, direct wet-seeding culture, dry and wet seasons
National recommendation in irrigated lowland areas, direct wet-seeding culture, dry and wet seasons
National recommendation in irrigated lowland areas, direct wet-seeding culture, dry and wet seasons
Regional recommendation in irrigated lowland areas in Mindanao particularly in the provinces of Bukidnon, Agusan del Norte, Zamboanga Sibugay and North Cotabato transplanting culture; and in Bohol APC in Visayas, direct wet-seeding Culture.
Regional recommendation in irrigated lowland areas in Visayas particularly in the provinces of Iloilo and Bohol, direct seeding culture, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons

National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons.
National recommendation in rainfed lowland areas, dry seeding culture
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Ilocos Norte, Albay, Bulacan, Pangasinan and Pampanga and in the Visayas provinces in Bohol, Leyte, Negros Occidental, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Albay, Bulacan and Pangasinan dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Cagayan, Ilocos Norte, Pampanga, Bulacan and Pangasinan dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Cagayan, Ilocos Norte, Pampanga, Albay and Pangasinan and in the provinces of Iloilo, Leyte, Negros and Bohol in Visayas, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Isabela, Cagayan, Ilocos Norte, Pampanga, Albay and Pangasinan, dry and wet season.

Regional recommendation in irrigated lowland areas in Luzon particularly in the provinces of Nueva Ecija, Cagayan, Camarines Sur, Albay, Ilocos Norte, Isabela, Bulacan and Pangasinan, dry and wet seasons
Regional recommendation in irrigated lowland areas in Luzon particularly in the province of Nueva Ecija, Isabela, Cagayan, Bulacan, Pampanga, Albay, Ilocos Norte, dry and wet seasons.
Regional recommendation in irrigated lowland areas in Mindanao particularly in the provinces of Bukidnon, North Cotabato, Agusan del Norte and Davao del Norte, dry and wet seasons.
National recommendation in irrigated lowland areas, dry and wet seasons
Regional recommendation in irrigated lowland areas in Visayas particularly in Bohol, direct wet seeding culture.
National recommendation in irrigated lowland areas, direct wet-seeding culture.
National recommendation in irrigated lowland areas, direct wet-seeding culture.
National recommendation in irrigated lowland areas, direct wet-seeding culture.
Regional recommendation in irrigated lowland areas in Mindanao particularly in the provinces of Bukidnon, Agusan del Norte, Zamboanga Sibugay and North Cotabato and in the provinces of Bohol and Iloilo in Visayas, direct wet seeding culture.
National recommendation in irrigated lowland areas, direct wet seeding culture.
National (subject to existing rules and regulations on GM crops)

National recommendation in irrigated lowland areas.
National recommendation in saline prone irrigated lowland areas.
Recommended in irrigated lowland areas in Central Luzon and Western Mindanao.
Recommended in irrigated lowland areas in Central Luzon
Recommended in irrigated lowland areas in North West Luzon, Central Luzon and Eastern Mindanao
Recommended in irrigated lowland areas in North East Luzon, Central Luzon, Southern Luzon, Western Visayas and Southern Luzon, Western Mindanao and Eastern Mindanao.
Recommended in irrigated lowland areas in North West Luzon, Central Luzon and Southern Luzon.
Recommended in irrigated lowland areas in North East Luzon, Central Luzon, Southern Luzon and Western Mindanao.
Recommended in irrigated lowland areas in North West Luzon and Western Visayas and Southern Luzon.
Recommended in irrigated lowland areas in Central Luzon and Western Mindanao.

Recommended in irrigated lowland areas in Central Luzon.
Recommended in irrigated lowland areas in Central Luzon.
Recommended in irrigated lowland areas in Central Luzon and Western Mindanao.
Recommended in irrigated lowland areas in Western Mindanao.
Recommended in irrigated lowland areas in North East Luzon and Central Luzon.

Recommended in irrigated lowland areas in Central Luzon

Recommended in irrigated lowland areas of North East Luzon, North West Luzon, Central Luzon, Southern Luzon, Western Visayas and Southern Luzon, Eastern Mindanao
Recommended in irrigated lowland areas of Southern Luzon and Western Mindanao
Recommended in irrigated lowland areas of Southern Luzon
Recommended in irrigated lowland areas of Central Luzon and North West Luzon
Recommended in irrigated lowland areas of Central Luzon.

Recommended in irrigated lowland areas of Central Luzon.
Recommended in national irriigated lowland areas.
Recommended in national rainfedlowland areas.
Recommended in national rainfedlowland areas.
Recommended in national rainfedlowland areas.
Recommended in national irrigated lowland areas.
Recommended in national irrigated lowland areas.
Recommended in irrigated lowland areas in North East Luzon
Recommendded in irrigated lowland areas in Southern Luzon
Recommended in irrigated lowland areas in North East Luzon
Recommended in irrigated lowland areas in Zamboanga Peninsula and North Southern Eastern and North Central Mindanao

Recommended in irrigated lowland areas in Zamboanga Peninsula, North East Luzon, North West Luzon, and North South Eastern and North Central Mindanao
Recommended in irrigated lowland areas in Zamboanga Peninsula, North West Luzon, and North South Eastern and North Central Mindanao
Recommended in irrigated lowland areas in Zamboanga Peninsula, North East Luzon, and North South Eastern and North Central Mindanao
Recommended in irrigated lowland areas in Zamboanga Peninsula and North South Eastern and North Central Mindanao
Recommended in irrigated lowland areas in North South Eastern and North Central Mindanao, North West Luzon
National recommendation in irrigated lowland areas
National recommendation in irrigated lowland areas
National recommendation in irrigated lowland areas
National recommendation in irrigated upland areas
National recommendation in irrigated lowland areas
Recommended in irrigated lowland areas in North West Luzon and North South Eastern and North Central Mindanao

Recommended in irrigated lowland areas in North West Luzon, Southern Luzon, Zamboanga Peninsula
Recommended in irrigated lowland areas in North East Luzon, North West Luzon, Central Luzon
Recommended in irrigated lowland areas in North East Luzon, North West Luzon, Central Mindanao,
Recommended in irrigated lowland areas in Central Mindanao, Zamboanga Peninsula and North South Eastern
Recommended in irrigated lowland areas in Zamboanga Peninsula and North South Eastern and
Recommended in irrigated lowland areas in Zamboanga Peninsula and North Central Mindanao
Recommended in irrigated lowland areas North Eastern Luzon and North South Eastern and Central Mindanao

Recommended in irrigated lowlan areas in National level
Recommended in irrigated lowland areas of North East Luzon (NEL) and Zamboanga Peninsula and Western Mindanao (ZPWM)
Recommended in irrigated lowland areas of Southern Luzon (SL), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of North East Luzon (NEL), Southern Luzon (SL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of Zamboanga Peninsula and Western Mindanao (ZPWM)
Recommended in irrigated lowland areas of North West Luzon (NWL), Central Luzon (CL), Southern Luzon (SL), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)

Recommended in irrigated lowland areas of North East Luzon (NEL), Central Luzon (CL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM).
Recommended in irrigated lowland areas of North East Luzon (NEL), Central Luzon (CL), Southern Luzon (SL), Central Mindanao (CM)
Recommended in irrigated lowland areas of Central Luzon (CL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM) North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of North West Luzon (NWL), Zamboanga Peninsula and Western Mindanao (ZPWM,) North South Eastern and North Central Mindanao (NSENCM), Southern Luzon (SL)
Recommended in irrigated lowland areas of North East Luzon (NEL), North West Luzon (NWL), Central Luzon (CL), Southern Luzon (SL), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)

Recommended in irrigated lowland areas of North East Luzon (NEL), North West Luzon (NWL), Central Luzon (CL), Southern Luzon (SL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of Central Luzon (CL), Southern Luzon (SL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM)
Recommended in irrigated lowland areas of North East Luzon (NEL), North West Luzon (NWL), Central Luzon (CL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of Southern Luzon (SL), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of North West Luzon (NWL)

Recommended in irrigated lowland areas of North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of Central Mindanao (CM)
Recommended in irrigated lowland areas of North East Luzon (NEL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of Southern Luzon (SL): 7888, Zamboanga Peninsula and Western Mindanao (ZPWM): 8729

Recommended in irrigated lowland areas of Zamboanga Peninsula and Western Mindanao (ZPWM)
Recommended in irrigated lowland areas of North East Luzon (NEL), Central Mindanao (CM), Zamboanga Peninsula and Western Mindanao (ZPWM), North South Eastern and North Central Mindanao (NSENCM)
Recommended in irrigated lowland areas of Zamboanga Peninsula and Western Mindanao(ZPWM)

ARIETIES

L.

Remarks and other Varietal Characteristics

Head rice recovery 40.4%, amylose content intermediate, fair eating quality

Amylose content intermediate, fair eating quality
High head rice recovery, high amylose content, good eating quality
Glutinous variety with bold grains comparable to native glutinous rice
High tillering capacity and eating quality is highly adaptable
Milling recovery is satisfactory with medium slender grain and good eating quality
Grain is medium long, slender and succulent
Grain is medium long, slender and succulent
Grain is medium long, slender and succulent
Grain is slender, medium and translucent, resistant to lodging
Grain is long and slender
Medium long and slender grains of very high acceptibility (93%)
Has an excellent grain quality with intermediate amylose content, thus comparable to C-4-63 in appearance
Long slender grain
92% acceptability when raw, 72% when cooked
Better milling recovery and acceptability than IR-36
Acceptability is 92.5%, slightly higher than malagkit sungsong's 89%
Better in quality and more acceptable th an IR-50, long slender grain
Higher milling recovery and more accepatble than IR-50, high stable yields across seasons/locations
Eating quality is more acceptabel than IR-50, high stable yields across seasons/locations
Grain quality is long, slender and translucent, milling recovery high
Variety of areas practice multiple cropping, medium long slender grain
Has better lodging resistance and better stable yield than other early maturing varieties.
Highly acceptable and has higher and stable yield than malagkit sungsong.
Higher and stable yield potential, better grain quality and lodging resistance, tolerant to alkalinity, salinity and phosphorous deficiency, long slender grain
Has wide adaptability, recommended for general planting with certain limitatins in areaaas with high pressure blast.
Showed superiority over IR-50, high amylose content with long slender grain.

High milling and head rice recovery, low gelatinization temperature, excellent long, slender translucent grains
High milling and head rice recovery, acceptable as raw and cooked, high amylose content, intermediate gelatinization temperature
High milling recovery, high amylose content and low gelatinization temperature, excellent long, slender and translucent grains.
Slightly high milling recovery but high head rice recovery, high amylose content, intermediate gelatinization temperature, excellent medium long and translucent grain.
Slightly lower headrice and milling recovery, high percentage of chalkiness, eating quality is less preferred.
Milling recovery 64.25%, Head rice 48.41%, Chalky 4.49%, Amylose Content 23.48%
Milling recovery 65.33%, Head rice 35.78%, Amylose content 27.18%, Acceptability 68.2% for cooked, 76.8% for raw
Milling recovery 64.71%, Head rice 41.53%, Chalky 2.78%, Amylose content 23.85%
Milling recovery 66.84%, Head rice 47.66%, Chalky 8.59%, Amylose content 28.2%
Milling recovery 66.93%, Head rice 49.65%, Chalky 4.82%, Amylose content 27.6%
Milling recovery 66.62%, Head rice 38.42%, Chalky 7.06%, Amylose content 26.86%
Milling recovery 66.75%, Head rice 52.39%, Chalky 5.02%, Amylose content 23.75%
Milling recovery 65.78%, Head rice 51.87%, Chalky 5.1%, Amylose content 24.33%
Acceptable grain quality
80% grain acceptability when cooked and 71.7% when raw.
79.1% grain acceptability when cooked and 87.3% when raw.
76.88% grain acceptability when cooked and 77.94% when raw.
76.88% grain acceptability when cooked and 77.94% when raw.
68.9% grain acceptability when cooked and 89.6% when raw.
The selection has higher yield potential and better lodging resistance. It has better grain qualities especially its milling potentials, physical attributes and sensory properties.
Higher potential yield with comparable disease and insect reactions than IR 72. Also with superior grain quality traits than IR72 and even better than IR64 in sensory qualities.
Has high acceptability in both cooked and raw form. Eating quality is better than PSB Rc4.

Has better physico-chemical traits, superior sensory qualities but with comparable milling and physical properties than IR74.
Higher yield potential in unfavorable rainfed lowland.
Comparable grain quality characteristics, however, it has better eating quality.
Comparable physio-chemical properties, physical attributes and milling potentials and with better sensory qualities (raw and cooked form).
Earliness and drought resistance at vegetative growth are the outstanding merits of MRC22929-24-2. It has better physico-chemical properties which relate to the high preference and acceptability than C4-137.
Has a higher yielding potentials earlier maturity, than the traditional varieties, and it has comparable grain qualities.
Moderate resistance to cool temperature during the reproductive phase but its earliness escapes the cold spell.
Medium maturing type whose salt tolerance is the best among the entries tested even in other soil related stresses.
Medium early maturing line which gave better yield than IR74. Its chalky character makes it less preferred in raw form.
Its low amylose of 19.7%, high milling, and headrice recoveries has a potential higher market value better than IR72 and comparable to IR68.
Early maturing and high yielding line. It is more adaptable during the wet season with higher % acceptability in the cooked and raw forms than IR72 and IR64. It has intermediate amylose.
Early maturing line is well suited for planting in both dry and weat seasons and registered wide.
It is medium early maturing selection that is comparable to IR68. It also has excellent grain quality especially in terms of milling potentials and sensory qualities.
High yielding ability, it has moderate resistance to major insect pest and diseases, good milling and sensory qualities.
High yield and resistance to blast and other major pests, it has prominently better grain qualities.
This medium maturing line is recommended for dry season planting although satisfactory yiekd were also obtained during wet season. Better eating quality that IR 64 with highest preferemce and % acceptability
This medium maturing line is recommended for dry season planting although satisfactory yield were also obtained during wet seasonpreferemce and % acceptability. It has better preference and % acceptability in the raw form than IR 64.

This early maturing line has 14.6% yield advantage over UPL Ri7. Comparable with check variety regarding grain quality characteristics.
This early maturing line produced 10.0% yield advantage over UPL Ri7. This line showed better physio-chemical characteristics, % headrice recovery and % acceptability form than UPL Ri7.
It has good grain quality with high acceptability in both cooked and raw forms.
This medium early maturing line produced higher yield than the check, UPL Ri5. It matures earlier than UPL Ri5. It is highly acceptable in raw form.
This hybrid line which is medium maturing showed higher yields in the wet season with yield advantage of 16.20% over IR64 in Phase I and 11.0% in Phase II in 23 sites nationwide.
It has excellent grain quality in terms of milling potentials and sensory qualities. It is highly preferred both in the raw and cooked forms than IR72 and comparably acceptable with IR 64.
This hybrid gave superior performance at specific locations, notably, Camarines Sur, Laguna and Bukidnon with an average yield superiority of 23.8% across seasons over the high yielding check variety IR72.
It has intermediate amylose content, better than IR72, and comparable in terms of milling and physical attributes. It is highly acceptable both in cooked and raw forms.
It has intermediate amylose content, better than IR72, It has good eating quality in terms of cooked and raw forms.
This line showed superior performance across locations over the two checks varities; IR68 and PSB Rc28 with an average yield advantage of 10.0%. It has intermediate amylose content and with good grain qualities ccomparable to PSB Rc28, especially in terms of milling potential. It is more preffered in raw form and highly acceptable both in cooked and raw forms than PSB Rc28.
It has good grain qualities, comparable to PSB Rc48 but generally better thana PSB Rc50. Is is highly preferred than PSB Rc48 and PSB Rc50 in cooked forms and highly acceptable in raw forms.
It has good grain qualities, comparable to PSB Rc48 but generally better thana PSB Rc50. Is is highly preferred than PSB Rc48 and PSB Rc50 in cooked forms and highly acceptable in raw forms.
This line can be best recommend as variety because of its earliness and excellent eating quality. It is exhibited an intermediate amylose content, generally better than PSB Rc48. It is highly preferred both in cooked and raw forms than PSB Rc48and PSB Rc50.
High performance score and percent acceptability in raw and cooked forms. In 15 farmers' field, yield advantage of 5-100% was obtained in 13 locations.
With better percentage headrice than PSB Rc46 and very high percentage acceptability in raw and cooked form. Comparable or better than PSB Rc46 in 4 trials in 9 farmers' field.
Better headrice recovery than PSB Rc46. High acceptability in cooked form. Comparable yield with PSB Rc46 in 6 trials in 9 farmes' field.

Higher headrice recovery than PSB Rc46. High percentage acceptability score both in raw and cooked form. Comparable or better yield than PSB Rc46 in 6 trials in 9 farmers' field.
High percet acceptability for raw and cooked form. Yield comparable or better than the highest yielding check in 9 trials in 14 farmers' field.
Very high percentage acceptability in both raw and cooked form. Yield comparable with highest yielding check in 7 trials in 15 farmers' field.
Intermediate amylose content. Yield comparable or higher than the yielding check in 9 trials in the farmers' field.
Highly acceptable than IR72 in cooked form.
Comparable to highest yielding check in 4 trials in 11 farmers' field.
As to grain quality, this japonica rice selection has low amylose and hard gel consistency better than some of the traditional varieties such as Pinidua and Tinawen.
As to grain quality, it passed all the standars for physio-chemical, physical, milling and sensory qualities.
Except for the low amylose content, all grain quality characteristics are within the recommended value. As to percent acceptability, it is highly acceptable both in cooked and raw forms.
This line had high amylose content with medium gel consistensy. It also passed the standards for grain quality particularly on milling potentials. It was acceptable both in cooked and raw forms.
This line showed intermediate amylose content with acceptable eating quality.
Intermediate amylose content. Good milling recovery. Highly acceptable in cooked and raw forms.
Good milling and headrice recovery. With intermidiate amylose content.
Intermediate amylose content, acceptable in cooked and raw forms.
Intermediate amylose content and acceptable milling and headrice recovery.
Fairly acceptable yield under both transplanted and direct seeding culture. Suitable for both TPR and DSR.

Acceptable grain qualities with intermediate amylose.
Acceptable grain quality and high yield potential.
Intermediate amylose content. Acceptable milling potentials. Better preference score in the cooked form and acceptability rating in both raw and cooked forms than IR64, PSB Rc72H and PSB Rc18.
Intermediate amylose content. Highly acceptable both in cooked and raw forms.
Slightly aromatic. Good milling and headrice recovery. Highly acceptable in cooked and raw forms.
Photoperiod insensitive. Acceptable milling recovery.
Intermediate amylose content. Good and headrice recovery. Has very high cooking expansion. Better preference score in the raw form over PSB Rc72H and PSB Rc28.
Acceptable milling and head recovery. Better preference score and accepatbility rating in cooked form than IR65.
Acceptable grain qualities.
Good milling and headrice recovery. Acceptable both in cooked and raw forms.
Has good grain eating quality, having intermediate amylose content and low gelatinization temperature. PJ21 apparently retained the good grain and eating quality characteristics of its parents IR64, Burdagol and Sasanishiki.
Intermediate good grain quality.
Good milling quality and milling potentials. Fairly high consumer acceptability in both raw and cooked forms.
Good eating quality. Good milling potentials and physical attributes. And good ratooning ability.
Adaptability to both transplanting and direct seeding systems. Acceptable grain quality characteristics.
Good milling quality and physical attributes. Low amylose and with good eating quality.

Intermediate amylose content and with good eating quality. Highly acceptable and preferred in both raw and cooked forms. With acceptable milling and physical attributes.
Require optimum fertilization to avoid blast problem. Suggest to harvest at 80% maturity to reduce shattering losses. Good eating quality. High milling and headrice recovery.
Intermediate amylose content. Good milling recovery. Highly acceptable especially in the raw form. Superior in aroma in cooked form as compared to ordinary rice.
Intermediate amylose content. Good milling and headrice recovery. Acceptable both in cooked and raw forms.
Good milling and heardice recovery. Highly acceptable in the raw form.
Intermediate amylose content. Good milling and headrice recovery. Acceptable both in cooked and raw forms.
Intermediate amylose content, with good milling and headrice recovery. Preffered both in cooked and raw forms as compared to PSB Rc82. Diverse parentals.
Superior performance across all seasons in MAT trial. Intermediate amylose content. Highly acceptable both in the cooked and raw forms.
High milling recovery. High percentage in the cooked and raw forms as compared to PSB Rc82.
Good grain quality and eating quality.
Acceptable milling and physical attrributes. High amylose but with low gel temperature score. Acceptable in cooked and raw forms.
High milling recovery. Intermediate amylose content with good grain quality better than IR64. Better preference and acceptability in raw forms than IR 64 and PSB Rc82
Good milling recovery. High acceptability in raw and cooked forms. With intermediate amylose and excellent eating quality like IR64.
Early maturity with yield of at least 4t/ha, resistant to blast, very good milling potential, excellent eating quality.
High iron content, early maturity for crop intesification, good field resistanfce to RTV, intermediate to blast, sheath blight BPH biotypes 2 and , excellent seeding vigor.

Good milling and head rice recovery, early matuing selection at 112 days, intermediate amylose content, intermediate to blast, sheath blight, GLH and WH
Early maturing at 113 days, good milling and headrice recovery, high percentag of acceptability in cooked and raw forms.
Acceptable both in cooked and raw forms, good milling recovery, intermediate reaction to blast, early maturing at 104 at DAS earlier by at least 5 days during WS
High percentage of acceptability in cooked and raw forms, good milling headrice recovery, intermediate amylose content, early maturing at 110 days
Intermediate amylose content. Godd milling recovery.
Good milling recovery. Preffered both in cooked and raw forms. High percentage acceptability both in cooked and raw forms.
Intermediate amylose content. Good milling and headrice recovery. Long and slender grain. Acceptable both in cooked and raw forms.
Premium milling recovery, very good headrice. Percent acceptability comparable to the check variety.
Very good headrice and milling recovery. Long and slender grain. High percentage acceptability both in cooked and raw forms.
Intermediate amylose content. Very good milling recovery. High percentage acceptability as compared to IR64 and PSB Rc14.
High tolerance to complete submergence at seedling to vegetative stage up to 14 days and retaining the good traits of high yield eating quality.
Good milling ability, long and slender grain. Highly acceptable both in cooked and raw forms.
Intermediate amylose content. Good milling recovery. High percentage acceptability in the raw form as compared to the check variety.
Intermediate amylose content. Good milling recovery, long and slender grain. Highly preferred both in cooked and raw forms as compared to the check variety.

Good milling and headrice recovery, long and slender grain. Highly preferred especially in the raw form as compared to the check varieties.
Intermediate amylose content. Good milling and headrice recovery. Long and slender grain. High percentage acceptability both in cooked and raw form as compare to check varieties.
Intermediate amylose content. Hih percentage milling recovery. Long and slender grain. High percentage acceptability in the raw form.
High percentage milling and headrice recovery inspite of high chalky character. Preferred in cooked and raw forms as compared to the check varieties. High acceptability both in cooked and raw forms. Better eating quality than IR64.
High milling recovery, long and slender grain. Highly preferred with high acceptability both in cooked and raw forms. Better eating quality than IR64.
Intermediate amylose content. Good milling recovery. Preferred both in cooked and raw forms as compared to the check variety. High percentage acceptability in the cooked and raw forms.
Intermediate amylose content. Good milling recovery. Preferred both in cooked and raw forms as compared to the check variety. High percentage acceptability in the cooked and raw forms.
Intermediate amylose content. Good milling and headrice recovery. Long and slender grain. High percentage acceptability especially in the raw form.
Comparable yield performance against the check variety, Burdagol. Very good milling and headrice recovery. Aromatic, high sensory preference and acceptability, tender, tasty.
Good milling recovery. Eating quality comparable to Koshihikari.
Intermediate amylose content. Good milling recovery. Long and slender grain. Highly preferred in the raw form as compared to the check variety.
Early maturing 111 days, good milling and headrice recovery with long slender grains.
Early maturing at 112 days as TPR and 104 days as DWSR, very good milling recovery with llong and intermediate grain.
Very early maturing days at 108 days, intermediate amylose content with high milling potentials and acceptable physical attributes

Very early maturing at 108 days, intermediate amylose content with good milling potentials.
Early maturing at 110 days, intermediate amylose content, long slender grain, exhibited better yield performance in 6 out of 10 sites
Early maturing at 115 days, intermediate reaction to blast, bacterial leaf blight, white stemborer, yellow stemborer and brown planthopper under induced method
Early maturing at 114 days, moderate resistant to WSB, good milling and headrice recovery with long and slender grain.
Premium milling recovery with fair brown rice and grade 1 headrice recovery. Intermediate amylose content with long and slender grain. Very high percentage acceptability in cooked and raw forms as compared to IR64 and NSIC Rc138 owing to better eating quality.
Excellent milling potentials with premium milling headrice recovery. Intermediate amylose content with long and intermediate grain. High percentage acceptability in the cooked and raw forms as compared to the check variety, NSIC Rc138.
High milling and headrice recovery with medium and intermediate grain. Slightly aromatic when cooked.
Good milling and headrice recovery. Highly acceptable in the cooked form as compared to the check variety, MS11.
Intermediate amylose content with long and slender grains. Percent acceptability in cooked and raw forms comparable to the inbred check, PSB Rc82.

Good grain quality, low amylose content, high milling potentials and highly acceptable in cooked and raw forms.
Good grain quality, with intermediate amylose content, high milling potentials with long and slender grains. High acceptability in the raw form.
Intermediate amylose content with long and slender grains, acceptable milling potentials. High percentage acceptability both in cooked and raw forms as compared to the inbred check, PSB Rc82 comparable to IR64.
Intermediate amylose content with good milling and headrice recovery. Long and slender grain with better sensory qualities to PSB Rc82 in terms of whiteness, translucency in raw form. In cooked form it is comparable to the hybrid check Mestiso7.
Intermediate amylose content with good milling potentials. Comparable to check variety PSB Rc18 in terms of sensory qualities.
Intermediate amylose content with long and slender grains. Very good milling potentials. High acceptability in the raw form but comparable only in the cooked form than PSB Rc82.
Good grain quality, high milling recovery, intermediate amylose content, with long and slender grains. Comparable to IR64, the eating quality check in both cooked and raw forms and with high percentage acceptability as compared to the inbred check, PSB Rc82.

Long and slender grain with good milling potentials. High percentage acceptability in the cooked and raw forms as compared to PSB Rc82.
Very good milling potentials with long and slender grains. High percentage of acceptability both in cooked and raw forms as compared to the inbred check, PSB Rc82.
Very good milling recovery, with intermediate amylose content. Highly acceptable both in cooked and raw forms as compared to the inbred check, PSB Rc18, and comparable in terms of raw form to IR64, the eating quality check.
High milling recovery with long and slender grains. Highly acceptable in the cooked form as compared to the two check varieties, PSB Rc82 and Mestiso 7 and comparable to IR64, the eating quality check.
High milling recovery with long and slender grains. Highly acceptable in the cooked form as compared to the inbred check, PSB Rc18, and comparable to IR64, the eating quality check in cooked form.
Intermediate amylose content with extra long and slender grains. Grade 1 milling recovery. High percentage acceptability especially in the raw form as compared to inbred check, PSB Rc82.
Intermediate amylose content with high milling recovery.

Long and slender grain with high milling recovery.
Long and slender grain with high milling and headrice recovery. Comparable to IR64, The eating quality check in the raw form.
Intermediate amylose content, long and immediate grain. High milling recovery. High percentage acceptability in the raw form as compared toIR64 and PSB Rc14.
Intermediate amylose comtent with extra long and slender grain. Grade 1 milling recovery. Aromatic with high percentage acceptability especially in the cooked form as compared to IR64, the eating quality check and PSB Rc14. Slightly aromatic in the raw form.
Very good milling recovery and headrice recovery, long and slender grain. Comparable in terms of raw to the check varieties, IR64 and PSB Rc14, but inferior in eating quality.
Intermediate amylose content with extra long and slender grain. High milling recovery with acceptable headrice and physical attributes. Highly preferred in the raw form as compared to the two check varities, IR64 & PSB Rc14.
Premium milling recovery, fair brown rice and grade 1 milling recovery with long and slender grains. Very high percentage acceptability in the raw form as compared to IR64 and PSB Rc14.

High milling and headrice recovery. Better sensory qualities in cooked form, comparable in raw form than UPL Ri7.
Intermediate amylose content. Good milling recovery.
High milling and headrice recovery with fair broqn rice. Highly acceptable in the cooked and raw forms as compared to PSB Rc50.
High milling and headrice recovery and fair brown rice with long and slender grains. Comparable quality in the raw form than PSB Rc50 but inferior in eating quality.
Good grain quality with high milling and headrice recovery, intermediate amylose with long and slender grain. Comparable with PSB Rc50 in the raw form.
Intermediate amylose content with medium and slender grains. Premium milling recovery and Grade 1 headrice recovery, and fair brown rice. Better sensory qualities in cooked and raw forms in terms of whiteness, glossiness and translucency.
Intermediate amylose content, grade 1 milling recovery with long and slender grains. High percentage acceptability both in cooked and raw forms than PSB Rc82 vand comparable to IR64.
Intermediate amylose content with good headrice and premium milling recovery. High percentage acceptability both in cooked and raw forms as compared to PSB Rc83 and with comparable to eating quality of IR64.
Intermediate amylose content with extra long and slender grains. Fair brown rice and grade 1 milling recovery with less inferior eating quality than IR64.
Grade 1 milling and headrice recovery with short and bold grains. High percentage acceptability especially in the cooked form, comparable to MS11 and IR64.

Intermediate amylose content with extra long and slender grains. Highly acceptable both in cooked and raw forms as compared to the check variety, PSB Rc82. High percentage acceptability in the cooked form as compared to IR64, the eating quality check.
Intermediate amylose content (22.2%) with long (7.1mm) and slender grain (3.2). Premium milling recovery (71.1%) fair brown rice (78.1%) and grade 1 headrice recovery (55.2%). Highly acceptable especially in the cooked form as compared to IR64, the eating quality check.
Slightly aromatic with intermediate amylose content (19.1%). Premium melling recovery (70.6%), fair brown (78.9%) and grade 2 headrice recovery (48.2%).
Intermediate amylose content (20.1%), fair brown rice (76.6%) and grade 1 milling (68.6%), and headrice recovery (48.1%).
Intermediate amylose content (19.1%), fair brown rice (76.5), grade 1 milling (68.0) and headrice recovery (49.2). Highly acceptable especially in the cooked form as compared to IR64, the eating quality check.
Intermediate amylose content (22.4%) with low to intermediate gel temp score. Grade 1 milling recovery (68.9%) and fair brown rice (76.6%). Long (7.4mm) and slender grain (3.3).
Intermediate amylose content (21.8%) with long (7.4mm) and slender grain (3.3). Premium miling recovery (70.9%), fair brown rice (78.1%) and grade 1 headrice recovery (52.8%).

Grade 1 milling (69.5%) and headrice recovery (49.7%) with fair brown rice (77.6%). High percentage acceptability especially in the cooked form as compared to IR64, the eating quality check and PSB Rc82.
Intermediate amylose content (19.6%) with long (7.0mm) and slender grain (3.1). Grade 1 milling recovery (68.4%) and grade 2 (43.9%) headrice recovery. High percentage acceptability both in cooked and raw forms as compared to the check variety, PSB Rc82 and IR64, the eating quality check.
Intermediate amylose content (20.0%). Grade 1 milling recovery (66.2%), fair brown rice (76.5%) and grade 2 headrice recovery (44.5%). Highly acceptable boyh in cooked and raw forms.
Intermediate amylose content (21.3%). Long (7.0mm) and slender grain (3.2). Grade 1 milling recovery (68.0%) and fair brown rice (77.4%).
Intermediate amylose content (20.7%) with long (7.0mm) and slender grain (3.0). Grade 1 milling recovery (68.7%), fair brown rice (76.8%) and grade 2 headrice recovery (43.1%).
Grade 1 milling recovery (69.0%) and fair brown rice (76.5). Medium (6.3m) and intermediate grain (2.7). Highly acceptable both in cooked and raw forms as compared to IR64, the eating quality check.
Intermediate amylose content (21.6%). Grade 1 milling recovery (69.4%) and fair brown rice (78.2%). Long (7.0mm) and slender grain (3.2).
Intermediate amylose content (24.0%). Grade 1 milling recovery (67.1%) and fair brown rice (76.9%) Extra long (7.7mm) and slender grain (3.8). Highly accepatble both in cooked and raw forms.
Intermediate amylose content (21.4%) with long (7.4mm) and slender grain (3.7). Grade 1 milling recovery (66.7%) and fair brown rice (76.6%). High percentage acceptability both in cooked and raw forms as compared to IR64, the eating quality check.
Intermediate amylose content 921.6%). Grade 1 milling recovery (67.8%), fair brown rice (76.8%) and grade 2 headrice recovery (39.6%). High percentage acceptability in the raw form as compared to IR64, the eating quality check.
Grade 1 milling recovery (65.6%) and fair brown rice (75.9%). Long (7.2mm) and slender grain (3.1). Highly acceptable especially in the raw form as compared to IR64, the eating quality check.
Intermediate amylose content (20.7%), slightly aromatic when freshly cooked after 6 hours. Grade 1 milling (69.5%) and headrice recovery (55.2%). High percentage acceptability in the cooked form as compared to NSIC Rc128.

Slightly aromatic when freshly cooked and after 6 hours. Intermediate amylose content (21.9%) with long (7.2mm) and slender grain (77.8%). Grade 1 milling (66.5%) and headrice recovery (48.8%0 and fair brown rice (77.8%). High percentage acceptability in the cooked form as compared to the check variety, NSIC Rc128.
Intermediate amylose content (21.5%) with long (6.8mm) and slender grain (3.2). Slightly aromatic when freshly cooked and after 6 hours. Grade 1 milling recovery (68.8%). Fair brown rice (77.0%) and grade c2 headrice recovery (44.3%).
Intermediate amylose content (22.1%), grade 1 milling recovery (68.4%) and fair brown rice (76.7%). High percentage acceptability in the raw form as compared to IR64, the eating quality check.
Grade 1 milling and headrice recovery. High percentage acceptability in cooked forms as compared to the check variety, PSB Rc82 and IR64, the eating quality check.
Intermediate amylose content (19.3%) with medium (6.4mm) and intermediate grain (2.8). Grade 1 milling (69.1%) and headrice recovery (50.2%) and fair brown rice (75.7%). Acceptable in raw and cooked forms but less inferior than IR64 than IR64 and comparable to NSIC Rc192.
Intermediate amylose content (19.8%) with extra long (7.5mm) and slender grain (3.7). Grade 1 milling (68.2%), grade 2 headrice recovery (46.7%) and fair brown rice (76%). Comparable to IR64, the eating quality check in terms of raw and cooked forms.
Intermediate amylose content (18.5%) with long (7.1mm) and slender grain (3.3). Grade 1 milling (68.2%), headrice recovery (46.6%) and fair brown rice (76.8%). High percentage acceptability in the raw form as compared to the check variety, NSIC Rc192, and comparable to IR64 in both raw and cooked forms.
Intermediate amylose content (18.9%) with long (7.2 mm) and slender grain (3.3). Fair brown rice and grade 1 milling recovery.
Intermediate amylose content (21.0%) with long (6.9mm) and slender grain (3.0). High milling recovery (69.9%), fair brown rice (77.4%). Comparable to IR64, the grain quality check in the raw form but less acceptable in cooked form.

Intermediate amylose content (20.1%) with long (7.4mm) and slender grain (3.3). Fair brown rice (75.6%) and grade 2 milling (64%) headrice recovery (39.9%). Better acceptability in the raw and cooked forms than IR64.
Intermediate amylose content (19.7%) with long (6.8mm) and slender grain (3.2). Premium milling recovery (72.4%), fair brown rice (78.6%) and grade 1 (54.6%) headrice recovery. High percentage acceptability especially in the raw form as compared to IR 64, the grain quality check.
Intermediate amylose content (18.4) with long (6.8mm) and slender grain (3.2). Grade 1 milling recovery (65.8%). High percentage acceptability in the cooked form as compared to IR64.
Intermediate amylose content (18.5%) with long (6.7mm) and slender grain (3.2). Grade 1 milling (69.5%) and headrice recovery (50.7%) and fair brown rice (77.1%). Very high acceptability in the cooked form as compared to IR64, the eating quality check and PSB Rc 82 and Mestiso 7, the check varieties.
Premium milling recovery (70.7%), fair brown rice (78.3%) and grade 2 (47.7%) headrice recovery. Acceptable in cooked and raw forms but less inferior to IR64.
Premium milling recovery (70.4%) and fair brown rice (78.0%). Long (7.1mm) and slender grain (3.2). Acceptable in cooked and raw forms but less inferior to IR64 and Mestiso 7.
Intermediate amylose content (18.9%) with long slender grain (3.3). Premium milling recovery (70.4%), fair brown rice (77.5%) and grade 1 (52.2%) headrice recovery. High percentage acceptability both in cooked and raw forms as compared to IR64, the eating quality check and PSB Rc82, the check variety.
Intermediate amylose content (21.0%) with long (7.1mm) and slender grain (3.1). Grade 1 miling recovery (69.9%), farin brown rice (77.9%) and grade 2 (47.6%) headrice recovery. Comparable to PSB Rc82 in raw and cooked forms.

Intermediate amylose (20.9%) content with medium (6.1mm) and intermediate grade (2.5). Premium milling recovery (72.3%), fair brown rice (79.8%) and grade 1 headrice recovery (54.0%). Acceptable in raw and cooed forms but less inferior than IR64.
Premium milling recovery (70.8%), fair brown rice (78.0%) and grade 1 (51.9%) headrice recovery. Long (7.0mm) and slender grain (3.1). High percentage acceptability especially in the raw form as compared to IR64, the eating quality check and PSB Rc82 and Mestiso7, the check varieties.
Premium milling recovery (70.7%), fair brown (78.8%) and grade 2 headrice recovery (47.7%). Long (7.1mm) and slender grain (3.1). High percentage acceptability especially in the raw form as compared to the check variety, PSB Rc82 and IR64, the eating quality check. Percent acceptability in the raw form comparable to Mestiso 7.
Intermediate amylose content (20.1%) with long (7.3mm) and slender grain (3.5). Premium milling recovery (71.0%), fair brown rice (79.1%) and grade 1 (48.7%) headrice recovery. High percentage acceptability both in cooked and raw forms as compared to the check varieties, PSB Rc82 and Mestiso 7 and IR64, the eating quality check.
Intermediate amylose (20.1%) content with long (7.1mm) and slender grain (3.5). Premium milling recovery (71.8%), fair brown rice (78.9%) and grade 1 (55.1%) headrice recovery. Acceptable in raw and cooked forms but less inferior than IR64 and Mestiso 7.
Intermediate amylose content (18.1%) with long (7.2mm) and slender grain (3.3). Premium milling recovery (70.3%) fair brown rice (78.6%) and grade 1 (51.0%) headrice recovery. Highly acceptable both in cooked and raw forms as compared to IR64, the eating quality check and PSB Rc82 and Mestiso 7, the check varieties.
Intermidiate amylose content (19.2%) with long (7.3mm) and slender grain (3.1). Premium milling recovery (72.1%), fair brown rice (78.6%) and grade 1 (53.7%) headrice recovery. High percentage acceptability especially in the raw form as compared to the check varieties, PSB Rc82 and Mestiso 7 and IR64, the eating quality check.

Premium milling (71.4%) and headrice recovery (54.3%), fair brown rice (78.4%). Long (7.3mm) and slender grain (3.1). High percentage acceptability in the raw form as compared to IR64, the eating quality check and PSB Rc82 and Mestiso 19, the check varieties.
Premium milling recovery (71.7%), fair brown (78.8%) and grade 1 (50.8%) headrice recovery. Long (7.1mm) and slender grain (3.1). High percentage acceptability in the raw form as compared to IR64, the eating quality check, and PSB Rc82, the check variety.
Intermediate amylose content (21.0%) with long (7.2mm) and slender grain (3.3). Premium milling recovery (71.6%), fair brown rice (79.4%) and grade 1 headrice recovery (49.6%). Highly acceptable both in cooked and raw forms as compared to IR64, the eating quality check.
Intermediate amylose content (21.0%) with long (7.3mm) and slender grain (3.3). Premium milling recovery (72.5%), fair brown rice (79.2%) and grade 1 headrice recovery (50.8.6%). High percentage acceptability both in cooked and raw forms as compared to IR64, the eating quality check.
Intermediate amylose content (19.9%), fair brown rice (78.4%), long (7.2mm) and slender (3.2mm) grain. Premium milling (72.3%) and grede 1 (48.9%) headrice recovery. High percentage acceptability in the raw and cooked forms as compared to the check variety, NSIC Rc224 and IR64, the eating quality check.
Fair brown rice (77.3%) and grade 1 milling recovery (69.9%), premium headrice (62.5%) recovery with medium (6.3mm) and intermediate (2.7mm) grain shape. Better acceptability (83.3%) in the raw form against IR64 (79.6%).
Intermediate amylose content (20.5%) fair brown rice (78.0%) with long (6.7mm) and intermediate (2.6mm) grain shape. Premium milling (71.4%) ang grade 1 (50.5%) headrice recovery. High percentage acceptability especially in raw form as compared to the checkvariety, NSIC Rc224 and IR64, the eating quality check.
Grade 1 milling (69.3%) and headrice (53.8%) recovery, fair brown rice (76.2%), medium grain length (6.4mm) and intermediate (2.8mm) grain shape, slightly aromatic when cooked.

Intermediate amylose content (19.2%), long (6.8mm) and slender grain (3.0mm), premium milling (70.3%) and headrice (62.4%) recovery, fair brown rice (77.5%). Comparable % acceptability in cooked form to IR64 with better acceptability in the raw form.
Fair brown rice (76.8%), grade 1 milling (69.8%) and premium headrice recovery (57.2%), long (6.7mm) and slender (3.1mm) grain. Passed acceptability rating but not better than IR64.
Intermediate amylose content (20.0%), fair brown rice (79.9%) with medium (6.4mm) and intermediate (2.8mm) grain. Premium milling (73.3%) and grade 1 headrice (52.3%) recovery; passed acceptability rating but not better than IR 64.
Intermediate amylose content (19.8%) with long (7.1mm) and slender (3.4mm) grain. Premium milling (72%), grade 1 headrice (52.6%) recovery and fair brown rice (78.9%). Acceptable but not better than IR64.
Intermediate amylose content (19.9%) with long (6.9mm) and slender (3.4mm) grain. Premium milling recovery (72.0%), fair brown rice (79.4%) and grade 1 (49.2%) headrice recovery. High percentage acceptability especially in the cooked form as compared to IR64, the eating quality check.
Intermediate amylose content (19.3%) with long (6.9mm) and intermediate (3.0mm) grain. Premium milling recovery (71.6%), fair brown rice (78.6%) and grade 1 (54.5%) headrice recovery. High percentage acceptability both in cooked and raw forms.
Fair brown rice (75.3%), short (5.1mm) and bold (1.8mm) grain. Percent acceptability comparable to Koshihikari, the eating quality check.
Intermediate amylose content (19.5%) with medium (6.3mm) and intermediate (2.8mm) grain shape. Premium milling (71.2%) and grade 1 headrice (48.2%) recovery and fair brown rice (78.7%). Acceptability in cooked form comparable to IR64.

Intermediate amylose content (21.1%), fair brown rice (77.0%). Grade 1 milling (70.0%) and headrice (51.0%) recovery. Long (7.2mm) and slender (3.4mm) grain. High percentage acceptability both in cooked and raw forms as compared to two check varieties, PSB Rc14 and NSIC Rc192.
Intermediate amylose content (21.0%), fair brown rice (77.5%) and grade 1 milling (69.1%) and headrice (49.6%) recovery. Long (7.3mm) and slender (3.4mm) grain. High percentage acceptability in the cooked form as compared to the two check varieties, PSB Rc14 and NSIC Rc192.
Intermediate amylose content (19.9%), fair brown rice (78.2%), premium milling (70.5%) and headrice (57.4%) recovery. Long (6.8mm) and slender (3.3mm) grain, very high percent acceptability both in cooked and raw forms as compared to the check varieties, PSB Rc14, NSIC Rc192 and IR64, the eating quality check.
Intermediate amylose content (20.8%), fair brown rice (78.1%) with long (6.7mm) and slender grain (3.0mm). Premium milling (71.4%) and headrice (57.4%) recovery. High percentage acceptability both in cooked and raw form as compared to the two check varieties, PSB Rc14 and NSIC Rc192.
Intermediate amylose content (19.1%), fair brown rice (78.2%) and premium milling (71.0%) recovery, long (7.0mm) and slender (3.1mm) grain. High percentage acceptability both in cooked and raw forms as compared.
Intermediate amylose content (18.9%), fair brown rice (76.8%), with long (6.9mm) and slender (3.1mm) grain, grade 1 milling (69.9%) and headrice (50.8%) recovery. High percentage acceptability especially in the cooked and raw form as compared to the two check varieties, PSB Rc14 and NSIC Rc192.
Intermediate amylose content (19.0%), fair brown rice (77.9%) with long (6.7mm) and slender (3.1mm) grain. Premium milling (70.5%) and headrice (58.7%) recovery, high percentage acceptability both in cooked and raw forms as compared to the two check varieties, PSB Rc14 and NSIC Rc192 and IR64, the eating quality check.
Premium milliing (71.6%), fair brown rice (78.5%) and grade 1 headrice(52.6%). Extra log (7.5mm) and Slender (3.4mm) grain. High percentage acceptability in the raw form as compared to the check varieties, M1 and IR64, the eating quality check.
Intermediate amylose content (19.4%), fair brown rice (78.2%), premium miling (70.7%) and grade 1 headrice (48.1%), Long (7.2mm) and Slender (3.3mm) grain. High percentage acceptability especially in the cooked form as compared to the two check varities, PSB Rc14 and NSIC Rc192.

Premium milling recovery (70.7%),fair brown rice (77.3%) and grade 2 (42.8%) headrice recovery. Medium (6.4mm) and slender (3.2mm) grain shape. High percentage acceptability in cooked form as compared to the check variety, PSB Rc82 and IR64,the eating quality check.
Intermediate amylose content (19.3%), Premium milling recovery (72.6%),fair brown rice (77.9%) and grade 2 (40.7%) headrice recovery, medium (6.5mm) and intermediate (2.9mm) grain shape.High percentage acceptability in th raw form as compared to the check variety, PSB Rc82 and IR64, the eating quality check.
Intermediate amylose content (19.2%), premium milling recovery (71.4%) and grade 1 (53.4%) headrice recovery, long (3.0mm) and slender(3.3mm) grain shape, high percentage acceptability in the cooked form as compared to the check variety, PSB Rc82 and IR64, the eating quality check.
Intermediate amylose content (17.85%), premium milling recovery (71.1%), fair brown rice (77.3%) and grade 2 (47.2%) headrice recovery, long (6.9mm) and slender (3.2) grain shape.
Premium milling recovery (70.8%), fair brown rice (78.8%), long (7.1mm) and slender (3.3mm) grain shape, high percentage acceptability in the raw form as compared to IR64, the eating quality check.
Intermediate amylose content (22.0%), premium milling recovery (71.3%),and grade 1 (54.0%) headrice recovery , long (7.1mm) and slender (3.2) grain shape. High percentage acceptability in the raw form as compared to IR64, the eating quality check.
Intermediate amylose content (19.1%), premium milling recovery(71.0%),fair brown rice (78.9%) and grade 1 (50.2%) headrice recovery, long (7.1mm) and slender (3.5mm) grain shape.

Premium milling recovery (72.5%), fair brown rice (79.6%) and grade 1 (51.1%) headrice recovery. Long (6.9mm) and slender (3.1mm) grain shape.
Intermediate amylose content (19.1%), premium milling recovery(71.0%),fair brown rice (78.4%) and grade 1 (50.4%) headrice recovery. High percentage acceptability in the raw form as compared to IR64, the eating quality check.
Premium milling recovery (72.0%), fair brown rice (78.6%) and grade 1 (48.7%) headrice recovery. Long (7.1mm) and slender (3.1mm) grain shape.
Premium milling recovery (70.9%) and headrice (59.5%) recovery,long (6.9mm) and slender (3.1mm) grain shape.High percentage acceptability in the raw form as compared to IR64, the eating quality check. Slightly creamish when cooked.
Intermediate amylose content (20.1%), premium milling recovery(72.2%),fair brown rice (78.5%) and grade 1 (54.5%) headrice recovery. Long (6.9mm) and slender (3.1mm) grain shape. High percentage acceptability in the raw form as compared to the check variety, Mestiso 1 and IR64, the eating quality check.
Intermediate amylose content (18.3%), premium milling recovery(72.1%),fair brown rice (78.7%) and headrice(58.1%) recovery. Long (7.2mm) and slender (3.3mm) grain shape. High percentage acceptability in cooked and raw forms as compared to IR64, the eating quality check.
Intermediate amylose content (18.3%), premium milling recovery(70.8%),fair brown rice (78.7%) and grade 1 (54.5%) headrice recovery. Long (7.1mm) and slender (3.2mm) grain shape.
Intermediate amylose content (20.2%), premium milling recovery(72.2%),fair brown rice (79.0%) and grade 2 (39.4%) headrice recovery. Comparable to IR64, the eating quality check in terms of cooked form.
Intermediate amylose content (19.1%), premium milling recovery(71.7%),fair brown rice (78.9%) and grade 2 (39.9%) headrice recovery. Long (6.9mm) and slender (3.2mm) grain shape.

Premium milling recovery (70.3%), fair brown rice (77.5%) and grade 1 (56.1%) headrice recovery. Long (7.1mm) and slender (3.1mm) grain shape.High percentage acceptability in both in cooked and row forms as compared to IR64, the eating quality check.
Premium milling recovery (70.7%), fair brown rice (78.3%) and grade 2 (42.1%) headrice recovery. Medium (6.2mm) and intermediate (2.6mm) grain shape.
Premium milling recovery (74.5%) and headrice (59.6%) recovery. Medium (5.9mm) and intermediate (2.6mm) grain shape.
Grade 1 (69.8%) milling and headrice (54.6%) recovery, fair brown rice (77.2%), medium (6.1mm) and intermediate (2.9mm) grain shape. High percentage acceptability both in cooked and raw forms as compared to the check varieties, NSIC Rc192 and PSB Rc14.
Intermediate amylose content (19.1%), premium milling recovery (72.3%), fair brown rice (79.1%) and grade 1 (50.1%) headrice recovery.
Intermediate amylose content (20.1%), premium milling recovery(71.1%),fair brown rice (78.4%) and grade 2 (49.8%) headrice recovery. Long (7.3mm) and slender (3.5mm) grain shape.High percentage acceptability in both in the row form as compared to the checked varieties, NSIC Rc192 and PSB Rc14.
Premium milling recovery (72.8%), fair brown rice (78.9%) and grade 1 (52.2%) headrice recovery. Long (6.6mm) and slender (3.3mm) grain shape.
Premium milling (71.7%), and headrice (64.7%) recovery. Fair brown rice (79.1%), Short (5.1 mm) and bold (1.8) grain shape.Percent acceptability comparable to Koshihikari with better sensory qualities in terms of glossiness, cohesiveness, tenderness, smoothness and taste.
Premium (73.0%) milling and head rice (64.2%) recovery. Acceptable both in cooked and raw forms.Comparable to Koshihikari, the eating quality check.

Grade 1 (69.8%) milling and headrice (55.6%) recovery. Long (6.88 mm) and intermediate (2.9) grain shape. Higher percentage acceptability in the raw form as compared to IMS 2. Better sensory qualities in terms of glossiness, cohesiveness, tenderness and smoothness, very opaque.
Premium (71.1%) milling and headrice (62.1%) recovery. Fair (77.4%) brown rice. Long (6.8 mm) and slender (3.1) grain. Slightly creamish when cooked and in raw form, cooked rice is glossy, cohesive, tender, smooth better than M19.
Premium (70.7%) milling recovery. Fair (78.5%) brown rice. Long (7.0 mm) and slender (3.2) grain. Higher percentage acceptability both in cooked and raw forms as compared to IR64, the eating quality check. Slightly aromatic, tastier and better transluscency than M1.
Premium (70.7%) milling recovery. Grade 1 (50.5%) headrice recovery. Long (7.1 mm) and slender (3.4) grain. Higher percentage acceptability especially in the cooked form as compared to IR64, the eating quality check and Mestizo 1.
Intermediate (20.5%) amylose content. Premium (71.6%) milling recovery and Grade 1 (50.6%) headrice recovery. Medium (6.4 mm) and intermediate (2.9) grains. Less chalky, comparable sensory qualities as M1.
Intermediate (20.9%) amylose content. Grade 1 (69.5%) milling and headrice (50.3%) recovery. Long (6.9 mm) and slender (3.3) grain. Higher percentage acceptability both in cooked and raw forms as compared to IR64, the eating quality check, comparable to Mestizo 1. Less chalky with comparable sensory qualities.
Premium (70.6%) milling recovery. Grade 1 (48.5%) headrice recovery. Extra long (7.5 mm) and slender (3.5) grain. More acceptable in raw form, transluscent than IR64 and Mestizo 1.

Premium (71.2%) milling recovery. Grade 1 (50.6%) headrice recovery. Fair (79.0%) brown rice. Long (6.7 mm) and slender (3.0) grain. Higher percentage acceptability in the raw form as compared to IR64. Comparable sensory qualities with IR64 and Mestiso 19.
Premium (70.9%) milling recovery. Fair (78.4%) brown rice. Long (7.2 mm) and slender (3.2) grain. Higher percentage acceptability in the raw form as compared to IR64, the eating quality check. High amylose but low/intermediate gelatinization temperature.
Premium (70.8%) milling recovery. Fair (77.9%) brown rice. Long (6.7 mm) and slender (2.8) grain width. Higher percentage acceptability both in cooked and raw forms as compared to IR64, the eating quality check. Slightly aromatic when cooked, better glossiness in cooked and raw forms.
Intermediate (21.4%) amylose content. Premium (70.5%) milling recovery and fair (78.3%) brown rice. Long (7.2 mm) and slender (3.4) grain. Better acceptability in the raw form than IR64.
Intermediate (17.4%) amylose content. Premium (73.2%) milling and grade 1 (52.0%) headrice recovery. Long (6.7mm) and intermediate (3.0) grain shape.
Intermediate (19.1%) amylose content. Premium (70.3%) milling and grade 1 (49.5%) headrice recovery.Fair (78.6%) brown rice. Long (6.9mm) and slender (3.3) grain shape.
Intermediate (20.0%) amylose content. Premium (72.6%) milling recovery. Fair (79.8%) brown rice. Long (6.8mm) and slender (3.1) grain shape.Slightly creamish in the raw form.
Premium (71.1%) milling recovery and fair (79.5%) brown rice. Long (6.7mm) and slender (3.1) grain shape. Higher percentage acceptability in the cooked form as compared to IR64, the eating quality check. Slightly creamish when cooked.

Premium (70.1%) milling and grade 1 (56.5%) headrice recovery. Fair (78.4%) brown rice. Long (6.7mm) and slender (3.4) grain shape. Slightly creamish in the raw form. High (23.2%) amylose content and low gelatinization temperature.
Intermediate (20.&%) amylose content. Grade 1 (69.3%) milling and headrice (53.5%) recovery. Long (6.8mm) and slender (3.2) grain shape. Higher percentage acceptability in the raw for as compared to IR64.
Intermediate (21.5%) amylose content. Premium (70.1%) milling recovery. Grade 1 (51.2%) headrice recovery; fair (77.7%) brown rice. Glossy, slightly cohesive, slightly tender, slightly smooth.
Intermediate (20.2%) amylose content. Grade 1 (68.3) milling recovery, fair (77.3%) brown rice. Extra long (7.5mm) and slender (3.3) grain shape. Higher percentage acceptability both in cooked and raw forms as compared to IR64. Slightly glossy, cohesive, tender and smooth.
Premium (71.2%) milling and grade 1 (55.3%) headrice recovery. Long (6.7mm) and intermediate (2.9) grain shape. High amylose (23.6%) but with intermediate/low gelatinization temperature. Very white, slightly glossy, slightly cohesive, slightly tender and smooth.
Intermediate (20.6%) amylose content. Grade 1 (70.0%) milling recovery; fair (77.3%) brown rice. Long (7.0mm) and slender (3.3) grain shape. Percentage acceptability comparable to IR64.
Intermidiate (21.0%) amylose content. Grade 1 (69.4) milling recovery; fair (76.6%) brown rice. Long (7.1mm) and intermediate (3.0) grain shape. Slightly glossy, slightly cohesive, slightly tender and slightly smooth.
Premium (70.2%) milling and grade 1 (54.0%) headrice recovery. Fair (77.4%) brown rice. Long (6.5mm) and slender (3.1) grain shape. Percentage acceptability comparable to IR64. Glossy, cohesive and slender, better than PSB Rc90.

Intermediate (21.4%) amylose content. Grade 1 (70.0%) milling recovery; fair (77.5%) brown rice. Medium (6.5mm) grain length and slender (3.2) grain shape. Higher percentage acceptability in the raw form as compared to IR64. Slightly creamish, slightly glossy, slightly tender when cooked.
Premium (70.9%) milling recovery. Grade 1 (51.2%) headrice recovery; fair (77.5%) brown rice. Medium (6.4mm) grain length and slender (3.3) grain shape. High (22.2%) amylose content and low gelatinization temperature. Better sensory qualities than PSB Rc90.
Premium (71.1%) milling and headrice (63.7%) recovery. Fair (76.8%) brown rice. Medium (6.1mm) grain length and intermediate (3.0) grain shape. Higher percentage acceptability in cooked form and comparable to IR64 in the raw form. Sromatic, slightly tasty when cooked. Slightly creamish both in cooked and raw forms.
Intermediate (18.6%) amylose content. Fair (77.0%) brown rice. Grade 1 (68.8%) milling recovery. Medium (6.1mm) grain length and intermediate (2.8) grain shape. Percentage acceptability comparable to IR64. Better sensory qualities in cooked form, cohesive, tender and smooth.
Grade 1 (69.6%) milling and headrice (52.4%) recovery. Long (6.9mm) and slender (3.1) grain shape. Higher percentage acceptability both in cooked and raw forms. White, cohesive and tender when cooked.
Grade 1 (69.5%) milling recovery. Fair (77.2%) brown rice. Long (6.9 mm) and intermediate (2.9) grain shape. Higher percentage acceptability in the raw form asa compared to IR64, the eating quality check. Sensory qualities comparable to Mestiso 20.
Grade 1 (68.3%) milling recovey. Fair (76.6%) brown rice. Medium (6.4 mm) and intermediate (2.8) grain shape.

Premium (70.1%) milling recovery. Fair (77.4%) brown rice. Long (7.4 mm) and dlender (3.4) grain shape. Higher percentage acceptability in the raw fform sa compared to IR64, the eating quality check.
Grade 1 (69.1%) milling and headrice (49.6%) recovery. Fair (77.2%) brown rice. Long (7.1 mm) and slender (3.2) grain shape.
Premium (70.6%) milling recovery. Fair (77.2%) brown rice. Medium (6.5 mm) grain length and intermediate (2.8) grain shape. Higher percentage acceptability in the cooked form as compared to IR64, the eating quality check. Slightly creamish when cooked.
Premium (70.7%) milling and headrice (58.3) recovery. Fair (77.7%) brown rice. Long (6.9mm) and slender (3.3) grain shape. Slightly creamish both in cooked and raw forms.
Intermediate (18.6%) amylose content. Fair (77.8%) brown rice. Premium (70.8%) milling recovery. Long (7.1mm) and slender (3.2) grain shape. High percentage acceptability in the raw form. Slightly creamish in the raw form.
Grade 1 (68.5%) milling and headrice (52.4%) recovery. Fair (75.6%) brown rice. Short (5.4 mm) and intermediate (2.4) grain shape. Higher percentage acceptability in the cooked form as compared to IR64, the eating quality check. Slightly creamish when cooked, creamish in the raw form.
Premium (70.1%) milling and Grade 1 (51.1%) headrice recovery. Fair (77.1%) brown rice. Medium (5.7 mm) and intermediate (2.5) grain shape. Higher percentage acceptability in the cooked form as compared to PSB Rc90. Slightly creamish both in cooked and raw form.

Intermediate (19.2%) amylose content. Premium (57.1%) headrice and grade 1 (70.0%) milling recovery. Fair (77.7%) brown rice. Medium (6.5 mm) and slender (3.1) grain shape. Better sensory qualities than PSB Rc90.
Yield advantage of 28.5% over the check variety, NSIC Rc104. Yield range: 2307-6597kg/ha. Earlier maturity at 143 days than NSIC Rc104. Intermediate reaction to blast. Premium (73.2%) milling recovery and grade 2 (44.4%) headrice recovery. Short (5.0mm) and bold (1.8) grains. Higher percentage acceptability in the raw form than NSIC Rc104. Slightly creamish and slightly tasty when cooked.
Higher yield potential of 4.3t/ha under moderate cold stress. Yield advantage of 37.9% over the check variety, NSIC Rc104. Yield range : 893-6895kg/ha. Earlier maturity days of 143 than NSIC Rc104. Intermediate reaction to blast, green leafhopper and brown planthopper. Premium (72.1%) milling recovery. Grade 2 (47.2%) headricE and brown rice (80.5%). Short (4.8mm) and bold (1.6) grains. Comparable to NSIC Rc104 in terms of cooked and raw forms. Moderately creamish both in cooked and raw forms.
Higher yield potential of 3.8t/ha. With yield advantage of 23.5% over the check variety, NSIC Rc104. Yield range :3026-4961kg/ha. Earlier maturity of 143 days than NSIC Rc104. Intermediate reaction to blast. Premium (71.7%) milling and grade 1 (55.8%) headrice recovery. Short (4.7mm) and bold (1.6) grains. Higher percentage acceptability in the raw form than NSIC Rc104. Moderately creamish both in cooked and raw forms.
Yield advantage of 29.5% over the check variety, PSB Rc96. Yield range: 1382-4068kg/ha. Comparable to PSB Rc96 in terms of maturity days (145). Resistant reaction to Whiteheads. Intermediate reaction to blast and yellow stemborer. Premium (73.0%) milling and headrice (64.0%) recovery. 100% acceptability in the cooked form and 90% in the raw form. Slightly tasty when cooked.
Yield advantage of 31.5% over the check variety, PSB Rc 14. Yield range: 1526-5267kg/ha. Early maturing at 114 days. Intermediate reaction to blast, bacterial leaf blight and sheath blight. Resistant reaction to deadhearts and whiteheads (YSB). Intermediate reaction to whiteheads (WSB), brown planthopper and green leafhopper. Fair (77.8%) brown rice. Grade 1 (69.2%) milling recovery. Medium (6.3 mm) and intermediate (2.7) grain shape. Slightly glossy, slighly cohesive, slightly tender and slightly smooth in terms of sensory qualities.

Yield advantage of 18.4% over the check variety, PSB Rc 14. Yield range: 1576-4493kg/ha. Early maturing at 115 days. Intermediate reaction to blast, bacterial leaf blight, sheath blight and rice tungro virus (modified). Resistant reaction to deadhearts and whiteheads (YSB). Intermediate reaction to brown planthopper and green leafhopper. Intermediate (21.2%) amylose content. Premium (73.9%) milling recovery. Fair (79.7%) brown rice and grade 1 (51.5%) headrice recovery. Long (7.3mm) and slender (3.3) grain shape. Higher percentage acceptability in the raw form as compared to IR64. Slightly creamish, slightly glossy in the raw form in terms of sensory qualities.
Yield advantage of 42.9% over the check variety, NSIC Rc 192. Yield range : 1397-4531kg/ha. Early maturing at 109 days. Intermediate reaction to blast, bacterial leaf blight and rice tungro virus (modified). Resistant reaction to deadhearts and whiteheads. Intermediate reaction to green leafhopper and brown planthopper. Intermediate (20.9%) amylose content. Premium (72.2%) milling recovery. Fair (78.9%) brown rice and grade 1 (53.4%) headrice recovery. Long (6.7mm) and slender (2.9) gran shape. High percentage acceptability in the cooked form as compared to IR64.
Yield advantage of 18.0% over the check variety, PSB Rc 14. Yield range : 1817-3577kg/ha. Early maturing at 113 days. Intermediate reaction to blast, bacterial leaf blight and sheath blight. Resistant reaction to deadhearts. Intermediate reaction to whiteheads. Intermediate reaction to brown planthopper and green leafhopper. Intermediate (19.2%) amylose content. Premium (72.5%0 milling recovery. Fair (79.4%) brown rice and grade 1 (53.6%) headrice recovery. Long (7.1 mm) and slender (3.3) grain shape. 100% acceptability in the cooked form as compared to IR64.
Yield advantage of 19.8% over the check variety, PSB Rc 14. Yield range: 1481-4334kg/ha. Early maturing at 113 days. Intermediate reaction to blast, bacterial leaf blight and sheath blight. Resistant reaction to deadheart and whiteheads (YSB). Intermediate reaction to brown planthopper and green leafhopper. Intermediate (19.5%) amylose content. Premium (72.5%) milling recovery. Fair (79.3%) brown rice. Long (6.9mm) and slender (3.1) grain shape. Sl. glossy, slightly cohesive, slightly smooth and slightly tender in terms of sensory qualities.
Yield advantage of 18.9% over the check variety, PSB Rc 14. Yield range: 1528-4361kg/ha. Early maturing at 113 days. Intermediate reaction to blast and bacterial leaf blight. Resistant reaction to deadhearts and whiteheads (YSB). Intermediate reaction to brown planthopper and green leafhopper. Intermediate (19.7%) amylose content. Premium (72.6%) milling recovery; fair (79.7%) brown rice. Long (6.7 mm) and intermediate (2.9) grain shape. High percentage acceptability in the raw form as compared to IR64. Sligthly creamish when cooked.

Yield advantage of 9.8% and 9.9% in Bohol APC in 2017WS and 2018DS, respectively, over the check variety, NSIC Rc 222. Yield advantage of 9.0% in WESVIARC in 2018WS. With yield advantage of 12.7% in CMU, 18.3% in PhilRice Agusan and 9.9% in ZAMPIARC during the 2018WS. Yield range : Visayas= 3062-7161kg/ha; Mindanao=2839-7904kg/ha. Early maturing at 111 days. Intermediate reaction to blast, bacterial leaf blight, sheath blight and rice tungro virus (modified). Resistant reaction to deadhearts. Intermediate reaction to whiteheads, brown planthopper and green leafhopper. Intermediate (18.2%) amylose content. Grade 1 (68.8%) milling and headrice (49.2) recovery. Fair (77.9%) brown rice. Long (6.6 mm) and slender (3.1) grain shape. High percentage acceptability both in cooked and raw forms as compared to IR64.
Yield advantage of 13.1% in 2017WS and 29.0% in 2018DS in Bohol APC, over the check variety, NSIC Rc 222. 39.1% yield advantage during the 2017WS in UEP. Yield range : 3054-7342kg/ha. Early maturing at 109 days. Intermediate reaction to blast, bacterial leaf blight, sheath blight and rice tungro virus (modified). Resistant reaction to deadheart. Intermediate reaction to whiteheads, brown planthopper and green leafhopper. Low (17.2%) amylose content. Grade 1 (68.7%) milling recovery. Fair (76.2%) brown rice. Long (6.7mm) and slender (3.1) grain shape. High percentage acceptability both in cooked and raw forms as compared to IR64. Slightly creamish both in cooked and raw forms in terms of sensory qualities.
Yield advantage of 13.8% across season, over the check variety, MS 11. Early maturing at 112 days. Yield range : 1545-6221kg/ha. Intermediate reaction to blast and bacterial leaf blight. Resistant reaction to deadheart. Intermediate reaction to whiteheads and brown planthopper. Low (12.1%) amylose content. Grade 1 (69.7%) milling recovery. Fair (78.1%) brown rice. Premium (58.8%) headrice recovery. Creamish both in cooked and raw forms in terms of sensory qualities.
With average yield advantage of 15.5% during the 2017DS in PhilRice CES, SCRC, PREC and PhilRice Bicol, over the check variety, Mestiso 19. Average yield advantage of 8.8% during the 2017WS in DA-Tabuk, PhilRice Bicol and INREC. High yield advantage of 24.6%, 41.6% and 48.6% in SCRC, PhilRice Isabela and INREC, respectively, during the 2018WS. Average yield advantage of 30.2% was obtained in Agusan del Sur and Davao in 2018DS and 25.6% in 2018WS in CMU, PhilRice Midsayap, Agusan. Yield range : Luzon= 4233-9964kg/ha; Mindanao = 2605-11264kg/ha. Early maturing at 112 days. Intermediate reaction to blast, bacterial leaf blight, sheath blight and rice tungro virus (modified). Resistant reaction to Deadheart. Intermediate reaction to whitehead, brown planthopper and green leafhopper. Low (14.4%) amylose content. Grade 1 (69.3%) milling and headrice (49.2) recovery. Fair (77.3%) brown rice. Long (7.2mm) and slender (3.5) grain shape. High percentage acceptability both in cooked and raw forms as compared to IR64. Slightly creamish in the raw form in terms of sensory qualities.

Yield advantage of 5.1% across season over the check variety, Mestiso 20. Yield range : 2320-12678kg/ha. Early maturing at 115 days. Intermediate reaction to blast, bacterial leaf blight, sheath blight and rice tungro virus (modified). Resistant reaction to deadheart. Intermediate reaction to whiteheads, brown planthopper and green leafhopper. Intermediate (16.5%) amylose content. Premium (70.6%) milling recovery. Fair (78.2%) brown rice and grade 1 (49.0) headrice recovery. Long (6.8mm) and slender (3.4) grain shape. High percentage acceptability in the raw form as compared to IR64. Slightly creamish in the raw form in terms of sensory qualities.
Yield advantage of 12.4% over the check variety, NSIC Rc 194, Resistant reaction to deadheart and whitehead. Intermediate reaction to blast, bacterial leaf blight, sheath blight, rice tungro virus (modified), brown planthopper and green leafhopper. Low (16.2%) amylose content. Grade 1 (66.7%) milling and headrice (56.6%) recovery. Long (6.7mm) and slender (3.1) grain shape. High percentage acceptability in the cooked (93.3%) and raw (100%) forms.
With yield advantage of 11.5% over the check variety, NSIC Rc 192.Very early maturing at 105 days. Resistant reaction to Deadheart (YSB) and Green leafhopper. Intermediate reaction to blast, bacterial leaf blight and rice tungro virus (modified). Intermediate reaction to Whiteheads and Brown planthopper. Intermediate (18.1%) amylose content. Premium (71.8%) milling recovery. Medium (6.5 mm) and intermediate (2.8) grain shape.
Yield advantage of 7.1% over the check variety, NSIC Rc 192. Early maturing at 109 days. Resistant reaction to Deadheart and Whitehead (YSB). Intermediate reaction to blast, bacterial leaf blight, brown planthopper and green leafhopper. Intermediate (17.2%) amylose content. Grade 1 (70.6%) milling recovery. Long (6.6mm) and intermediate (2.8) grain shape.
Yield advantage of 6.7% over the check variety, NSIC Rc 192. Early maturing at 113 days. Resistant reaction to Deadheart and Whitehead (YSB). Intermediate reaction to blast, bacterial leaf blight, sheath blight and rice tungro virus (modified). Intermediate reaction to brown planthopper and green leafhopper. Premium (71.2%) milling recovery. Grade 1 (55.3%) headrice recovery. Medium (6.5mm) and intermediate (2.9) grain shape.
With yield advantage of 10.1% over the check variety, NSIC Rc 192. Medium maturing at 118 days. Resistant reaction to blast, bacterial leaf blight and sheath blight. Intermediate reaction to Whitehead (WSB), brown planthopper and green leafhopper.Intermediate (20.3%) amylose content.Grade 1 (69.3%) milling and headrice (50.8%) recovery. Long (6.6mm) and intermediate (2.9) grain shape. 93.3% acceptability in the raw form.
With yield advantage of 57.2% over the check variety, N22.Early maturing at 110 days. Average spikelet fertility of 82%. Resistant reaction to Deadheart (YSB) and Whitehead (WSB and YSB). Intermediate reaction to blast, bacterial leaf blight, sheath blight, brown planthopper and green leafhopper. Intermediate (20.1%) amylose content. Grade 1 (66.9%) milling recovery. Long (6.8mm) and slender (3.3) grain shape. With 93.3% acceptability in the raw form.

With yield advantage of 54.1% over the check variety, N22. Early maturing at 109 days. Average spikelet fertility of 85%. Resistant reaction to Whitehead (YSB and WSB). Intermediate reaction to blast, bacterial leaf blight, sheath blight, deadheart (YSB),brown planthopper and green leafhopper. Intermediate (19.8%) amylose content. Grade 1 (65.15) milling recovery. Long (6.7 mm) and slender (3.2) grain shape. High percentage acceptability both in cooked (96.7%) and raw (90.0%) forms. Slightly creamish when cooked and creamish in raw form in terms of sensory qualities.
Yield advantage of 23.1% and 68.6% across seasons over the check varieties, NSIC Rc 182 and NSIC Rc 184. High yield advantage of 80.7% in the WS and 41.9% in the DS over NSIC Rc 184. Yield advantage of 42.1% in the DS over NSIC Rc 182. Very early maturing at 104 days. Resistant reaction to blast and whitehead (YSB). Intermediate reaction to bacterial leaf blight, sheath blight, brown planthopper and green leafhopper. Intermediate (19.6%) amylose content. Premium (71.4%) milling recovery. Grade 1 (51.4%) headrice recovery. Long (6.6 mm) and intermediate (2.8) grain shape. High percentage acceptability (93.3%) in the raw form.
Yield advantage of 16.6% and 28.2% across seasons over the check varieties, NSIC Rc 392 and PSB Rc 90. Early maturing at 109 days. Resistant reaction to blast and deadheart (YSB). Intermediate reaction to bacterial leaf blight, sheath blight, whitehead (YSB), brown planthopper and green leafhopper. Low (13.8%) amylose content. Grade 1 (69.4%) milling and headrice (56.9%) recovery. Long (7.1 mm) and slender (3.2) grain shape. High percentage acceptability both in cooked (95%) and raw (98.3%) forms. Slightly creamish both in cooked and raw forms in terms of sensory qualities.
Yield advantage of 16.7 across season over NSIC Rc 392 and 8.2% over PSB Rc 88. With 39.3% yield advantage during the DS over NSIC Rc 392. Early maturing at 110 days. Resistant reaction to deadheart (YSB). Intermediate reaction to blast, bacterial leaf blight, whitehead (YSB), brown planthopper and green leafhopper. Low (16.6%) amylose content. Grade 1 (68.7%) milling recovery. Long (6.9mm) and slender (3.1) grain shape. High percentage acceptability both in cooked and raw forms.
Yield advantage of 51.1% across season over the check variety, NSIC Rc 184 and 14.5% over NSIC Rc 392. Early maturing at 107 days. Resistant reaction to deadheart and whitehead (YSB). Intermediate reaction to blast, bacterial leaf blight, brown planthopper and green leafhopper. Intermediate (17.8%) amylose content. Grade 1 (67.2%) milling recovery. Long (7.3 mm) and slender (3.3) grain shape. Acceptable both in cooked and raw forms.

Yield advantage of 26.6% and 39.2% across seasons over the check varieties, NSIC Rc 392 and PSB Rc 90. Early maturing at 109 days. Resistant reaction to blast and deadheart (YSB). Intermediate reaction to bacterial leaf blight, whitehead (YSB) brown planthopper and green leafhopper. Intermediate (17.2%) amylose content. Grade 1 (67.7%) milling recovery. Medium (5.5 mm) and intermediate (2.2) grain shape.
High yield advantage of 61.9% in PhilRice Isabela and 58.1% in INREC in 2018WS over the check variety, Mestiso19. Yield advantage of 41.3% in 2019DS in SCRC over the check variety, Mestiso 20. Average yield advantage of 6.9% across sites in 2019WS over the check variety, Mestiso 20. Early maturing hybrid at 112 days. Resistant reaction to blast, rice tungro virus (modified) and deadheart (YSB). Intermediate reaction to bacterial leaf blight, sheath blight, whitehead, brown planthopper and green leafhopper. Intermediate (20.9%) amylose content. Grade 1 (68.8%) milling and headrice (51.4) recovery. Long (6.7 mm) and slender (3.2) grain shape.93.3% acceptability in the raw form.
With yield advantage of 48.9% in PhilRice Isabela and 22.1% in Ilocos Norte in 2018WS over the check variety, Mestiso 19. Yield advantage of 12.5%, 18.7% and 37.3% in 2019DS over the check variety, Mestiso 20 in PhilRice CES, SCRC and DA-Tabuk, respectively. High yield advantage of 60.1% in 2019WS in PhilRice CES over the check variety, Mestiso 20. Early maturing hybrid at 108 days. Resistant reaction to blast, Deadheart (YSB) and Whitehead (WSB/YSB). Intermediate reaction to bacterial leaf blight, sheath blight, brown planthopper and green leafhopper. Intermediate (21.9%) amylose content. Premium (70.1%) milling and headrice (57.3%) recovery. Long (6.7mm) and slender (3.2) grain shape. 100% acceptability in the raw form.
With an average yield advantage of 17.3% across sites in 2018WS over the check variety, Mestiso 20. High yield advantage of 52.7% in 2019WS in PhilRice CES over Mestiso 20. Yield advantage of 14.6%, 11.6% and 19.9% in 2019WS over Mestiso 20 in Pampanga,PhilRice Bicol and DA-Tabuk, respectively. Early maturing hybrid at 115 days. Resistant reaction to blast, deadheart and whitehead (YSB). Intermediate reaction to bacterial leaf blight, sheath blight, rice tungro virus (modified),brown planthopper and green leafhopper. Low (12.2%) amylose content. Premium (74.0%) milling recovery. Grade 1 (54.1%) headrice recovery. Medium (6.3 mm) and intermediate (2.9) grain shape. 90% acceptability in the cooked for. Slighly creamish in the raw form in terms of sensory qualities.
Intermediate amylose (19.1%) content. Premium (72.0%) milling recovery. Early maturing at 113 days.

Intermediate reaction to white stemborer, yellow stemborer, brown planthopper and green leafhopper. Intermediate (19.7%) amylose content. Grade 1 (65.6%) milling and headrice (50.6%) recovery. Long (6.8mm) and intermediate (2.9) grain shape. High percentage acceptability both in cooked and raw forms.
Grade 1 (68.2%) milling and headrice (53.6%) recovery. Medium (6.0mm) and intermediate (2.6) grain shape. High percentage acceptability both in cooked and raw forms.
Intermediate (18.6%)amylose content. Extra long(7.7mm) and slender (3.4) grain shape. High percentage acceptability in cooked form.
Intermediate(18.0%) amylose content. Grade 1 (65.6%) milling and headrice (50.8%) recovery. Long (7.2mm) and slender (3.3) grain shape. High percentage acceptability both in cooked and raw forms.
Low (17.0%) amylose content. Grade 1 (66.4%) milling recovery. Long (7.1mm) and slender (3.1) grain shape.
Grade 1 (69.9%) milling and headrice (54.1%) recovery. Medium (6.3mm) and intermediate (2.7) grain shape. High percentage acceptability both in cooked and raw forms. Slightly creamish in cooked and raw forms.
Intermediate (19.4%) amylose content. Grade 1 (66.4%) milling recovery. Medium (6.0mm) and intermediate (2.3) grain shape.
Low (12.4%) amylose content. Grade1 (66.5%) milling recovery. Medium (5.7mm) and intermediate (2.4) grain shape.
Low (12.9%) amylose content. Grade 2 (68.2%) milling recovery. Premium (57.4%) headrice recovery. Long (6.9mm) and slender (3.4) grain shape. Slightly tasty when cooked.
Low (13.4%) amylose content. Grade 1 (65.7%) milling and headrice (53.7%) recovery. Long (6.8mm) and slender (3.1) grain shape. Slightly tasty when cooked.
Low (12.9%) amylose content. Grade 1 (67.0%) milling and headrice (54.2%) recovery. Long (6.8mm) and slender (3.2) grain shape. High percentage acceptability both in cooked and raw forms.

Premium (71.1%) milling and headrice (58.0%) recovery. Medium (6.5mm) and intermediate (2.9) grain shape.
Low(11.9%) amylose content. Grade 1 (51.0%) headrice recovery. Medium (6.1mm) and intermediate (2.8) grain shape. Slightly tasty when cooked.
Intermediate (19.5%) amylose content. Grade 1 (66.8%) milling and headrice (49.8%) recovery. Long (6.8mm) and slender (3.5) grain shape.
Intermediate (20.4%) amylose content. Grade 1 (68.9%) headrice recovery. Long (6.9mm) and slender (3.5) grain shape.
Premium (71.8%) milling and headrice (57.5%) recovery. Long (6.7mm) and slender (3.2) grain shape.
Premium (70.8%) milling recovery. Fair (77.2%) brown rice. Long (68 mm) and intermediate (3.0) grain shape. Slightly creamish when cooked.
Intermediate (20.1%) amylose content. Premium (75.0%) milling and headrice (60.4%) recovery. Long (6.7mm) and slender (3.1) grain shape. Slightly creamish when cooked.
Intermediate (21.1%) amylose content. Premium (73.9%) milling recovery. Grade 1 (55.1%) headrice recovery. Long (6.8mm) and slender (3.3) grain shape.
Intermediate (19.0%) amylose content. Grade 1 (69.3%) milling recovery. Premium (73.2%) headrice recovery. Long (6.7mm

Intermediate (18.6%) amylose content. Premium (72.8%) milling recovery. Long (6.7mm) and intermediate (2.9) grain shape. Slightly creamish when cooked.
Extra long (7.8mm) and slender (3.5) grain shape. Resistant reaction to blast, rice tungro virus (modified), yellow stemborer and white stemborer. Intermediate reaction to bacterial leaf blight, sheath blight, brown planthopper and green leafhopper.
Low (14.8%) amylose content. Premium (71.0%)milling and headrice (57.1%)recovery.Long (6.9mm) and slender (3.1) grain shape.
Intermediate (20.1%) amylose content. Premium (71.1%) milling recovery. Long (7.4mm) and slender (3.6) grain shape.
Intermediate (20.5%) amylose content. Premium (70.6%) milling recovery and Grade 1 (56.6%) headrice recovery. Medium (6.2 mm) and intermediate (2.8) grain size and shape. Slightly creamish when cooked and creamish in raw form.
High (23.1%) amylose content. Premium (71.7%) milling recovery and Grade 1 (53.3%) headrice recovery. Long (7.1mm) and slender (3.4mm) grain size and shape.
Intermediate (18.5%) amylose content. Premium (71.5%) milling recovery and Grade 1 (59.7%) recovery. Extra Long (7.7mm) and slender (3.6mm) grain size and shape.
Hard (22.1%) amylose content. Grade 1 (69.5%) milling recovery and headrice (50.6%) recoveries. Long (7 mm) and slender (3.4mm) grain size and shape. Slightly creamish in the raw form
Intermediate (19.0%) amylose content. Premium (70.9%) milling recovery and Grade 1 (50.2%) headrice recovery. Long (7.0mm) and slender (3.2mm) grain size and shape.
Intermediate (18.4%) amylose content. Premium (70.1%) milling recovery and Grade 1 (53.9%) headrice recovery. Long (6.8mm) and slender (3.3mm) grain size and shape.
Intermediate (18.4%) amylose content. Fair (79%) brown rice. Premium milling (73.5%) and headrice (63.1%) recovery. Medium (6.4mm) and intermediate (3.0) grain size and shape. Soft Instron cooked rice hardness (1.52 kg/cm2). Total carotenoids of 4.6 (3.8-6 ug/g). Essentially-derived line of PSB Rc 82 with 98.27% genetic similarity.

Intermediate amylose content (17.5%). Fair brown rice (78.4%). Premium milling recovery (70.8%). Grade 1 head rice recovery (48.0%). Long (7.3mm) and slender grain size and shape (3.4). Slightly creamish in the raw form. Zinc content of 23.2 (mg kg-1) and iron content of 3.2 (mg kg-1).
Hard amylose content (24.1%). Premium milling recovery (71.1%). Fair brown rice (78.0%). Long (6.6mm) and intermediate grain shape (2.9).
Intermediate amylose content (22.0%). Fair brown rice (79.1%). Grade 1 milling (69.0%) and headrice recoveries (53.5%). Medium (6.1mm) and intermediate grain shape (2.8). 100% acceptability in the cooked form. Slightly creamish in the raw form.
Intermediate amylose content (20.1%). Fair brown rice (79.2%). Premium milling (70.9%) and headrice recoveries (60.1%). Long (6.7mm) and slender grain shape (3.2). 93.3% acceptability in the raw form. Slightly creamish when cooked.
Low amylose content (15.2%). Fair brown rice (79.2%). Premium milling (70.2%) and headrice recoveries (61.3%). Long (6.8mm) and intermediate grain shape (2.9). 90% acceptability in the raw form.
Low amylose content (15.2%). Fair brown rice (79.2%). Premium milling (70.2%) and headrice recoveries (61.3%). Long (6.8mm) and intermediate grain shape (2.9). 90% acceptability in the raw form.
Low amylose content (13.5%). Fair brown rice (78.9%). Premium milling (71.5%) and headrice recoveries (63.2%). Long (6.7mm) and slender grain shape (3.1).
Low amylose content (14.3%). Good brown rice (80.2%). Premium milling (71.7%) and headrice recoveries (64.9%). Long (6.6mm) and slender grain shape (3.1).
Intermediate amylose content (19.3%). Fair brown rice (79.1%). Grade 1 milling (70.0%) and headrice recoveries (49.1%). Long (7.1mm) and slender grain shape (3.3). Slightly creamish when cooked.
Intermediate amylose content (20.7%). Fair brown rice (79.1%). Grade 1 milling recovery (68.9%). Long (7.4mm) and slender grain shape (3.5). 90% acceptability in the cooked form.

Intermediate amylose content (18.4%). Fair brown rice (79.6%). Premium milling recovery (70.8%). Grade 1 headrice recovery (51.5%). Long (7.1mm) and slender grain shape (3.4). 96.7% acceptability both in the cooked and raw forms. Slightly creamish in the raw form.
Intermediate amylose content (20.4%). Fair brown rice (77.0%). Grade 2 milling recovery (65.0%). Long (7.4mm) and slender grain shape (3.6.).
Intermediate amylose content (20.4%). Fair brown rice (78.9%). Grade 1 milling recovery (69.4%). Grade 2 headrice recovery (45.0%). Long (6.7mm) and intermediate grain shape (3.0). 90% acceptability in the cooked form.
Intermediate amylose content (17.0%). Fair brown rice (79.1%). Grade 1 milling recovery (69.5%). Grade 2 headrice recovery (47.2%). Long (6.9mm) and intermediate grain shape (2.9). Slightly creamish in the raw form.
Intermediate amylose content (20.9%). Fair brown rice (79.1%). Grade 1 milling (69.7%) and headrice recoveries (48.9%). Extra long (7.6 mm) and slender grain shape (3.5). 90% acceptability in the cooked form.
High amylose content (23.5%). Fair brown rice (79.4%). Premium milling recovery (71.0%). Grade 2 headrice recovery (47.9%). Long (7.0mm) and slender grain shape (3.2). 90% acceptability in the cooked and 96.7% in the raw forms.
High amylose content (23.5%); Fair brown rice (79.0%). Grade 1 milling recovery (69.2%). Long (7.0mm) and intermediate grain shape (2.9). 90.0% acceptability in the cooked and 96.7% in the raw forms
Intermediate amylose content (20.8%); Fair brown rice (78.7%). Grade 1 milling (69.3%) and headrice recoveries (53.7%). Extra long (7.7mm) and slender grain shape (4.0). Slightly creamish in the raw form.
Intermediate amylose content (21.9%); Fair brown rice (78.1%). • Grade 1 milling (69.0%) and headrice recoveries (51.2%). • Extra long (7.9mm) and slender grain shape (3.8). • 93.3% acceptability in the cooked and 96.7% in the raw forms.
Intermediate amylose content (19.0%); Fair brown rice (78.4%). Grade 1 milling recovery (67.7%). Long (7.1mm) and intermediate grain shape (3.0).
Low amylose content (13.7%). Fair brown rice (78.9%). Grade 1 milling (69.0%) and headrice recoveries (48.3%). Long (7.3 mm) and slender grain shape (3.5). 90% acceptability in the cooked and 93.3% in the raw forms. Slightly creamish when cooked.

Low amylose content (13.3%). Fair brown rice (79.1%). Grade 1 milling (69.9%) and headrice recoveries (52.0%). Long (7.1 mm) and slender grain shape (3.2). 96.7% acceptability in the cooked and 93.3% in the raw forms. Slightly creamish in the raw form.
Intermediate amylose content (18.5%). Fair brown rice (79.1%). Premium milling recovery (70.4%). Grade 1 head rice recovery (52.6%). Long (7.2mm) and slender grain shape (3.2). Slightly aromatic and slightly tasty when cooked.
High amylose content (23.2%). Fair brown rice (78.9%). Premium milling recovery (70.9%). Grade 1 head rice recovery (52.2%). Long (7.1mm) and intermediate grain shape (3.4). 100% acceptability in the raw form.
High amylose content (23.8%). Fair brown rice (79.4%). Premium milling recovery (72.5%). Grade 1 head rice recovery (56.0%). Grade 1 head rice recovery (56.0%). Long (6.8mm) and intermediate grain shape (3.0). 100% acceptability in the raw form.
Intermediate amylose content (21.3%). Fair brown rice (79.8%). Premium milling recovery (72.1%). Grade 1 head rice recovery (49.7%). Long (7.0mm) and slender grain shape (3.4). 90% acceptability in the raw form.
Intermediate amylose content (19.8-22.6%). Fair brown rice (77.6-79.7%). Premium milling recovery (71.8-73.7%). Grade 1 head rice recovery (44.1-59.5%). • Medium (6.4-6.6 mm) and slender grain shape (2.7-3.0). Slightly creamish when cooked and raw forms.
Intermediate amylose content (19.0-21.2%). Fair brown rice (77.0-79.0%). • Premium milling recovery (71.4-72.0%). • Grade 1 head rice recovery (44.4-57.6%). • Medium (6.4-6.6 mm) and slender grain shape (2.9-3.0). • Slightly creamish when cooked and raw forms.
• High amylose content (23.3%). • Fair brown rice (79.7-79.8%). • Premium milling recovery (71.2-71.6%). • Grade 2 head rice recovery (43.3-51.3%). • Long (7.1-7.2 mm) and intermediate grain shape (3.0-3.1).
Intermediate amylose content (20.0 - 21.0%). Fair brown rice (77.8 - 80.1%). Grade 1 milling recovery (66.7-72.3%). Grade 1 headrice recovery (48.3-56.8%). Medium (6.2-6.4mm) and slender grain shape (3.2). Slightly creamish in cooked and raw forms
Intermediate amylose content (20.7 - 21.9%). Good brown rice recovery (80.0-81.1%). Premium milling recovery (70.8-73.5%). Grade 1 headrice recovery (51.2-61.3%). Medium (6.2-6.5mm) and intermediate grain shape (2.9). Slightly creamish in cooked form
Intermediate amylose content (19.5-21.9%). Good brown rice recovery (79.8-80.8%). Premium milling recovery (69.6-72.2%). Grade 1 headrice recovery (52.9-54.3%). Long (6.7-6.8mm) and slender grain shape (3.3). Slightly creamish in cooked form.

Intermediate amylose content (19.9%). Fair brown rice recovery (79.1-79.4%). Premium milling recovery (69.8-72.7%) and headrice recovery (52.5-63.8%). Medium (6.4mm) and slender grain shape (3.2-3.3)
Intermediate amylose content (15.-21.9%). Fair brown rice recovery (79.1-79.5%). Premium milling recovery (71.0-71.8%) and headrice recovery (52.4-65.1%). Medium (6.5-6.6mm) and intermediate grain shape (3.0)
High amylose content (23.4-24.1%). Good brown rice recovery (80.2-80.1%). Premium milling recovery (71.2-72.2%) and G1 headrice recovery (45.9-53.4%). Long (7.2 mm) and slender grain shape (3.3-3.4)
High amylose content (23.1-23.2%). Fair brown rice recovery (79.6-80.0%). Premium milling recovery (70.9-72.3%) and G1 headrice recovery (47.5-53.7%). Long (6.9-7.1 mm) and slender grain shape (3.2). 100% acceptability and slightly creamish in the raw form
Low amylose content (16.8-16.9%). Fair brown rice recovery (77.9-79.2%). Premium milling recovery (70.2-70.8%) and headrice recovery of 64.1-65.6%. Long (6.4-6.8 mm) and slender grain shape (3.0-3.2). Slightly creamish in cooked and raw forms.
Intermediate amylose content (17.8%). Fair brown rice (78.6%). Premium milling (72.4%) and headrice recovery (61.5%). Medium (6.2mm) and intermediate grain shape (2.9). Zinc content (25.5mg.kg-1). Slightly creamish in the raw form
Intermediate amylose content (16.5%). Fair brown rice (78.6%). Premium milling (70.8%) and Grade 2 headrice recovery (43.8%). Extra long (7.5mm) and slender grain shape (23.6). Zinc content (27.3 mg.kg-1).
Low amylose content (16.4%). Fair brown rice (78.8%). Premium milling (72.5%) and Grade 1 headrice recovery (46.4%). Long (7.2mm) and slender grain shape (3.4). Zinc content (27.3 mg.kg-1).
Intermediate amylose content (18.6%). Fair brown rice (78.4%). Premium milling recovery (70.9%) and Grade 1 headrice recovery (56.3%). Long (6.6mm) and slender grain shape (3.1). Aromatic and tasty when cooked
Intermediate amylose content (20%). Fair brown rice (78.4%). Premium milling recovery (70.3%). Grade 1 headrice recovery (54.4%). Long (7.1mm) and slender grain shape (3.4). No aroma and slightly creamish when cooked and with 95% acceptability.
Intermediate amylose content (20.4%). 6.3% Protein Content. Fair brown rice (78.6%). Premium milling recovery (70.6%). Grade 1 head rice (54.9%). Long (6.6 mm) and intermediate grain shape (3.0). No aroma and slightly white in color when cooked, with 75% acceptability.

Low amylose content (14.1%). Fair brown rice (78.4%). Premium milling (70.7%) and headrice recovery of 62.9%. Long (6.8mm) and intermediate grain shape (2.9). Slightly creamish when cooked.
Low amylose content (14.10%). Fair brown rice (78.4%). Premium milling recovery (70.6%). Grade 1 headrice recovery (56.4%). Long (6.9mm) and slender grain shape(3.0). Slightly creamish when cooked.
Low amylose content (14%). Fair brown rice (79.1%). Premium milling recovery (71.2%). Grade 1 headrice recovery (53%). Long (7mm) and slender grain shape (3.1). Slightly aromatic, slightly creamish and slightly tasty when cooked.
Low amylose content (12.2%). Fair brown rice (79.5%). Premium milling (71.6%) and headrice recovery (59.1%). Long (6.9mm) and intermediate grain shape (3.0). Slightly creamish in cooked and raw forms. 100% acceptability in cooked form.
Low amylose content (12.2%). Fair brown rice (78.4%). Premium milling recovery (70.5%). Grade 1 headrice recovery (55.0%). Long (6.6mm) and intermediate grain shape (2.9). Slightly creamish in cooked and raw form.
Low amylose content (12.5%). Fair brown rice (73.4%). Grade 1 milling recovery (68.4%). Grade 2 headrice recovery (45.8%). Long (7.1 mm) and intermediate grain shape (2.8). Slightly white in cooked and raw form with no aroma
Intermediate amylose content (20 %). Grade 1 brown rice (80%). Premium milling recovery (72.6%). Grade 1 headrice recovery (49.1%). Long (7.1 mm) and intermediate grain shape (3.0). Slightly white in cooked and raw form with no aroma

Intermediate amylose content (19.6 %). Fair brown rice (78.6%). Premium milling recovery (71.1%). Grade 2 headrice recovery (44.2%). Long (7.0 mm) and slender grain shape (3.4). Slightly creamish in cooked and raw forms with no aroma.
Low amylose content (12.7 %). Fair brown rice (78.3%). Premium milling recovery (70.2%). Grade 1 headrice recovery (51.2%). Long (6.9 mm) and intermediate grain shape (3.0). Slightly grayish in cooked and raw forms with no aroma.
Low amylose content (12.4 %). Fair brown rice (78.2%). Grade 1 milling recovery (69.4%). Grade 1 headrice recovery (50.5%). Long (6.7 mm) and slender grain shape (3.1). Slightly whitish in cooked and raw forms with no aroma.
Low amylose content (15.8 %). Fair brown rice (78.9%). Grade 1 milling recovery (69.6%). Grade 1 headrice recovery (56.2%). Long (7.0 mm) and slender grain shape (3.3). Slightly creamish in raw form with no aroma and slightly whitish in cooked form with no aroma.
Intermediate amylose content (18.7 %). Fair brown rice (78.9%). Grade 1 milling recovery (69.6%). Grade 1 headrice recovery (56.2%). Long (7.0 mm) and slender grain shape (3.3). Slightly whitish in cooked form with no aroma and slightly creamish in raw form with no aroma.
Intermediate amylose content (18.7 %). Fair brown rice (78.9%). Grade 1 milling recovery (69.6%). Grade 1 headrice recovery (56.2%). Long (7.0 mm) and slender grain shape (3.3). Whitish in raw and cooked form with no aroma.

Intermediate amylose content (19.8 %).Good brown rice (80.6%). Premium milling recovery (71.7%). Grade 1 head rice recovery (51.2%). Medium (6.0 mm) and intermediate grain shape (2.7). Slightly creamish in raw and cooked form with no aroma.
High amylose content (23.1 %).Fair brown rice (79.4%). Premium milling recovery (71.4%). Grade 1 head rice recovery (49.2%). Medium (6.1 mm) and intermediate grain shape (2.6). Whitish in raw and cooked form with no aroma.
Intermediate amylose content (19.8 %).Fair brown rice (79.1%). Premium milling recovery (70.1%). Grade 2 head rice recovery (44.7%). Long (7.0 mm) and slender grain shape (3.3). Slightly creamish in cooked form and slightly whitish in raw form. No aroma.
Low amylose content (14.5 %).Fair brown rice (78.1%). Grade 1 milling recovery (69.7%). Premium head rice recovery (57%). Long (7.0 mm) and slender grain shape (3.3). Slightly grayish in cooked and raw form with no aroma.
Low amylose content (13.8 %).Fair brown rice (77.8%). Grade 1 milling recovery (69.1%). Grade 1 head rice recovery (51.6%). Long (6.5 mm) and intermediate grain shape (2.8). Slightly whitish in cooked form and grayish in raw form. No aroma.

Low amylose content (13.2%).Fair brown rice (78.4%). Grade 1 milling recovery (69.3%). Grade 1 head rice recovery (55.5%). Long (6.8 mm) and intermediate grain shape (3.1). Whitish in cooked and raw form. No aroma.
High amylose content (23.4%).Fair brown rice (78.3%). Grade 1 milling recovery (69.2%). Grade 2 head rice recovery (43.8%). Long (7.2 mm) and slender grain shape (3.5). Slightly whitish in cooked and raw form. No aroma.
Intermediate amylose content (19.8%).Fair brown rice (78.4%). Grade 1 milling recovery (69.2%). Grade 2 head rice recovery (42.6%). Long (7.3 mm) and slender grain shape (3.6). Slightly whitish in raw form and whitish in cooked form. No aroma.
High amylose content (24.4%).Fair brown rice (79.9%). Premium milling recovery (70.3%). Grade 2 head rice recovery (39.5%). Long (6.8 mm) and slender grain shape (3.3). Whitish in rawand cooked form. No aroma.
Intermediate amylose content (18.3%).Fair brown rice (79.9%). Premium milling recovery (71.9%). Grade 1 head rice recovery (49.6%). Long (6.9 mm) and slender grain shape (3.1). Slightly creamish in cooked form and slightly whitish in raw form. No aroma.

Intermediate amylose content (18.3%).Fair brown rice (79.9%). Premium milling recovery (71.9%). Grade 1 head rice recovery (49.6%). Long (6.9 mm) and slender grain shape (3.1). Slightly whitish in cooked form and slightly creamish in raw form. No aroma.
Intermediate amylose content (18.2%).Fair brown rice (78.7%). Grade 1 milling recovery (67.5%). Grade 3 head rice recovery (33.4%). Long (7.3 mm) and slender grain shape (3.7). Whitish in cooked form and slightly whitish in raw form. No aroma.
Intermediate amylose content (18.2%).Fair brown rice (78.7%). Grade 1 milling recovery (67.5%). Grade 3 head rice recovery (33.4%). Long (7.3 mm) and slender grain shape (3.7). Slightly whitish in cooked and raw form. No aroma.
Intermediate amylose content (17.7%).Fair brown rice (79.4%). Premium milling recovery (70.9%). Grade 2 head rice recovery (41.2%). Long (6.8 mm) and intermediate grain shape (3.0). Whitish in cooked and raw form. No aroma.
Low amylose content (13.1%).Fair brown rice (78.2%). Grade 1 milling recovery (69.1%). Grade 1 head rice recovery (48.1%). Long (7.0 mm) and slender grain shape (3.4). Sligtly whitish in cooked and raw form. No aroma.

Intermediate amylose content (19.1%).Fair brown rice (78.7%). Grade 1 milling recovery (69.9%). Grade 1 head rice recovery (48.4%). Long (6.8 mm) and slender grain shape (3.3). Sligtly creamish in cooked form and whitish raw form. No aroma.
Intermediate amylose content (19.8%).Good brown rice (80%). Premium milling recovery (70.7%). Grade 2 head rice recovery (44.5%). Long (7.0 mm) and slender grain shape (3.1). Sligtly whitish in cooked form and whitish raw form. No aroma.
Intermediate amylose content (21.9%).Fair brown rice (79.1%). Premium milling recovery (70.6%). Grade 3 head rice recovery (36.1%). Long (7.3 mm) and slender grain shape (3.2). Sligtly creamish in cooked and raw form. No aroma.