Comparison of Quantum Krylov and UCCSD for Molecular Ground-State Approximation via Subspace Projection

Jihyeon Park*

This work compares Quantum Krylov and UCCSD as non-variational basis generators for molecular ground-state approximation in a projected-subspace framework. Rather than optimizing one ansatz, I compare how well each basis family generates a useful low-energy subspace. Across the benchmark set, Quantum Krylov reaches chemical accuracy with substantially fewer basis states than the non-variational UCC constructions, showing a stronger expressivity-per-basis trend as system size grows. At the same time, per-state resource estimates show that Quantum Krylov states are much more expensive to prepare because they rely on Trotterized Hamiltonian evolution, whereas selected UCCSD states remain shallow and inexpensive. The resulting full-stack conclusion is a clear tradeoff: Quantum Krylov is favored at the algorithmic layer by its smaller required subspace, while selected UCCSD is favored at the logical and hardware layers by its lower circuit cost.

I. INTRODUCTION AND MOTIVATION

Quantum algorithms are widely expected to offer an eventual advantage over classical methods for describing quantum many-body systems, especially in electronic-structure problems where the Hilbert space grows exponentially with system size. Quantum phase estimation (QPE) is one of the representative examples because, in principle, it can recover eigenvalues to high precision when provided with a sufficiently good initial state [1]. In practice, however, QPE requires deep coherent time evolution and fault-tolerant resources that remain beyond the reach of current hardware for most chemically relevant systems.

These limitations motivated the development of hybrid algorithms such as the variational quantum eigensolver (VQE) [2]. VQE replaces long coherent evolution with repeated preparation and measurement of a parameterized trial state, while a classical optimizer updates the circuit parameters to minimize the measured energy. This hybrid structure is much more compatible with near-term hardware, but it introduces a different bottleneck: the variational optimization itself can be costly, sensitive to initialization, and difficult to scale when the ansatz contains many parameters.

As an alternative, subspace methods have also been developed for molecular ground-state estimation. Instead of optimizing a single wavefunction, these methods prepare a set of basis states and solve a projected eigenvalue problem within the span of those states. In this setting, the quality of the chosen basis becomes the central issue. A better basis can improve the accuracy of the projected ground-state estimate, while a lower-cost basis can reduce circuit depth, gate count, and measurement overhead.

This motivates the present study. Among the many basis-construction strategies proposed in the literature, Quantum Krylov states generated by Hamiltonian evolution [3] and UCCSD states generated by chemically motivated single and double excitations [4] are two representative state-of-the-art directions. Quantum Krylov uses the Hamiltonian itself to generate candidate directions in Hilbert space, whereas UCCSD uses an excitation structure inherited from coupled-cluster theory. Although many variations of both approaches exist, these two families serve here as representative Hamiltonian-guided and chemistry-guided basis constructions.

A. Problem Statement

There are several distinct ways to evaluate a quantum state-preparation strategy for ground-state estimation. In a QPE-type setting, a prepared state is used as the initial state for phase estimation, so its quality is judged mainly by its overlap with the target eigenstate. In a VQE-type setting, the state or ansatz is used as the trial form in a variational optimization, so its quality is judged by the minimum energy reachable after parameter updates. In a subspace method, by contrast, a family of prepared states is treated as a basis set, and the quality of that family is judged by the projected eigensolver built from their span.

This work adopts the third viewpoint. The main object of interest in this paper is not just one trial state, but the basis generator itself. Evaluating Quantum Krylov and UCCSD through the subspace they generate makes it possible to compare their collective expressivity, which can be richer than what is visible from a single state alone. Quantum Krylov is therefore treated here as a basis-family construction rather than as a one-state ansatz, and subspace projection provides the natural framework for comparing it with UCCSD. The concrete problem studied in this paper is molecular ground-state energy estimation. Starting from the same Hartree-Fock reference and comparing basis families at the same target subspace size K, this study asks which family delivers the lower projected ground-state energy for a given state-preparation cost. Exact FCI is used as the reference on the small calibration systems, while the larger-system comparisons are made against a TrimCI reduced-

* park687@wisc.edu

space reference that is first validated against FCI in the tractable cases.

II. METHODS / MODEL ASSUMPTIONS

A. Quantum Basis

1. Quantum Krylov Basis

Quantum Krylov methods construct a basis from repeated Hamiltonian action on a reference state and are naturally suited to projected eigensolvers [3]. Formally, the Quantum Krylov subspace of order K generated from $|\phi_{\mathrm{HF}}\rangle$ is

$$\mathcal{K}_K(\hat{H}, |\phi_{\mathrm{HF}}\rangle) = \mathrm{span}\{|\phi_{\mathrm{HF}}\rangle, \hat{H}|\phi_{\mathrm{HF}}\rangle, \hat{H}^2|\phi_{\mathrm{HF}}\rangle, \ldots\}. \tag{1}$$

In the circuit-based implementation used here, a time-evolved version is more convenient, and each additional Quantum Krylov basis state is generated from the Hartree-Fock reference using Trotterized real-time evolution,

$$|\phi(t_k)\rangle \approx e^{-it_k\hat{H}}|\phi_{\mathrm{HF}}\rangle. \tag{2}$$

The main characteristic of the Quantum Krylov basis is that it is Hamiltonian-guided: the Hamiltonian itself determines the directions added to the span. Its most important continuous parameter is the evolution time t_k assigned to each basis state, while the Trotter order and number of Trotter steps are fixed globally. In principle, the time steps can be selected by grid search or by heuristic spacing rules for time-propagated basis states [5, 6]. In this paper, the Quantum Krylov times are chosen as an equally spaced sequence $t_k = k\Delta t$, and the spacing Δt is selected by grid search over candidate values. Candidate spacings that lead to a poorly conditioned overlap matrix are rejected, and the final choice is the spacing that gives the lowest projected ground-state energy among the remaining candidates. Quantum Krylov is therefore treated as a non-variational basis generator whose parameter-selection step is restricted to this explicit grid-search protocol rather than to a full continuous optimization loop.

2. UCCSD Basis

UCCSD constructs basis states from a chemically motivated excitation manifold built on top of the Hartree-Fock reference [4]. Its general form is

$$|\phi(\boldsymbol{\theta})\rangle = e^{\hat{T}(\boldsymbol{\theta})-\hat{T}^\dagger(\boldsymbol{\theta})}|\phi_{\mathrm{HF}}\rangle, \tag{3}$$

where the cluster operator is truncated to single and double excitations,

$$\hat{T}(\boldsymbol{\theta}) = \sum_{ia}\theta_i^a \hat{a}_a^\dagger \hat{a}_i + \frac{1}{4}\sum_{ijab}\theta_{ij}^{ab}\hat{a}_a^\dagger\hat{a}_b^\dagger\hat{a}_j\hat{a}_i. \tag{4}$$

Here i, j denote occupied spin orbitals and a, b denote virtual spin orbitals. The main characteristic of UCCSD is that it is chemistry-guided rather than directly Hamiltonian-guided: the basis is restricted to the singles-and-doubles excitation structure expected to capture a large fraction of weak-to-moderate correlation effects.

The main parameters of UCCSD are the excitation amplitudes θ_i^a and θ_{ij}^{ab} attached to the singles-and-doubles pool. In a standalone VQE implementation, these amplitudes would be determined by variational energy minimization. In the present non-variational comparison, they are instead assigned classically using warm-start rules such as second-order Møller-Plesset perturbation theory (MP2) [7], which provides perturbative amplitudes, or coupled-cluster singles and doubles (CCSD) [8], which provides amplitudes from an iterative coupled-cluster calculation. Two UCCSD constructions are used in this work. The first is a one-shot full-versus-screened comparison, in which the full singles-and-doubles pool is evaluated directly against a screened or selected variant that retains only the larger-amplitude excitations. The second is the head-to-head K-dependent basis construction used later in the paper: starting from one MP2 or CCSD amplitude vector, the individual excitation generators are ranked by amplitude magnitude, and each added basis state is formed by applying one ranked generator at its fixed classical amplitude to the Hartree-Fock reference. This latter construction is not a conventional single-state UCCSD ansatz; rather, it is deliberately chosen to be closer in spirit to the GCIM idea [9], where one builds a nonorthogonal basis from several UCC-generated states and then solves the projected eigenvalue problem in their span instead of optimizing one combined UCC circuit. Thus, MP2 or CCSD provides the amplitudes, while the full-versus-screened or generator-ranked choice determines how those amplitudes are turned into benchmark basis states.

B. TrimCI-Based Reduced-Space Setup

The working space is reduced to a TrimCI-selected determinant space before the Quantum Krylov and UCCSD benchmark states are constructed. The reason is that the expensive steps of the unreduced workflow scale with the size of the full many-body space. In particular, an explicit qubit-state representation of a generic n-qubit Quantum Krylov or UCCSD state scales as 2^n, while exact FCI in the fixed-electron sector scales combinatorially with the number of spin orbitals and electrons, through the determinant-space dimension $\binom{n}{N_e}$. In the naive full-space setting of this project, this keeps the routine end-to-end benchmark only in the small-system regime, practically through LiH in the main workflow, with H_2O / STO-3G still manageable only as a standalone reference check. Such a regime is too small to support a meaningful extrapolation in qubit count, so

larger systems are needed. In practice, this means that qubit-scaling trends should be probed at least beyond the 20-qubit scale, and preferably into the 40–50 qubit range. By projecting first into a compact selected determinant space, the main linear-algebra cost is controlled instead by the number of kept determinants rather than the full Hilbert-space dimension.

For a given molecular system, the first step is to run TrimCI in a shared restricted Hartree-Fock molecular-orbital basis and obtain an approximate ground state in determinant form,

$$|\Psi_{\mathrm{TrimCI}}\rangle = \sum_{\mu=1}^{M} c_\mu \, |D_\mu\rangle \, . \tag{5}$$

In the present workflow, the practical TrimCI limiter is not an explicit energy-accuracy threshold but a determinant budget, controlled mainly by the parameter `ndets`, together with optional exploration controls such as `ndets_explore` and `nexploration`. After the TrimCI run, the selected determinants are ranked by weight $|c_\mu|^2$.

The working subspace for the benchmark is then defined by truncating this list to the top N determinants,

$$\mathcal{S}_N = \mathrm{span}\{|D_1\rangle \, , |D_2\rangle \, , \ldots , |D_N\rangle\}, \tag{6}$$

with $N \leq M$. This additional cutoff removes the low-weight tail of the TrimCI expansion and gives a compact determinant basis for the later Quantum Krylov and UCCSD constructions. The adequacy of this reduced determinant space is validated explicitly against exact FCI on the smaller systems before the larger reduced-space benchmarks are interpreted.

All operators used in the benchmark are then represented directly in this N-determinant basis. The projected Hamiltonian is

$$H^{(N)}_{\mu\nu} = \langle D_\mu | \, \hat{H} \, | D_\nu \rangle \, , \tag{7}$$

and the UCCSD excitation generators are represented in the same way,

$$G^{(a,N)}_{\mu\nu} = \langle D_\mu | \left(\hat{\tau}_a - \hat{\tau}_a^\dagger \right) | D_\nu \rangle \, . \tag{8}$$

Because the determinant basis itself is orthonormal, its intrinsic overlap matrix is the identity. The nontrivial overlap matrix appears only after the Quantum Krylov or UCCSD benchmark states are generated inside this reduced space.

Given these projected operators, the benchmark basis states are constructed entirely within $\mathcal{S}_N$. Quantum Krylov states are generated from the projected Hamiltonian and the projected Hartree-Fock reference. UCCSD states are generated from the projected excitation generators, either as one-shot full-versus-screened UCCSD states or, in the later head-to-head comparison, as ranked single-generator states built from MP2 or CCSD amplitudes. If a comparison uses K benchmark basis states, then K Quantum Krylov or UCCSD states are constructed in this reduced determinant space before the final projected eigensolver is applied.

TABLE I. Benchmark systems used in the reduced-space study. All systems are treated in the STO-3G basis. The Hamiltonian Pauli-term column reports the number of terms in the mapped qubit Hamiltonian; the C_3H_8 entry is listed as a coarse extrapolated estimate because direct compilation of the full 46-qubit Hamiltonian was prohibitively slow in the present environment.

System	Qubits	Electrons	Hamiltonian Pauli terms	Role
H_2	4	2	15	Exact-FCI calibration
H_4	8	4	185	Exact-FCI calibration
LiH	12	4	631	Exact-FCI calibration
N_2	20	14	3,327	Reduced-space scaling
C_2H_4	28	16	8,919	Reduced-space scaling
C_3H_8	46	26	$\sim 3.8 \times 10^4$	Reduced-space scaling

C. Benchmark Systems

The benchmark ladder is chosen to serve two purposes at once: direct small-system validation against exact FCI and larger-system scaling tests in the TrimCI-reduced workflow. All systems are treated in the STO-3G basis. The smaller systems H_2, H_4, and LiH are used to calibrate and validate the reduced-space construction against exact reference data, while N_2, C_2H_4, and C_3H_8 extend the study into a regime where full-space simulation would already be impractical in the end-to-end workflow but the TrimCI-based reduced-space approach remains tractable.

D. Projected Subspace Construction

All basis families start from the same Hartree-Fock reference state $|\phi_{\mathrm{HF}}\rangle$. Given a set of prepared basis states $\{|\phi_i\rangle\}_{i=0}^{K-1}$, the projected Hamiltonian and overlap matrices are formed as

$$H_{ij} = \langle \phi_i | \, \hat{H} \, | \phi_j \rangle \, , \tag{9}$$

$$S_{ij} = \langle \phi_i | \phi_j \rangle \, , \tag{10}$$

and the lowest generalized eigenvalue of

$$Hc = E_{\mathrm{sub}} S c \tag{11}$$

is used as the subspace energy estimate.

E. Generalized Eigensolver and Overlap Conditioning

The projected problem takes the generalized form rather than the ordinary form because the Quantum Krylov and UCCSD basis states are generally nonorthogonal. Thus the subspace coefficients satisfy $Hc =$

$E_{\text{sub}}Sc$, and this reduces to an ordinary eigenvalue problem only in the special case of an orthonormal basis with $S = I$.

Because the nonorthogonal basis can contain nearly dependent directions, the overlap matrix may become ill-conditioned. To control this, S is eigendecomposed as

$$S = U \operatorname{diag}(s_i) U^\dagger, \tag{12}$$

and directions with overlap eigenvalues below a threshold are discarded. In the current implementation, this is done with an absolute eigenvalue test $s_i > s_{\text{eval_cutoff}}$ rather than with a relative rule. The released benchmark notebooks use $s_{\text{eval_cutoff}} = 10^{-8}$ in one notebook and a looser 10^{-10} cutoff in the H_4 and LiH notebooks, so the working basis is orthogonalized only on the numerically stable part of the span. The transformation

$$X = U_{\text{keep}} \operatorname{diag}(s_i^{-1/2}) \tag{13}$$

then maps the retained nonorthogonal basis to an orthonormal basis, and the Hamiltonian is transformed as

$$\widetilde{H} = X^\dagger H X. \tag{14}$$

The final projected estimate is obtained by solving the ordinary eigenvalue problem for $\widetilde{H}$ in this retained orthogonal subspace.

In a hardware-oriented execution, the matrix elements of H and S would have to be obtained from Hadamard tests or related overlap-estimation techniques. In the present work, however, the purpose is to compare Quantum Krylov and UCCSD as basis generators rather than to benchmark matrix-element measurement protocols. The benchmark states are therefore represented analytically in the reduced TrimCI space, and the entries of H and S are formed directly from inner products and projected operators before solving the generalized eigenvalue problem. For larger reduced-space instances, iterative eigensolvers such as Davidson or LOBPCG are natural choices.

F. Comparison Protocol

The comparison in this work is non-variational for both basis families. Quantum Krylov and UCCSD are not judged by the result of a full parameter-optimization loop, but by the projected subspaces generated from their method-specific state-construction rules. This choice isolates the basis generator itself rather than the trainability of a variational ansatz.

Under this protocol, both methods are evaluated for the same molecular Hamiltonian, in the same orbital basis, from the same Hartree-Fock reference state, inside the same TrimCI-defined reduced determinant space, and at the same target benchmark-basis size K. The projected ground-state estimate is always obtained from the same Rayleigh-Ritz procedure together with the same overlap-conditioning procedure. Exact FCI is used as the validation reference on the smaller calibration systems, while the larger systems are compared within the same TrimCI-reduced workflow validated against those smaller exact benchmarks.

The two families are allowed to keep their natural parameterization. For Quantum Krylov, the defining parameter is the time spacing Δt that generates the equally spaced evolution times $t_k = k\Delta t$ used for the time-evolved basis states. In this paper, Δt is selected by grid search. For UCCSD, the defining ingredients are the amplitudes attached to the singles-and-doubles excitation pool together with the choice of whether to keep the full pool or a screened subset. In the present non-variational setting, these amplitudes are supplied by classical chemistry information such as MP2 or CCSD rather than by a VQE optimization loop. The comparison therefore does not claim to identify the best standalone Quantum Krylov or the best standalone UCCSD-VQE result. Instead, it asks which family provides the more expressive projected subspace, within the same reduced-space representation, for the cost of preparing its states and operators.

For the explicit K-by-K head-to-head comparison reported later, the UCCSD side is organized as two separate ranked generator families, one built from MP2 amplitudes and one built from CCSD amplitudes. In each family, the individual singles-and-doubles generators are ranked by amplitude magnitude, and each added basis state is formed by applying one ranked generator at its fixed classical amplitude to the Hartree-Fock reference. Thus $K = 1$ is the Hartree-Fock state itself, $K = 2$ adds the top-ranked single-generator state, $K = 3$ adds the second-ranked single-generator state, and so on. This makes the later UCCSD comparison closer in spirit to a GCIM-style [9] nonorthogonal basis built from UCC generators than to a nested full-UCCSD truncation ladder. Curves may terminate before the target K if the overlap-conditioning rule rejects additional nearly dependent states.

G. Assumptions and Parameters

Several assumptions are held fixed so that the ansatz comparison remains controlled.

- The main benchmark is noiseless and focuses on basis quality and preparation cost, not hardware noise or full shot-noise modeling.
- The primary fairness axis is basis size K; total trainable parameter count is tracked only as a secondary diagnostic.
- Quantum Krylov and UCCSD use the same molecular Hamiltonian, the same Hartree-Fock reference state, the same projected-subspace objective, and the same validation reference.

- The comparison is non-variational. Quantum Krylov basis states are specified by chosen evolution times, while UCCSD basis states are specified either as one-shot full-versus-screened ansätze or as ranked single-generator states built from MP2 or CCSD amplitudes.

- Candidate states are rejected when the overlap matrix becomes numerically ill-conditioned. In the current code, this is implemented with an absolute smallest-eigenvalue cutoff on S, with $s_{\text{eval_cutoff}} = 10^{-8}$ in one benchmark notebook and 10^{-10} in the looser H_4 and LiH notebooks.

- Small calibration systems are used intentionally so that the reduced-space workflow can be checked against exact FCI data before moving to the larger systems.

III. RESULTS AND VALIDATION

A. Quantum Krylov Time-Step Selection by Grid Search

Even though the benchmark avoids a full variational optimization loop, the Quantum Krylov basis still depends on a method-specific parameter, namely the time spacing Δt used to define the equally spaced sequence $t_k = k\Delta t$. For this reason, the first stage of the results is a grid-search comparison over candidate Δt values. For each molecule, candidate spacings are scanned, choices that lead to an ill-conditioned overlap matrix are discarded, and the remaining spacing with the lowest projected ground-state energy is retained. In this grid-search stage, the target Quantum Krylov size is fixed at $K = 4$ so that only the time spacing is varied; for H_2, the retained TrimCI space is only two-dimensional, so the largest nontrivial Quantum Krylov basis is $K = 2$. This fixed-K scan is used as a calibration step before the later head-to-head comparison extends Quantum Krylov to the largest numerically stable target up to $K = 20$.

Figure 1 shows the representative N_2 scan. The smallest spacings are rejected because the overlap matrix becomes too ill-conditioned, while a shallow minimum appears once the time-evolved states become sufficiently distinct. At this fixed $K = 4$ calibration point, the best spacing is $\Delta t = 0.150$ for N_2, with a reduced-space Quantum Krylov gap of 0.698 mHa relative to the TrimCI reference. The same grid-search protocol was applied to every benchmark system, and the resulting fixed-K calibration spacings are summarized in the table below. A striking feature of this summary is that the preferred spacing is very stable across the ladder: $\Delta t = 0.150$ is selected for H_4, LiH, N_2, and C_2H_4, while the largest system C_3H_8 shifts only slightly to $\Delta t = 0.300$.



FIG. 1. Representative Quantum Krylov grid search for N_2 in the TrimCI-reduced space. The plotted quantity is the projected energy gap $E_{\text{QK}} - E_{\text{ref}}$ as a function of the spacing Δt at $K = 4$. Red crosses mark candidate spacings rejected by the overlap cutoff, and the vertical dotted line marks the selected spacing.

System	Qubits	N_{det}	Best Δt	Best gap (mHa)
H_2	4	2	0.100	-0.000
H_4	8	20	0.150	0.244
LiH	12	69	0.150	0.631
N_2	20	73	0.150	0.698
C_2H_4	28	73	0.150	0.584
C_3H_8	46	71	0.300	0.408

B. Full and Screened UCCSD with MP2/CCSD Parameters

The second stage of the results examines whether the full UCCSD excitation pool is actually needed once amplitudes are fixed by classical chemistry information. To do this, I evaluate non-variational UCCSD states at MP2 and CCSD amplitudes, and then compare the full singles-and-doubles pool against a screened variant that keeps only operators with amplitude magnitude above the fixed threshold 10^{-2}. In this work, the screened ansatz is the selected-UCC surrogate used for hardware-cost reduction. Adaptive UCC variants are not included, because they would reintroduce an iterative variational loop outside the intended scope of this non-variational comparison. The detailed screening study is reported on H_2, H_4, and LiH, where exact FCI remains available and full one-shot UCCSD evaluation is still routine in the present implementation.

Figure 2 shows the representative LiH tradeoff. In the full UCCSD pool, the MP2- and CCSD-parameterized states have FCI gaps of 11.224 and 9.002 mHa, respectively. After screening, those gaps change only slightly, to 11.308 mHa for MP2 and 9.073 mHa for CCSD, while the operator counts collapse from 92 to 6 and 8. This is the central screening result of the subsection: at least



FIG. 2. Representative LiH comparison for full and screened one-shot, non-optimized UCCSD with MP2- and CCSD-parameterized amplitudes. Top: energy gaps relative to exact FCI, with the dashed line marking chemical accuracy. Bottom: retained excitation-operator counts for the same four variants.

on the calibration systems, most of the one-shot UCCSD benefit survives after discarding the small-amplitude tail of the excitation pool.

The same pattern appears across the rest of the small-system set. For H_2, screening reduces the operator count from 3 to 1 with no visible change in the reported gap. For H_4, screening reduces the operator count from 26 to 10 while changing the gap by less than 0.03 mHa. In all three systems, the CCSD-initialized ansatz is more accurate than the MP2-initialized ansatz, but the main hardware-relevant message is that the screened construction tracks the full construction very closely at substantially lower operator count.

For the later cross-family comparison, this subsection therefore motivates a nested UCCSD construction rather than a single fixed UCCSD state. In that later comparison, the excitation amplitudes are ranked by importance and used to build growing MP2-based and CCSD-based UCCSD hierarchies, so that the UCCSD side can also be studied as a function of basis size K.

System	Variant	Gap to FCI (mHa)	Ops
H_2	MP2 full	9.473	3
	MP2 screened	9.473	1
	CCSD full	5.104	3
	CCSD screened	5.104	1
H_4	MP2 full	23.613	26
	MP2 screened	23.613	10
	CCSD full	18.793	26
	CCSD screened	18.768	10
LiH	MP2 full	11.224	92
	MP2 screened	11.308	6
	CCSD full	9.002	92
	CCSD screened	9.073	8

TABLE II. Full and screened one-shot UCCSD variants on the small calibration systems. Reported gaps are relative to exact FCI, and 'Ops' counts the retained excitation operators.

C. Head-to-Head Quantum Krylov versus UCCSD as K Increases

With the Quantum Krylov spacing fixed by grid search, the final stage is a direct comparison of how the basis families themselves improve as K grows across the full six-system benchmark ladder. The figure therefore spans H_2, H_4, LiH, N_2, C_2H_4, and C_3H_8, so that the head-to-head comparison can be read from the smallest to the largest reduced-space instances used in this work. In every case, the projected energy is reported relative to the same TrimCI reduced-space reference energy. On the Quantum Krylov side, the plotted curve uses the best grid-searched spacing for the largest numerically stable Quantum Krylov target up to $K = 20$. On the UCCSD side, I use two separate non-variational generator-ranked families, one ranked by MP2 amplitudes and one ranked by CCSD amplitudes. Each added UCCSD basis state is generated from a single ranked excitation operator acting on the Hartree-Fock reference at its fixed classical amplitude, so the ordered sequence is HF at $K = 1$, then the top-ranked single-generator state, then the second-ranked single-generator state, and so on. In this sense, the head-to-head UCCSD comparison is closer to a GCIM-style nonorthogonal basis built from UCC generators than to a nested truncation of one full UCCSD circuit [9].

Figure 3 shows that the three families behave quite differently once the comparison is phrased in terms of how quickly their projected span improves with K. Quantum Krylov is the only family that consistently drives the projected energy close to the reduced-space reference across the full ladder: it is exact for H_2 by $K = 2$, reaches the numerical floor for H_4 by $K = 12$, and then continues to converge to 10^{-4}–10^{-6} mHa levels on LiH, N_2, C_2H_4, and C_3H_8. For this extended K sweep, the final benchmark-selected Quantum Krylov spacings are $\Delta t^* = 0.10$ (H_2), 1.95 (H_4), 1.70 (LiH), 1.05 (N_2), 1.20 (C_2H_4), and 0.95 (C_3H_8). The largest numerically stable Quantum Krylov targets range from $K = 2$ for H_2 up to the full $K = 20$



FIG. 3. Head-to-head comparison of Quantum Krylov, MP2, and CCSD basis families as the benchmark basis size K increases across the full benchmark ladder. The vertical axis shows the absolute projected-energy gap $|E_{\mathrm{sub}} - E_{\mathrm{ref}}|$ on a logarithmic scale, with values below 10^{-6} mHa clipped to that floor for display. The gray shaded region marks gaps below chemical accuracy (1.6 mHa). 'MP2' and 'CCSD' denote the generator-ranked UCC basis families built from fixed MP2 or CCSD amplitudes, respectively, where $K = 1$ is Hartree-Fock and each additional basis state is generated by one more ranked single UCC excitation generator. The horizontal dotted lines use the same MP2 and CCSD colors and mark the corresponding full-pool one-shot UCCSD baselines in the same reduced determinant space. Curves terminate when the overlap-conditioning rule rejects further nearly dependent states or when the available reduced-space dimension is exhausted.

target on N_2, C_2H_4, and C_3H_8.

The UCC side shows a different pattern. On the smallest systems the generator-ranked MP2 and CCSD families can also approach the reference, but as the systems grow they converge much more slowly and often terminate early because of overlap conditioning. The CCSD-ranked family is usually the stronger of the two on the small calibration systems, reaching 0.287 mHa on H_4 and 0.0209 mHa on LiH, but it still remains at 20.09 mHa on N_2, 21.16 mHa on C_2H_4, and 33.47 mHa on C_3H_8 at the largest sampled K. The MP2-ranked family stays above chemical accuracy as well on the larger systems, with final gaps of 19.82 mHa on N_2, 11.92 mHa on C_2H_4, and 21.98 mHa on C_3H_8. The main conclusion from the six-system figure is therefore that Quantum Krylov gains much more from increasing K than the amplitude-ranked single-generator UCC families once the problem moves beyond the small calibration regime. At the same time, the difference between the MP2 and CCSD curves shows that this GCIM-like UCC construction depends on two practical choices: the classical amplitude source changes how the excitation generators are ranked and therefore which basis states are added first, and the resulting nonorthogonal basis can have different numerical conditioning, which affects how far each family can be extended before the overlap matrix becomes ill-conditioned.

D. Extrapolated Basis Size at Chemical Accuracy

The qubit-scaling trend can be summarized more compactly by extracting, from Fig. 3, the basis size required to reach chemical accuracy. For each system and basis family, I define K_{chem} as the first sampled K for which $|E_{\mathrm{sub}} - E_{\mathrm{ref}}| \leq 1.6$ mHa. If the sampled data do not reach that threshold, I fit the tail of the convergence curve to the log-linear form

$$\log_{10} |E_{\mathrm{sub}} - E_{\mathrm{ref}}| = a + bK, \tag{15}$$

using the last up to five available points, and solve for the crossing with 1.6 mHa to obtain an extrapolated estimate of K_{chem}.

Figure 4 makes the scaling separation between the basis families explicit. Quantum Krylov reaches chemical accuracy with only a handful of basis states across the full ladder: $K_{\mathrm{chem}} = 2$ for H_2, 11 for H_4, 2 for LiH, 5 for N_2, and 4 for both C_2H_4 and C_3H_8. The



FIG. 4. Estimated basis size required to reach chemical accuracy as a function of qubit count. Filled markers denote systems where the sampled data already reach the 1.6 mHa target, while open markers denote extrapolated values obtained from a log-linear fit to the tail of the convergence curve. The dashed curves are descriptive family-wise linear least-squares fits of K_{chem} versus qubit count and should be read only as visual guides to the overall scaling trend. The right-side callout reports the corresponding descriptive 100-qubit estimates from those linear trend fits.



FIG. 5. Per-basis-state resource scaling versus qubit count, shown as two separate panels rather than a dual-axis plot. Panel (a) reports the logical T count for Krylov-trotterization, UCCSD-MP2, and UCCSD-CCSD. Panel (b) reports the compiled circuit depth for the same three methods together with a fixed-accuracy Krylov-qubitization depth proxy. Thin dashed guide curves show the power-law fits used for the right-side 100-qubit estimates. The Krylov-qubitization line appears only in panel (b), since no compiled PREPARE/SELECT oracle was available to estimate a comparable T-count. The C_3H_8 Krylov-trotterization point in both panels is an extrapolated proxy, while the remaining compiled values are taken directly from the current PennyLane resource estimates.

generator-ranked UCC families behave very differently. On the small systems, the MP2 and CCSD constructions still reach chemical accuracy with moderate basis sizes, namely $(2, 11, 5)$ for MP2 and $(2, 10, 7)$ for CCSD on H_2, H_4, and LiH. Beyond that regime, however, chemical accuracy is not reached within the sampled K window, and the extrapolated requirements rise into the tens of basis states. The extrapolated values are approximately $K_{\mathrm{chem}} \approx 44$, 34, and 63 for the MP2 family on N_2, C_2H_4, and C_3H_8, and $K_{\mathrm{chem}} \approx 47$, 41, and 85 for the corresponding CCSD family. This final scaling summary is consistent with the direct convergence plots: within the present reduced-space benchmark, Quantum Krylov maintains a much more favorable expressivity-per-basis trend than the non-variational UCC constructions as the system size increases.

IV. FULL-STACK IMPACT / RESOURCE IMPLICATIONS

A. Resource Estimation

Because the dependence on the number of basis states K has already been studied in the previous section, the resource comparison here focuses on the intrinsic cost of preparing one additional basis state beyond the shared Hartree-Fock reference. For Quantum Krylov, the reported values are based on first-order Trotterized time evolution with $n = 2$ steps at the benchmark-selected Δt^*. For UCCSD-MP2 and UCCSD-CCSD, the reported values are averaged over the ranked single-generator basis states used in the head-to-head comparison. The combined table below summarizes logical T count, circuit depth, and two-qubit-gate count per basis state. The C_3H_8 Quantum Krylov row is an extrapolated proxy based on the exact smaller-system trend, while the remaining rows are taken directly from the current PennyLane resource estimates.

For LiH, the same resource picture is reflected in the comparison between full and amplitude-selected UCCSD. The selected construction preserves the one-shot warm-start energy to within about 0.1 mHa while reducing the compiled circuit cost by more than an order of magnitude. In the MP2 branch, the selected construction lowers the gate count from 9344 to 720, and in the CCSD branch from 9344 to 748, while the energy gap changes only from 11.22 to 11.31 mHa and from 9.00 to 9.07 mHa, respectively. This makes selected UCCSD the clearly preferable chemistry-inspired basis when the objective is to reduce circuit cost without reintroducing a variational loop.

B. Full-Stack Impact

At the quantum-algorithm layer, Quantum Krylov remains attractive because its stronger expressivity lowers the basis size required for chemical accuracy, as shown in Fig. 4. Since the number of projected Hamiltonian and overlap elements grows roughly as K^2, this also reduces the shot burden of the subspace solve. Using the

Method	H_2			H_4			LiH			N_2			C_2H_4			C_3H_8		
	T	Depth	2Q	T	Depth	2Q	T	Depth	2Q	T	Depth	2Q	T	Depth	2Q	T	Depth	2Q
Krylov	1232	26	12	16192	362	56	55440	1250	148	261536	5926	380	784784	17774	756	4235201.0	98801.3	2289.4
UCCSD-MP2	352	20	14	352	20	14	352	20	14	352	20	14	352	20	14	352	20	14
UCCSD-CCSD	352	20	14	277.7	15.7	10.6	297.3	16.8	11.5	352	20	14	352	20	14	352	20	14

TABLE III. Per-basis-state resource summary for the current Quantum Krylov, UCCSD-MP2, and UCCSD-CCSD constructions. Each method block reports the average logical T count, circuit depth, and two-qubit-gate count for one additional basis state beyond the shared Hartree-Fock reference. The C_3H_8 Quantum Krylov entry is an extrapolated proxy based on the exact smaller-system trend; all remaining entries are drawn directly from the current PennyLane resource estimates.

extracted K_{chem} values, the number of unique basis pairs $K_{\text{chem}}(K_{\text{chem}}+1)/2$ is only 15 for Quantum Krylov on N_2 and 10 for Quantum Krylov on both C_2H_4 and C_3H_8, compared with 990 and 2016 for the MP2-ranked UCC family and 1128 and 3655 for the CCSD-ranked UCC family on N_2 and C_3H_8, respectively. In this sense, Quantum Krylov is algorithmically more efficient: it reaches the same reduced-space accuracy with a much smaller projected subspace.

At the logical-processor layer, however, Fig. 5 shows the opposite trend: each Quantum Krylov basis state is much deeper than the corresponding selected UCC basis state because it requires Trotterized Hamiltonian evolution. The added Krylov-qubitization depth curve indicates one possible route to reduce this Quantum Krylov depth bottleneck for larger systems, although it does so by introducing block-encoding oracle and ancilla overhead rather than by producing a shallow native circuit. A simple depth-dominated logical-fidelity estimate can be obtained by requiring the total logical failure probability for preparing the K_{chem} basis states to remain below a budget η,

$$p_L^{\text{req}} \approx \frac{\eta}{K_{\text{chem}} d_{\text{state}}}, \tag{16}$$

where d_{state} is the per-state circuit depth. In this estimate, I set $\eta = 10^{-2}$ as a rough one-pass budget for basis-state preparation. The estimate is optimistic because it ignores the repeated sampling needed to measure the projected H and S matrices, but it is still sufficient to expose the stack-level trend.

This estimate shows that selected UCCSD is substantially easier at the logical and error-correction layers. On N_2, for example, the resulting requirement is about 3.0×10^{-7} for Quantum Krylov, compared with 1.1×10^{-5} for both MP2- and CCSD-ranked UCC. On C_3H_8, the corresponding values are 2.4×10^{-8}, 7.9×10^{-6}, and 5.9×10^{-6}. Under any standard fault-tolerant architecture, a smaller target logical error rate translates into a larger code distance and therefore a heavier physical-processor overhead. In that sense, selected UCCSD is favored at the logical and physical layers even though it is less expressive at the algorithmic layer.

The overall full-stack conclusion is therefore mixed rather than one-sided. Quantum Krylov suppresses the basis-size and shot-scaling burden, while selected UCCSD relaxes the required logical fidelity per circuit. Which family is closer to practical implementation depends on the dominant bottleneck: if projected-subspace size and sampling cost dominate, Quantum Krylov is more appealing; if logical depth and error-correction overhead dominate, selected UCCSD is more realistic. This benchmark therefore suggests that the ansatz choice is not just an algorithmic question, but a stack-wide co-design question linking quantum chemistry, logical resources, and hardware feasibility.

V. CONCLUSION

This work compared Quantum Krylov and UCCSD as non-variational basis generators for molecular ground-state approximation in a projected-subspace framework. By placing both families in the same reduced determinant-space workflow, the benchmark focused on basis quality rather than on a variational optimization loop. The main numerical result is that Quantum Krylov reaches chemical accuracy with a substantially smaller basis size than the non-variational UCC constructions across the benchmark ladder, indicating a stronger expressivity-per-basis trend as the system size grows.

At the same time, the resource analysis shows that this algorithmic advantage does not translate into a simple hardware advantage. Quantum Krylov basis states are much more expensive to prepare because they require Trotterized Hamiltonian evolution, whereas selected UCCSD states remain shallow and comparatively cheap. The resulting picture is therefore a genuine trade-off: Quantum Krylov is favored when the dominant concern is projected-subspace size and sampling burden, while selected UCCSD is favored when the dominant concern is logical depth and hardware cost per basis state.

Taken together, these results suggest that basis design for quantum chemistry should be evaluated at more than one layer of the stack. In the present benchmark, Quantum Krylov is the more powerful basis family at the algorithmic level, but selected UCCSD remains attractive from the perspective of implementation. This makes the choice between them not just a question of accuracy, but a co-design problem linking ansatz expressivity, logical resources, and hardware feasibility.

VI. LIMITATIONS AND FUTURE WORK

A. Current Limitations

The main limitation of the present benchmark is that the working subspace is not the full FCI space for the entire system ladder. Instead, the Hamiltonian, reference energy, and benchmark basis states are all defined in a TrimCI-selected determinant space. The reduced space reproduces the exact FCI result very well on the small calibration systems, but the larger-system conclusions in this report should still be interpreted as reduced-space results rather than exact full-space statements.

A second limitation is that the resource comparison is not yet fully symmetric between the two basis families. In the current implementation, Quantum Krylov basis states are generated from Trotterized Hamiltonian evolution, whereas the UCCSD side is represented by selected single-generator states ranked by MP2 or CCSD amplitudes. This is a meaningful practical comparison, but it does not exhaust all possible hardware-aware variants of either family. In particular, a truncated or selected Quantum Krylov construction could make the resource comparison more balanced against selected UCCSD.

B. Future Work

One natural extension is to make the hardware-level benchmark more symmetric. This includes repeating the resource analysis with truncated-Hamiltonian Quantum Krylov constructions and comparing those results directly against warm-start and selected-UCCSD variants. Such a study would clarify whether lower-cost Quantum Krylov states can preserve enough of the expressivity advantage seen in the present benchmark. A closely related direction is to investigate whether qubitization- based implementations of Quantum Krylov time evolution can reduce the resource cost of the Quantum Krylov basis relative to the current Trotterized construction.

A second direction is to expand the choice of reference state beyond Hartree-Fock. For example, both Quantum Krylov and UCCSD could be initialized from a stronger warm-start state such as the TrimCI reduced-space ground state rather than the Hartree-Fock determinant. That would help separate the effect of basis generation from the effect of the initial reference itself.

It would also be valuable to connect the present resource estimates more directly to concrete hardware specifications. Mapping the benchmark circuits to specific platform assumptions, such as superconducting or trapped-ion devices, would make it possible to assess which basis family is closer to realistic implementation on a given hardware stack.

Finally, the present comparison suggests several algorithmic extensions. One possibility is to design a new hybrid basis family that combines the compact expressivity of Quantum Krylov with the shallower state-preparation structure of selected UCCSD. Another is to investigate related subspace constructions and variations of UCC/Quantum Krylov ideas, for example Q-SENSE [10]. A complementary theoretical direction is to study more carefully why Quantum Krylov bases so effectively describe the ground state in these electronic-structure problems, and whether that behavior can be predicted from spectral or correlation properties of the Hamiltonian.

VII. REPOSITORY AND REPRODUCIBILITY

This project is organized around a small set of benchmark notebooks and validation scripts. The main software stack uses PennyLane for quantum-state construction, PySCF for classical electronic-structure reference calculations, and TrimCI for reduced determinant-space selection.

Repository: `https://github.com/jhp1859/p759_final_project/draft_deliveries`

Main entry points:

- `Fair_Subspace_Benchmark.ipynb`, `Fair_Subspace_Benchmark_H4.ipynb`, and `Fair_Subspace_Benchmark_LiH.ipynb`: end-to-end small-system benchmarks for Quantum Krylov and UCCSD.
- `validate_trimci_subspace.py`: validation of the TrimCI reduced-space reference against exact FCI, together with the figure `TrimCI_FCI_Subspace_Validation.pdf`.
- `warmstart_uccsd_compare.py`: MP2- and CCSD-parameterized full and screened UCCSD warm-start comparisons.
- `uccsd_screening_benchmark.py`: compact full-versus-screened UCCSD benchmark producing the LiH representative figure and the H_2/H_4/LiH summary table.
- `krylov_vs_uccsd_k_comparison.py`: six-system head-to-head comparison of Quantum Krylov and nested MP2/CCSD-ranked UCCSD hierarchies as the basis size K increases.
- `selected_krylov_all_systems.py`: selected Quantum Krylov study reproducing the K-convergence, chemical-accuracy, and resource-scaling figures for the benchmark ladder.
- `h2o_trimci_krylov_uccsd_compare.py`: auxiliary larger-system reduced-space comparison script.
- `Resource_Estimation_Subspace.ipynb`: resource-estimation summaries for the benchmark ansatz families.

[1] D. S. Abrams and S. Lloyd, Quantum algorithm providing exponential speed increase for finding eigenvalues and eigenvectors, Physical Review Letters **83**, 5162 (1999).

[2] A. Peruzzo, J. McClean, P. Shadbolt, M.-H. Yung, X.-Q. Zhou, P. J. Love, A. Aspuru-Guzik, and J. L. O'Brien, A variational eigenvalue solver on a quantum processor, Nature Communications **5**, 4213 (2014).

[3] C. L. Cortes and S. K. Gray, Quantum krylov subspace algorithms for ground- and excited-state energy estimation, Physical Review A **105**, 022417 (2022).

[4] A. Anand, P. Schleich, S. Alperin-Lea, P. W. K. Jensen, S. Sim, M. Díaz-Tinoco, J. S. Kottmann, M. Degroote, A. F. Izmaylov, and A. Aspuru-Guzik, A quantum computing view on unitary coupled cluster theory, Chemical Society Reviews **51**, 1659 (2022).

[5] N. H. Stair, R. Huang, and F. A. Evangelista, A multireference quantum krylov algorithm for strongly correlated electrons, Journal of Chemical Theory and Computation **16**, 2236 (2020).

[6] J. Cohn, M. Motta, and R. M. Parrish, Quantum filter diagonalization with compressed double-factorized hamiltonians, PRX Quantum **2**, 040352 (2021).

[7] C. Møller and M. S. Plesset, Note on an approximation treatment for many-electron systems, Physical Review **46**, 618 (1934).

[8] G. D. Purvis and R. J. Bartlett, A full coupled-cluster singles and doubles model: The inclusion of disconnected triples, The Journal of Chemical Physics **76**, 1910 (1982).

[9] M. Zheng, B. Peng, A. Li, X. Yang, and K. Kowalski, Unleashed from constrained optimization: Quantum computing for quantum chemistry employing generator coordinate inspired method, npj Quantum Information **10**, 127 (2024).

[10] S. Patel, P. Jayakumar, R. Huang, T. Zeng, and A. F. Izmaylov, Quantum seniority-based subspace expansion: Linear combinations of short-circuit unitary transformations for the electronic structure problem, arXiv preprint arXiv:2509.01061 10.48550/arXiv.2509.01061 (2025).