

Tutorial 2

ST2137-2420

Material

This tutorial covers basic Python syntax and data manipulations with **pandas**. It provides practice on chapter 2 from the course textbook.

The following numpy functions may be useful: `np.where`, `np.exp`. The following pandas functions too: `pd.Series.value_counts`.

In the final section, we use the concept of MLE, which you would be familiar from ST2132. It is a method of deriving optimal estimators for distribution parameters.

Dataset: Liverpool

The dataset `liverpool_2223_season.csv` contains information on games that Liverpool Football Club¹ played. The data was obtained from [footballref](#). The team played 19 other teams, and played each of them Home and Away.

1. Read the dataset into Python:

	Date	Day	Venue	GF	GA	Opponent
0	2022-08-06	Sat	Away	2	2	Fulham
1	2022-08-15	Mon	Home	1	1	Crystal Palace
2	2022-08-22	Mon	Away	1	2	Manchester Utd
3	2022-08-27	Sat	Home	9	0	Bournemouth
4	2022-08-31	Wed	Home	2	1	Newcastle Utd

2. Tabulate the number of times Liverpool played games on the different days.
3. Add two columns to the dataset:
 - One string column named **result**, that contains W (for Win), D (for Draw) or L (for Loss).
 - One numeric column **pts** that contains the value of 3, 1 or 0, corresponding to a win, draw or loss respectively.
4. Write a **for** loop to produce the following output, which computes the total number of points Liverpool won off each team:

```
Points against Arsenal: 1
Points against Aston Villa: 4
Points against Bournemouth: 3
Points against Brentford: 3
Points against Brighton: 1
Points against Chelsea: 2
```

The output above has been truncated. Your output should include all 19 opponents.

5. It is often said that a team needs to accumulate 40 points to be safe from relegation (forced down to a lower division). Add a column that computes the cumulative number of points Liverpool obtains, and use it to identify when Liverpool first accumulated 40 points.

¹Liverpool play in the English Premier League.

6. Use the internet to figure out what this function does. Then use it to compute the length of the longest winning streak that Liverpool had.

```
import re
def fn1(str1):
    out_string = re.split('[LD]+', str1)
    st1 = [len(x) for x in out_string]
    return st1

rle_out = fn1(liv.result.str.cat())
np.max(rle_out)
```

7. The following questions are on the use of the slice operator in Python. Use `.iloc` along with the slice operator to retrieve:
1. The first 10 rows
 2. Alternate rows from the first 10 rows, starting with the first.
 3. Every alternate row, and columns Date, Venue, GF, GA and result.
 4. The last 5 rows.
 5. All rows in reverse order.

Truncated Poisson

Suppose that $X \sim \text{Pois}(\lambda)$, and the distribution of Y is given by

$$P(Y = y) = P(X = y | X > 0), \quad y = 1, 2, \dots$$

This is known as the truncated Poisson distribution. Given observations y_1, \dots, y_n , the Maximum Likelihood Estimate $\hat{\lambda}$ is given by the solution to

$$\bar{y} = \frac{\hat{\lambda}}{1 - \exp(-\hat{\lambda})}$$

Suppose we observe the following readings from 30 observations of Y :

$Y = 1$	$Y = 2$	$Y = 3$	$Y = 5$
12	14	3	1

8. Perform a grid search over $(0.5, 2)$ with spacing 0.01 to identify the MLE.