

Problem 1. [20 pts] Harmonic oscillator via Hamilton-Jacobi

Consider the Hamiltonian of a one-dimensional harmonic oscillator

$$H(q, p) = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 q^2, \quad (1)$$

where ω is a constant. What is the the solution to the corresponding Hamiltonian-Jacobi equation $W(q, t)$ (you don't need to evaluate the integral over q)? And using W , find the dynamical trajectories $q(t)$ and $p(t)$.

Problem 2. [40 pts] Wave- to ray-optics with Hamilton-Jacobi

The setup: If light is going through a medium with a local index of refraction $n(y)$ and is partially moving in the y and z direction, it is governed by the following wave equation

$$\frac{n(y)^2}{c^2} \frac{\partial^2 E}{\partial t^2} - \frac{\partial^2 E}{\partial y^2} - \frac{\partial^2 E}{\partial z^2} = 0, \quad (2)$$

in this problem, we will derive ray optics which gives a particle description of these waves.

- (a) To find how light moves in this medium when it is at frequency ω , substitute

$$E(y, z, t) = Ae^{-i(\omega t - \frac{\omega}{c}W(y, z))}$$

and keep terms only of order $(\omega/c)^2$ (high frequency approximation).

- (b) Solve for $\frac{\partial W}{\partial z}$ putting the equation in the form of a Hamilton-Jacobi equation

$$\frac{\partial W}{\partial z} = -H\left(y, \frac{\partial W}{\partial y}\right). \quad (3)$$

What is the Hamiltonian $H(y, p)$? (*Hint:* Many times in this problem you will encounter square roots. The **positive** square roots will give you what you want but think about what the negative square roots mean physically; they are physical.)

- (c) Compute the Lagrangian $L(y, \frac{dy}{dz}) = \frac{dy}{dz}p - H(y, p)$ eliminating (y, p) in favor of $(y, \frac{dy}{dz})$. *Note:* The spatial direction z is acting like “time” in this situation.
- (d) Show that the “action” $S = \int L(y, \frac{dy}{dz})dz$ we are minimizing is related to the *least time* of a ray between two points (this is called “Fermat’s principle”)

$$T = \int_A^B \frac{n(y)}{c} \sqrt{1 + \left(\frac{dy}{dz}\right)^2} dz = \int_A^B \frac{n(y)}{c} \sqrt{dy^2 + dz^2} \quad (4)$$

- (e) Write a paragraph (at least 1/2-page) about what we have shown as a correspondence between classical waves and particles/rays. Compare this with the massive particles in classical mechanics. Is there an equation like Eq. (2) in classical mechanics? What about quantum mechanics?
- (f) For $n(y) = n_0 \frac{y}{y_0}$, solve for $y(z)$ assuming $y(0) = y_0$ and $y'(0) = 0$. Furthermore, using $W(y, z) = bz + g(y)$, solve for $W(y, z)$ and write out the full form of $E(y, z, t)$ in this approximation.