Supplementary data: Reference-country composition and age-standardized PCOS-attributed infertility contrasts: China and 18 G20 countries, 1990–2023

Companion supplementary methods and results

29 October 2026

1 Scope and computational audit

This supplement documents the secondary computations accompanying the main article. The primary estimand is the 1990–2023 change in the total PCOS-attributed infertility difference under contemporary GBD reference shares. The analyses use aggregate GBD estimates and matched demographic data; the upstream GBD model was not reestimated.

The GBD inputs comprise total burden export `1b13e337`, matched population `916826fb`, annual primary/secondary Number export `416daf73` and separate endpoint Number/Rate export `76c5523c`. The full filenames, query specifications and SHA-256 values are in the accompanying README, input manifest and data-acquisition document. Expected coverage is 19 countries by 34 years by seven ages: 4,522 population cells under each source and 27,132 prevalence cells across two causes and three infertility types. The original GBD burden/stratum exports also contain other measures/metrics; the focused analysis selects prevalence counts and reconstructs rates from matching GBD populations.

The 1,064 original endpoint subtype rates agreed with the reconstructed rates to within 1.16×10^{-11} per 100,000. Core R/Python outputs agreed within 5.69×10^{-13}. The 1,440 decomposition sensitivity rows passed the exact additive identity with maximum residual 1.99×10^{-13}. Numerical tolerances are stated in code; no artificial research observations were created. Archived aggregate tables are used only for post-computation regression checks, not to generate results.

2 Sensitivity definitions and interpretation

Country exclusions renormalize within-age shares among the remaining reference countries. The age-20–49 specification sets the age-15–19 weight to zero and renormalizes the remaining six standard weights. Core sensitivity scenarios use both GBD and UN shares, 1990 or 2010 start years, and 2021, 2022 or 2023 endpoints. Complete results are in `composition_sensitivities.csv`; Table S2 displays 1990–2023 total PCOS results. A 2021 endpoint always uses GBD 2023 estimates for 2021, not a previous GBD release.

The symmetric identity is applied within age and subtype. Table S3 displays age-by-type components for the GBD main reference. Annual chaining applies the same identity to each con-

secutive year, then adds components from 1990 to 2023 (Table S4). For total PCOS under GBD shares, chaining gives +87.11 for reference composition rather than +88.65 from the endpoint calculation, while the net change remains +151.54. Neither path is an uncertainty interval.

Total-variation distance between GBD and UN reference-country share vectors is one half of the sum of absolute country-share differences for each age and year (Table S5). Age-weighted national shares used in Figure S5 average the age-specific country proportions using the same fixed standard age weights. They are descriptive summary weights, not actual pooled all-age population percentages.

The seven unnormalized published standard weights are 0.082891, 0.078012, 0.075914, 0.073217, 0.068281, 0.061474 and 0.055113, summing to 0.494902. We normalize them over ages 15–49. Using them without renormalization multiplies restricted-age standardized rates by 0.494902 (Table S6). These conventions must not be mixed. Table S8 evaluates all 128 corner perturbations of the unnormalized weights by half the last published decimal before renormalization. This rounding envelope quantifies numerical precision sensitivity alone.

3 Endometriosis terminal-year audit

The endometriosis example is secondary. China's modeled restricted-age standardized rate increased from 43.13 to 53.30 per 100,000 in 2022–2023, or +23.58%. The absolute increment was +10.17; primary and secondary contributions were +0.74 and +9.43. The historical maximum positive annual change over the preceding 1991–2022 intervals was +1.56%. Selected other 2022–2023 changes were +14.82% in the United States, +11.93% in Japan and -0.41% in the Republic of Korea. All 19 countries appear in Table S7 and Figure S6; the within-age/type accounting appears in Figure S7.

These are repeated computations from the same GBD release. They establish source-output concordance, not independent epidemiological corroboration, a pandemic effect, a new biological event or a clinical mechanism. The endometriosis net China-minus-reference change over 1990–2023 is small, so its decomposition percentage would be unstable; absolute components are provided in the machine-readable results. No mechanism is inferred for the final-year jump.

4 Joint uncertainty and acquisition status

The GBD input files contain point estimates and marginal lower/upper limits, without posterior draw columns. Availability of a provider-confirmed joint posterior array aligned across all dimensions required by this analysis was not established from the study inputs. This does not imply that suitable draws do not exist or cannot be supplied. The present analyses therefore do not quantify joint posterior uncertainty.

Until official joint alignment is confirmed across countries, ages, years, types and any uncertain denominators, the covariance of custom pooled estimates remains unknown. Original marginal limits must not be added or simulated independently to manufacture intervals. G20 resampling measures a different, hypothetical country-selection process. UN share replacement assesses benchmark construction, not GBD posterior uncertainty. All custom contrasts,

components, rankings and ratios remain descriptive point estimates. If draws are obtained, recompute each entire formula within each draw and report posterior quantiles, alongside diagnostics for near-zero net-change denominators.

5 GATHER reporting map

The map below concerns secondary calculations and does not claim replication of GBD's upstream evidence synthesis. Author-provided declarations remain XX pending author confirmation before submission.

Item	Topic	Location or limitation
1	Indicators/populations	Main Methods: females, ages, years, countries, infertility types and cause IDs.
2	Funding	Main Funding section: XX pending author completion.
3	Data identification	Official GBD and UN resources; acquisition specifications and hashes.
4	Inclusion/exclusion	Complete fixed country set; explicit sensitivity removals and age restrictions.
5	Source characteristics	Modeled GBD outputs and UN population counts; original evidence belongs to GBD.
6	Input bias	Main Discussion: heterogeneity, cause allocation, denominator and model limitations.
7	Other inputs	Matched populations and published rounded standard weights.
8	Input access	Galaxy source archive, IHME acquisition instructions and public UN subset.
9	Conceptual overview	Main equations: pooling, standardization, contrast and symmetric decomposition.
10	Analytical details	Main Methods, this supplement and executable code.
11	Model selection	No fitted trend/forecast candidate selection in this focused revision; benchmark definitions are explicit.
12	Evaluation	Deterministic sensitivity outputs, official endpoint checks and R/Python comparison.
13	Uncertainty approach	No joint draws; no custom posterior intervals.
14	Source code	Reproducibility archive and versioned GitHub repository.
15	Estimates	Aggregate CSV results and generated tables.
16	Quantitative uncertainty	Incomplete: available marginal limits do not identify derived uncertainty.
17	Interpretation	Main Discussion and established Kitagawa literature.
18	Limitations	Main Discussion and uncertainty section above.

Table S1: Endpoint sensitivity of total infertility under contemporary GBD shares.

Cause	Year	China	Reference	Difference
Endometriosis	1990	63.70	66.63	-2.93
Endometriosis	2021	43.48	51.93	-8.45
Endometriosis	2022	43.13	51.89	-8.76
Endometriosis	2023	53.30	53.51	-0.20
PCOS	1990	333.56	606.48	-272.92
PCOS	2021	601.79	745.23	-143.44
PCOS	2022	602.09	726.09	-124.00
PCOS	2023	605.08	726.45	-121.38

Rates/differences per 100,000 females, standardized within ages 15–49. All years come from GBD 2023.

Table S2: Reference-composition sensitivity for total PCOS, 1990–2023.

Shares	Specification	Net change	Share term	100S/net
GBD	Main	+151.54	+88.65	58.5
GBD	Ages 20–49	+183.78	+109.22	59.4
GBD	Without Argentina	+152.71	+89.88	58.9
GBD	Without Australia	+151.60	+89.24	58.9
GBD	Without Brazil	+143.40	+94.90	66.2
GBD	Without Canada	+152.10	+90.16	59.3
GBD	Without France	+151.04	+87.32	57.8
GBD	Without Germany	+151.68	+87.96	58.0
GBD	Without India	+141.07	+56.16	39.8
GBD	Without Indonesia	+188.27	+98.35	52.2
GBD	Without Italy	+134.12	+70.88	52.9
GBD	Without Japan	+117.03	+59.35	50.7
GBD	Without Mexico	+155.77	+94.91	60.9
GBD	Without Republic of Korea	+154.16	+90.25	58.5
GBD	Without Russian Federation	+162.61	+110.34	67.9
GBD	Without Saudi Arabia	+153.44	+89.87	58.6
GBD	Without South Africa	+149.37	+88.67	59.4
GBD	Without Türkiye	+153.63	+90.25	58.7
GBD	Without United Kingdom	+150.10	+86.94	57.9
GBD	Without United States of America	+136.85	+82.66	60.4
UN WPP	Main	+150.69	+86.76	57.6
UN WPP	Ages 20–49	+182.92	+107.12	58.6
UN WPP	Without Argentina	+151.96	+88.07	58.0
UN WPP	Without Australia	+150.67	+87.28	57.9
UN WPP	Without Brazil	+141.88	+92.39	65.1
UN WPP	Without Canada	+151.23	+88.28	58.4
UN WPP	Without France	+149.81	+85.05	56.8
UN WPP	Without Germany	+150.80	+86.01	57.0
UN WPP	Without India	+139.63	+52.67	37.7
UN WPP	Without Indonesia	+185.19	+95.35	51.5
UN WPP	Without Italy	+133.50	+69.18	51.8
UN WPP	Without Japan	+115.95	+57.20	49.3
UN WPP	Without Mexico	+155.64	+93.67	60.2
UN WPP	Without Republic of Korea	+153.32	+88.38	57.6
UN WPP	Without Russian Federation	+162.89	+109.28	67.1
UN WPP	Without Saudi Arabia	+153.13	+88.57	57.8
UN WPP	Without South Africa	+148.87	+87.12	58.5
UN WPP	Without Türkiye	+152.85	+88.41	57.8
UN WPP	Without United Kingdom	+148.79	+84.57	56.8
UN WPP	Without United States of America	+137.34	+82.18	59.8

Country-exclusion rows omit one reference country and renormalize shares; ages 20–49 renormalizes the age standard. The last column is a signed accounting ratio in percent, not an uncertainty interval.

Table S3: GBD-share age-by-type decomposition of PCOS contrast changes, 1990–2023.

Type	Age	China rates	Ref. rates	Ref. shares	Net change
Total	15–19	+20.02	-19.20	-2.27	-1.46
Total	20–24	+54.07	-41.65	+11.32	+23.75
Total	25–29	+50.64	-40.16	+15.36	+25.85
Total	30–34	+45.85	-36.00	+15.79	+25.64
Total	35–39	+43.43	-32.79	+20.42	+31.07
Total	40–44	+38.48	-28.65	+23.76	+33.60
Total	45–49	+19.01	-10.18	+4.27	+13.10
Primary	15–19	+8.82	-3.65	-0.75	+4.41
Primary	20–24	+19.84	-13.65	+8.74	+14.92
Primary	25–29	+12.18	-9.79	+10.09	+12.48
Primary	30–34	+4.86	-3.58	+5.95	+7.24
Primary	35–39	+2.71	-2.34	+4.32	+4.69
Primary	40–44	+2.67	-3.38	+3.29	+2.57
Primary	45–49	+1.59	-1.71	+0.21	+0.09
Secondary	15–19	+11.21	-15.55	-1.52	-5.87
Secondary	20–24	+34.24	-28.00	+2.58	+8.83
Secondary	25–29	+38.47	-30.37	+5.27	+13.37
Secondary	30–34	+40.98	-32.43	+9.84	+18.40
Secondary	35–39	+40.73	-30.45	+16.10	+26.38
Secondary	40–44	+35.82	-25.27	+20.48	+31.03
Secondary	45–49	+17.42	-8.47	+4.06	+13.01

Contributions per 100,000. Age/type contributions add exactly before rounding.

Table S4: Annual-chained decomposition summed across 1990–2023.

Cause	Type	Shares	Ref. rates	Ref. shares	Net change
Endometriosis	Primary	GBD	+6.95	-0.18	+6.79
Endometriosis	Primary	UN WPP	+6.98	-0.21	+6.79
Endometriosis	Secondary	GBD	+6.63	-0.28	-4.06
Endometriosis	Secondary	UN WPP	+6.50	-0.16	-4.07
Endometriosis	Total	GBD	+13.58	-0.46	+2.73
Endometriosis	Total	UN WPP	+13.48	-0.36	+2.72
PCOS	Primary	GBD	-35.11	+28.86	+46.40
PCOS	Primary	UN WPP	-35.22	+27.59	+45.02
PCOS	Secondary	GBD	-171.97	+58.25	+105.14
PCOS	Secondary	UN WPP	-172.44	+59.25	+105.67
PCOS	Total	GBD	-207.08	+87.11	+151.54
PCOS	Total	UN WPP	-207.66	+86.83	+150.69

The Chinese-rate component is identical under chaining and endpoint calculation. Results describe accounting-path sensitivity.

Table S5: Total-variation distance between GBD and UN country-share vectors.

Age	1990–2023 mean	2023
15–19	1.313	1.530
20–24	1.147	1.358
25–29	0.843	1.339
30–34	0.912	1.105
35–39	0.957	1.254
40–44	1.085	1.188
45–49	1.674	1.463

Distances are expressed in percentage points. The simple mean across seven age groups in 2023 is 1.32 percentage points.

Table S6: China endometriosis rate-convention reconciliation.

Year	Modeled cases	Restricted 15–49	Unnormalized weights
1990	192,845.45	63.697	31.524
2023	181,587.09	53.304	26.380

Rates per 100,000. The last column equals the preceding column multiplied by 0.494902. This is a numerical convention check, not independent validation.

Table S7: All-country endometriosis-attributed infertility terminal-year audit.

Country	2022–2023 change (%)	1991–2022 max (%)
Argentina	+2.14	+0.86
Australia	+2.01	+0.47
Brazil	-1.91	+3.62
Canada	-0.05	+10.07
China	+23.58	+1.56
France	-0.41	+0.99
Germany	-0.31	+8.49
India	+3.68	-0.39
Indonesia	+4.35	+0.07
Italy	+7.00	+3.62
Japan	+11.93	+1.38
Mexico	-1.87	+0.37
Republic of Korea	-0.41	+1.56
Russian Federation	+0.07	+1.86
Saudi Arabia	-0.45	-0.29
South Africa	-1.85	+3.78
Türkiye	-0.96	+0.26
United Kingdom	-0.00	+11.21
United States of America	+14.82	+3.88

The historical column is the largest signed annual percentage change over 1991–2022. All values are model estimates from one release; no significance test or mechanism is inferred.

Table S8: Standard-age-weight rounding envelopes for 2023 rate differences.

Cause	Type	Shares	Minimum difference	Maximum difference
PCOS	Total	GBD	-121.37685	-121.37559
PCOS	Total	UN WPP	-122.62203	-122.62076
PCOS	Primary	GBD	-123.53997	-123.53917
PCOS	Primary	UN WPP	-124.88451	-124.88370
PCOS	Secondary	GBD	2.16290	2.16381
PCOS	Secondary	UN WPP	2.26225	2.26317
Endometriosis	Total	GBD	-0.20251	-0.20240
Endometriosis	Total	UN WPP	-0.02923	-0.02912
Endometriosis	Primary	GBD	-7.93574	-7.93569
Endometriosis	Primary	UN WPP	-7.92095	-7.92090
Endometriosis	Secondary	GBD	7.73321	7.73331
Endometriosis	Secondary	UN WPP	7.89170	7.89180

All 128 corner perturbations of seven unnormalized weights by plus/minus half the last decimal, renormalized within ages 15–49. This envelope is numerical sensitivity, not a confidence interval.




Figure S1: 2023 total PCOS differences under country exclusions and age restriction. GBD population shares; all values are point estimates.




Figure S2: PCOS subtype differences under three benchmark definitions. UN current-year shares, UN fixed-1990 shares and equal-country weights are applied within age. Disease rates remain GBD estimates.




Figure S3: Signed PCOS composition ratios under reference and age specifications. The dashed line marks 50 percent, not a significance threshold. The 39.8–67.9 percent range is not a posterior interval.




Figure S4: Endometriosis total-infertility China-minus-reference trajectory. The highlighted endpoints show why the 2023 near-zero contrast does not summarize a stable trajectory.




Figure S5: Changes in age-weighted reference-country shares, UN WPP 2024. Country proportions within age are averaged with fixed standard age weights. These are composition changes, not disease-rate changes.




Figure S6: Endometriosis modeled rate changes across 19 countries, 2022–2023. This reproduces the source release. It does not establish an epidemiological event or external validation.




Figure S7: Age and infertility-type contributions to the Chinese endometriosis rate increment, 2022–2023. Standard-weighted contributions sum to +10.1723 per 100,000. Most of the signed modeled increment is secondary infertility; this is not an etiologic fraction.

Prior questions and the present estimand

This targeted comparison supports scientific positioning; it is not an exhaustive systematic review or proof that no prior analysis has used similar accounting. Bibliographic details and source URLs are provided in the main article reference list. No pooled-comparator error is attributed to a paper simply because it reports national age-standardized rates.

Table S9: Questions addressed by selected prior PCOS-attributed infertility analyses and the present study.

Study	Verified question or analysis	Relation to the present primary estimand
Liu et al., 2024	GBD 2019 global, regional and national prevalence, YLD and trends, 1990–2019. Published Methods describe direct age standardization.	Provides disease-burden context. It does not establish that a China-versus-18-country pool with our within-age weights was used. We do not reinterpret its individual national comparisons.
Liang et al., 2025	GBD 2021 burden and trends; the Methods describe a Das Gupta allocation to population growth, ageing and epidemiological factors.	A burden/count change and a national restricted-age standardized-rate difference against a changing pool are different targets. Our share term concerns countries within each age, not changing age structure.
Liang, 2026	Indexed abstract: GBD 2021 China and worldwide PCOS/related-infertility trends, including Joinpoint and ARIMA. Full comparator construction was not independently verified.	Supports a China/worldwide monitoring use case. No claim about an undocumented pooling formula is made; conclusions here do not test that paper's estimates.
Present study	GBD 2023 China-minus-contemporaneous-18-country pooled total-infertility difference, 1990–2023; fixed age weights.	Primary change and established symmetric allocation to national rates, reference rates and within-age country shares. UN shares test demographic-source sensitivity; fixed-1990/equal-country weights change the target.