Additional file 1: Supplementary methods and results

Companion to the Reproductive Health manuscript draft

25 October 2026

Additional arithmetic definitions

The same-population benchmark is $B_T = \sum_a P_{C,aT}(r_{C,aT} - r_{R,aT})/10^5$. It uses China's actual age distribution rather than standard weights and is not preventable burden. Age-specific contributions to the standardized difference are $w_a(r_{C,at} - r_{R,at})$. The seven published unnormalized standard weights, in ascending age order, are 0.082891, 0.078012, 0.075914, 0.073217, 0.068281, 0.061474 and 0.055113; they sum to 0.494902 before renormalization. References and outcome definitions are given in the main manuscript.

1 Supplementary methods and results

1.1 Decomposition of case-count changes

We expressed the case count as $F(N, \boldsymbol{f}, \boldsymbol{p}) = N \sum_a f_a p_a$, where N is the total female population aged 15–49, f_a is its age distribution, and p_a is age-specific prevalence as a proportion. For each factor, we averaged its marginal contribution across all six possible replacement orders between the starting and ending states. These Shapley contributions sum exactly to the observed modeled case-count change. We examined 1990–2023 and 2010–2023 for both causes in all 19 countries. Population size here refers specifically to females aged 15–49; age composition refers to redistribution within this range. The age-specific prevalence component includes epidemiological change, ascertainment and modeling changes and cannot be interpreted as a causal risk-factor contribution.

1.2 Trends, sensitivities and forecast validation

We fitted continuous piecewise linear models to log standardized rates, searching integer-year knot locations with zero to two knots and at least five observations per segment, counting shared boundary years in each segment. Model selection used the Bayesian information criterion, including a penalty for knot locations. Segment-specific annual percentage change was $100[\exp(\beta) - 1]$; average annual percentage change (AAPC) was obtained from duration-weighted log slopes. These fits summarize modeled trajectories and are not outputs of the proprietary Joinpoint program. AR(1) generalized least-squares refits retained the OLS-selected

knots; allowing up to three knots and restricting the period to 2010–2023 were additional sensitivities. Trend confidence intervals are conditional on selected knots and fixed GBD point estimates.

Reference sensitivities used age-specific country population shares fixed at 1990, excluded one reference country at a time, and restricted ages to 20–49 with renormalized weights. Following inspection of the terminal-year endometriosis increase, we added analyses ending in 2021 and 2022; these endpoint analyses were not prespecified. To evaluate numerical sensitivity to published weight rounding, we evaluated all 2^7 combinations of perturbing unnormalized weights by $\pm 0.5 \times 10^{-6}$. The resulting deterministic envelopes are not uncertainty intervals.

An exploratory forecasting module evaluated each Chinese age-specific prevalence series on the logit scale. Candidates were a constant-last-rate baseline, ARIMA with $p, q \in \{0, 1, 2\}$ and $d \in \{0, 1\}$ selected by training-set corrected AIC, and additive damped-trend exponential smoothing. ARIMA candidates failing convergence were excluded. Rolling training windows contained at least 20 years and assessed horizons 1, 3 and 5; a separate last-10-year holdout assessed horizons 1–10. We specified an exploratory retention rule before running this module: at least 10% lower mean absolute error (MAE) than the baseline in every age group in both validation schemes, with complete predictions. Future counts would additionally require compatible population projections; none were generated when validation failed. This module was not a validation against independent observed infertility data.

Table S1: Shapley decomposition of changes in China's modeled prevalent cases, thousands.

Cause	Period	Net change	Population size	Age composition	Prevalence
Endometriosis	1990–2023	-11.3	-8.7	30.8	-33.4
Endometriosis	2010–2023	6.1	-38.4	7.7	36.8
PCOS	1990–2023	783.4	-68.4	-8.9	860.7
PCOS	2010–2023	-129.8	-414.4	-18.2	302.9

Population size is the female population aged 15–49; age composition is its within-range distribution; prevalence is age-specific modeled infertility prevalence. Contributions may exceed the net change and need not have the same sign. Rounding can cause small displayed discrepancies.

Table S2: Restricted-age standardized YLD rates and YLD totals.

		YLDs, thousands		Rate per 100,000			
Cause	Year	China	Reference	China	Reference	D	RR
Endometriosis	1990	1.0	2.2	0.34	0.39	-0.05	0.872
Endometriosis	2023	1.0	2.6	0.29	0.31	-0.02	0.934
PCOS	1990	5.9	20.5	1.80	3.59	-1.79	0.502
PCOS	2023	10.1	35.5	3.31	4.22	-0.91	0.784

All values are derived point estimates. YLD: years lived with disability.

Table S3: Endpoint sensitivity added after inspection of terminal-year estimates.

Cause	End year	China ASPR	Ref. ASPR	D	RR	$D_T - D_{1990}$
Endometriosis	2021	43.48	51.93	-8.45	0.837	-5.52
Endometriosis	2022	43.13	51.89	-8.76	0.831	-5.83
Endometriosis	2023	53.30	53.51	-0.20	0.996	2.73
PCOS	2021	601.79	745.23	-143.44	0.808	129.47
PCOS	2022	602.09	726.09	-124.00	0.829	148.91
PCOS	2023	605.08	726.45	-121.38	0.833	151.54

The source release remains GBD 2023 in all rows. Ending in 2021 is not substitution of GBD 2021 estimates. Rate units: per 100,000 women.

Table S4: Conditional trend sensitivity for prevalence, 1990–2023.

Location	Cause	Model	AAPC (%)	Conditional 95% CI
China	Endometriosis	OLS, 0–2 knots	-0.86	(-1.00, -0.71)
China	Endometriosis	AR(1), fixed knots	-0.82	(-0.99, -0.65)
China	Endometriosis	OLS, 0–3 knots	-0.86	(-1.00, -0.71)
China	PCOS	OLS, 0–2 knots	1.89	(1.85, 1.93)
China	PCOS	AR(1), fixed knots	1.94	(1.81, 2.08)
China	PCOS	OLS, 0–3 knots	1.84	(1.80, 1.87)
Reference	Endometriosis	OLS, 0–2 knots	-0.74	(-0.78, -0.71)
Reference	Endometriosis	AR(1), fixed knots	-0.74	(-0.78, -0.69)
Reference	Endometriosis	OLS, 0–3 knots	-0.72	(-0.75, -0.68)
Reference	PCOS	OLS, 0–2 knots	0.53	(0.49, 0.58)
Reference	PCOS	AR(1), fixed knots	0.49	(0.42, 0.57)
Reference	PCOS	OLS, 0–3 knots	0.57	(0.53, 0.61)

Intervals condition on selected knots and fixed GBD point estimates. AR(1) retains the knots selected by OLS with up to two knots. These intervals exclude GBD, knot-selection and structural model uncertainty. Full periods, measures, knots and residual-correlation estimates are in the accompanying CSV.

Table S5: Age-specific forecast-validation MAE ratios relative to the naive baseline.

Cause	Model	Rolling: min–max	Last 10 years: min–max	Retained
Endometriosis	ARIMA	0.87–1.30	0.93–1.80	No
Endometriosis	ETS	1.00–1.51	0.86–2.10	No
PCOS	ARIMA	0.19–2.20	0.06–1.38	No
PCOS	ETS	0.20–1.79	0.09–1.28	No

Ranges span seven age groups. Values below 1 favor the candidate. Retention required every age-specific ratio to be below 0.9 in both schemes, with complete predictions. The rule was specified before running this exploratory module, not externally preregistered. Rolling MAE combines horizons 1, 3 and 5; the holdout combines horizons 1–10.

1.3 Acquisition and computation audit

The official query identifiers were:

Burden: `1b13e3376135ab41a4bc5e0ffc60e84f`.

Population: `916826fb285f8d3c3e53117dd2133406`.

Pilot: `0daed0e53f5ad8ebdf0dc00c23ccd110`.

Annual strata: `416daf735b9fc79bec35506514424e73`.

Endpoint strata: `76c5523c52f32caf342843daf2326fa4`.

Raw uncertainty limits remain in the original CSVs. Country and outcome mappings, source hashes, rounding sensitivities and all analytical tables are separate machine-readable files. Nine automated core checks plus the new reference-composition accounting checks cover core accounting identities, the complete data grid, denominator consistency and outcome gates; temporary artificial test fixtures were not used as research data.

The protocol was written before analysis but was not externally preregistered. The endpoint and weight-rounding sensitivities were added following quality inspection. Forecasting remained exploratory and was withheld after failed comparative validation. Primary/secondary stratification was completed using annual Number exports and independent endpoint Number/Rate checks. The symmetric reference-composition and age-by-type analyses were added during manuscript review and were not prespecified. External survey comparisons and future population scenarios were not completed; they were optional extensions. The annual stratum rate reconstruction reduced redundant metrics within the remaining GBD download allowance.

1.4 GATHER reporting map

This map applies GATHER to the secondary analysis; upstream input reconstruction and primary disease estimation belong to the original GBD study. It is not a claim of independent GBD model replication.

Item	Topic	Location or applicability
1	Indicators and populations	Methods: causes, ages, female sex, fixed countries and 1990–2023.
2	Funding	Declarations: author-supplied information remains pending.
3	Data identification and access	Official Results Tool, download date, query identifiers and cited GBD methods.
4	Eligibility and exclusions	Exact outcome filters and completeness gates; no country removals from the main set. Sensitivity exclusions are explicit.

Item	Topic	Location or applicability
5	Source characteristics	GBD model outputs, dimensional mappings and acquisition audit. Original source-study characteristics are documented by GBD.
6	Potential input biases	Discussion: sparse evidence, diagnostic heterogeneity, attribution and modeled denominators.
7	Other inputs	Matching GBD population and published standard weights with provenance.
8	Input data access	CSVs retained in the research archive; source institution IHME and exact queries identified for reproducibility.
9	Conceptual overview	Equations for within-age pooling, standardization, contrasts and decomposition.
10	Analytical detail	Normalization, validation, comparison, decomposition, trend and sensitivity methods and scripts.
11	Candidate evaluation and selection	Segmented-model BIC, training AICc and explicit forecast-retention criterion.
12	Evaluation results	Reference, endpoint, AR(1), rounding and forecast-validation results.
13	Uncertainty methods and scope	Joint draws unavailable; no propagated derived intervals; conditional trend interval scope stated.
14	Source code access	GitHub repository, pinned dependencies and executable scripts.
15	Machine-readable estimates	Complete derived CSV tables accompany the paper.
16	Quantitative uncertainty	Incomplete for derived estimates: marginal source limits retained but cannot substitute for joint uncertainty. Trend intervals are conditional only.
17	Interpretation and prior evidence	Discussion compares prior GBD studies. Release-to-release differences were not estimated.
18	Limitations	Discussion: endpoint, reference, denominator, source, rounding and inference limitations.




Figure S1: Sensitivity of the 2023 prevalence difference to reference-country exclusions, fixed 1990 country shares within ages, and restriction to ages 20–49. Dashed lines show the main 15–49 comparison. PCOS signs are stable; endometriosis signs vary. All are point estimates.




Figure S2: Restricted-age standardized YLD rates for both infertility causes, 1990–2023. These are infertility-impairment YLD rates rather than the total disability burden of either underlying disease.




Figure S3: Country-specific prevalence rates in 2023, restricted to ages 15–49. Dashed lines are the pooled reference. Descriptive point estimates only.




Figure S4: Age contrasts and standard-weighted contributions to changes in China minus reference. Contributions are not case counts or causal effects.




Figure S5: China's case-count decomposition and same-population benchmarks. The latter are signed arithmetic contrasts using China's actual age distribution, not preventable cases.