Reference composition and outcome definitions in PCOS- and endometriosis-related infertility comparisons: GBD 2023 and US survey analyses

Author names, institutional addresses and corresponding author
pending research-team completion

25 October 2026

Running title: Interpreting infertility comparisons

Extended abstract

Study question: How do reference composition, infertility subtype and outcome/denominator definitions affect the interpretation of PCOS and endometriosis infertility comparisons?

Summary answer: Reference-country shares explained a substantial part of the modeled China–G20 PCOS contrast change, while a separate survey showed that diagnosis-associated estimates varied with definitions and denominators.

What is known already: GBD studies describe infertility burden, trends and subtypes. Age standardization does not fix country composition within a pooled reference, and survey definitions are not interchangeable.

Study design, size, duration: Ecological secondary analysis of GBD 2023, 1990–2023, covering 19 sovereign G20 countries and seven female age groups, plus a post hoc cross-sectional NSFG 2022–2023 analysis with 4,855 respondents aged 20–49 at interview.

Participants/materials, setting, methods: China was compared with 18 other countries using pooling within age, restricted-age standardization, subtype accounting and symmetric endpoint decomposition, with annual-chained sensitivity. NSFG analysis used final weights, 20 strata and 80 clusters, preserving joint Taylor covariance. Independent R and Python implementations checked estimates and covariance.

Main results and the role of chance: The PCOS-related infertility contrast narrowed by 151.54 per 100,000; reference composition contributed 88.65 (58.5%), compared with 87.11 (57.5%) under annual chaining. Secondary infertility accounted for 69.4%. These GBD quantities lack propagated uncertainty. Survey prevalence ratios for broad fertility problems versus 12-month infertility in unions were 2.65 (95% design CI 2.30–3.04) versus 1.77 (1.22–2.57) for PCOS, and 1.78 (1.40–2.25) versus 0.84 (0.50–1.43) for endometriosis. The paired within-union PCOS definition shift in prevalence differences was 17.15 percentage points (10.24–24.06).

Limitations, reasons for caution: No jointly aligned GBD draws were obtained. Survey associations are unadjusted and partly sparse; the narrow endometriosis interval crosses one. Definitions and populations differ, and NSFG is not a proven held-out GBD validation sample. Several analyses were post hoc; endometriosis terminal-year instability is unresolved.

Wider implications of the findings: Comparisons should distinguish subtype, reference composition, definitions and denominators. Neither convergence nor definition-sensitive associations demonstrate improved care, disease causation or a protective endometriosis effect.

Study funding/disclosures: Actual human-author funding and disclosures remain to be verified; none are assumed.

Trial registration number: Not applicable; no interventional trial or external preregistration is claimed.

Key words: infertility; PCOS; endometriosis; reference population; survey design

1 Introduction

Infertility associated with polycystic ovarian syndrome (PCOS) and endometriosis is relevant to population reproductive health as well as individualized clinical care Teede et al. (2023); Becker et al. (2022). Estimates attributed to these causes within the Global Burden of Disease (GBD) infertility impairment describe a different outcome from total PCOS or endometriosis prevalence. They are also distinct from lifetime infertility or infertility measured exclusively among people currently trying to conceive GBD 2023 Collaborators (2025).

Much of the descriptive territory has already been covered. Xu and colleagues analyzed both underlying diseases and their attribution to total, primary and secondary infertility in GBD 2021 Xu et al. (2025). Lu and colleagues examined the same two infertility causes among women

aged 15–49 in China and globally, including projections Lu et al. (2026). Wang and colleagues subsequently analyzed infertility causes and primary/secondary attribution using GBD 2021 and 2023 Wang et al. (2026). Broad trend, decomposition and prediction analyses have also appeared in this journal Feng et al. (2025). Neither a newer GBD edition, subtype stratification alone, nor adding a G20 comparison constitutes a sufficient methodological advance.

A narrower unresolved question concerns the interpretation of an aggregate international contrast. A population-pooled reference combines country rates with country population shares that may change over time, even after age standardization. A narrowing difference could therefore reflect changing Chinese rates, changing reference-country rates, or changing reference-country composition. Primary and secondary infertility may contribute differently to each component. These are separable accounting questions; they do not identify causal mechanisms.

We compared China's cause-specific infertility estimates with a fixed set of 18 other sovereign G20 countries. We quantified subtype and age contributions to changes in the contrast and added an explicit decomposition of reference composition during manuscript review. We also assessed country and endpoint dependence. Endometriosis was retained as a contrasting cause, with terminal-year instability treated as a finding requiring caution rather than evidence of sustained convergence.

2 Materials and methods

2.1 Design, data and outcome definitions

This ecological secondary analysis used GBD 2023 Results Tool downloads obtained on 25 October 2026 IHME (2026). The analysis included females aged 15–49, seven five-year age groups and annual estimates for 1990–2023. China was compared with Argentina, Australia, Brazil, Canada, France, Germany, India, Indonesia, Italy, Japan, Republic of Korea, Mexico, Russian Federation, Saudi Arabia, South Africa, Türkiye, United Kingdom and United States of America. These are the other sovereign country members of the G20; regional organizations were excluded. The G20 is an economically defined comparison, not a matched control population or a global benchmark.

We selected infertility impairment 198, underlying causes endometriosis (607) and PCOS (605),

prevalence and years lived with disability (YLDs), and Number and Rate. Primary (220) and secondary (221) infertility were downloaded separately. Matched female populations came from the same release. Main analyses concern prevalence; YLDs, case-count decompositions, country distributions, trend models and failed forecast validation are secondary results in Additional file 1. The GBD component used no individual records; a separate post hoc analysis used deidentified NSFG public-use respondent records, as described below.

The GBD methods appendix describes survey-based primary infertility in couples without a previous live birth who desire a child, have been in a union for at least five years, and report no contraceptive use. Secondary infertility also uses the interval since the last live birth. Heterogeneous survey and clinical definitions are harmonized within GBD; the resulting estimates should not be equated with a uniform 12-month clinical definition. Underlying-cause allocation occurs within the modeled infertility envelope GBD 2023 Collaborators (2025). The denominator here is all females in each age group, not only couples attempting pregnancy.

2.2 Validation and standardization

We required complete dimensional coverage, unique cells, consistent cause and infertility identifiers, finite non-negative values and ordered original marginal uncertainty bounds. Number divided by matching population was compared with official Rate. Annual subtype rates were reconstructed from Number and matched population; independent original Number/Rate exports for 1990 and 2023 checked this reconstruction. Primary plus secondary point estimates were required to reproduce the total. Input hashes, mappings and retrieval specifications accompany the code PCOS research workspace (2026).

We used published GBD 2023 standard-population weights reproduced to six decimals, normalized over ages 15–49 Chen et al. (2026). The original machine-precision IHME weight file was not retrieved. Thus, all standardized rates are restricted-age rates. With normalized age weight w_a, country c, age a, year t, infertility type k, modeled cases C and female population P, rates per 100,000 were $r_{catk} = 10^5 C_{catk}/P_{cat}$. The age-specific reference was formed before standardization:

$$r_{Ratk} = \sum_{c \in R} s_{cat} r_{catk}, \quad s_{cat} = \frac{P_{cat}}{\sum_{j \in R} P_{jat}}, \quad A_{Ctk} = \sum_a w_a r_{Catk}, \quad A_{Rtk} = \sum_a w_a r_{Ratk}. \tag{1}$$

Here R always excludes China. The country-share weights s vary within age and year; the

standard age weights w are fixed. Rates were not constructed by averaging country standardized rates.

2.3 Subtype and reference-composition accounting

We calculated the signed difference $D_{tk} = A_{Ctk} - A_{Rtk}$ and ratio A_{Ctk}/A_{Rtk}. Because primary and secondary counts sum to total, their differences and changes also sum exactly. Their signed contributions are not attributable fractions or causal effects. Each age group's contribution is its standard weight multiplied by its age-specific difference change.

During manuscript review, we added a symmetric decomposition between start year 0 and end year T:

$$\Delta D_k = X_k + Q_k + S_k, \tag{2}$$

$$X_k = \sum_a w_a (r_{CaTk} - r_{Ca0k}), \tag{3}$$

$$Q_k = -\sum_a w_a \sum_{c \in R} \frac{s_{ca0} + s_{caT}}{2} (r_{caTk} - r_{ca0k}), \tag{4}$$

$$S_k = -\sum_a w_a \sum_{c \in R} \frac{r_{ca0k} + r_{caTk}}{2} (s_{caT} - s_{ca0}). \tag{5}$$

X is the Chinese rate component, Q the reference-country rate component and S the reference country-share component. The mean of the two replacement orders shares the product interaction symmetrically. Components sum exactly to the change in the standardized contrast, separately by subtype and age. A positive component moves China minus reference upward; with an initially negative contrast, it can indicate narrowing without indicating health improvement. This decomposition describes reference weighting, whereas the separate case-count decomposition concerns China's own population size and age distribution.

2.4 Sensitivity analyses and uncertainty

We excluded each reference country in turn, restricted ages to 20–49, and fixed within-age reference-country shares at their 1990 values. The symmetric decomposition was repeated for country exclusions, the restricted age range, periods starting in 1990 or 2010, and endpoints 2021, 2022 and 2023. Endpoint analyses were added after inspection of a terminal-year endometriosis increase; the reference-composition decomposition was added during manuscript review. Nei-

ther was externally preregistered. Fixed-share contrasts are a separate, order-specific sensitivity, not an interchangeable estimate of the symmetric share component.

We examined all 128 combinations of perturbing each unnormalized standard weight by half of its last published decimal. These deterministic rounding envelopes do not measure epidemiological uncertainty. Additional file 1 describes all secondary methods, including conditional trend intervals and forecast-validation rules.

Jointly aligned GBD draws across ages, countries and years were unavailable. We therefore report derived point estimates without propagated uncertainty intervals or significance tests. Original marginal limits are preserved but not summed or treated as standard errors. Country exclusions describe reference sensitivity, not sampling uncertainty. YLDs and prevalence share inputs and are not independent replications. Python 3.12 and pinned NumPy, pandas, SciPy, statsmodels and Matplotlib dependencies were used. Code and machine-readable outputs are available in Additional file 2 and the repository PCOS research workspace (2026). Applicable GATHER items, including incomplete propagated uncertainty reporting, are mapped in Additional file 1 Stevens et al. (2016).

2.5 Independent survey: definitions and denominators

We added a post hoc cross-sectional analysis of the 2022–2023 US National Survey of Family Growth (NSFG) public-use female respondent file NCHS (n.d.-a). This is a separate analysis, not a validation of Chinese GBD levels, China–G20 contrasts or GBD cause attribution. CDC's published estimates provided computational replication targets rather than new findings Nugent and Lochner (2026).

The original file contained 5,586 respondents. Our primary domain comprised 4,855 women aged 20–49 at interview. A sensitivity included one respondent aged 50 who had been 49 at screening according to the recode specification, yielding 4,856. Unknown diagnosis responses were excluded for each condition, not coded as no. Exposures were ever-reported professional diagnoses, not assigned causes of infertility.

Three estimands were distinguished: broad current fertility problems among all eligible women; that broad outcome among current opposite-sex married/cohabiting women; and 12-month infertility within the same union domain. Broad problems were FECUND categories 3–5 or INFERT=2. Surgical sterility (FECUND=1 or 2) was not classified as impaired fecundity. The

union domain used RMARITAL categories 1–2, and the narrower outcome INFERT=2 with valid INFERT categories 1–3. The recodes can reflect either partner's fertility and do not isolate a disease-specific cause Nugent and Chandra (2024); NCHS (n.d.-b).

Final weights WGT2022_2023, strata VEST and clusters VECL were used. All 80 clusters in 20 strata were retained, including zero domain contributions, giving 60 design degrees of freedom. Taylor-linearized covariance across 36 proportions preserved dependence between overlapping definitions. Pointwise 95% intervals used t_{60}, the log scale for ratios and the untransformed scale for differences. Paired changes in prevalence differences were evaluated in the same union domain. Analyses were exploratory, unadjusted and not multiple-comparison corrected. Narrow-outcome sparse-event counts were reported. R `survey` reconstructed proportions and the full covariance from original records using `svyratio` and a joint delta transformation of `svytotal`. The separate local Python calculation agreed to numerical precision; audit files record the comparison. This verifies computation under the design, not GBD validity.

2.6 Ethical considerations

Aggregate GBD estimates and deidentified public-use NSFG records were analysed without new recruitment or identifiable linkage. The research team must document the applicable institutional determination before submission; no approval or waiver is invented.

2.7 Use of artificial intelligence

OpenAI Codex was used for coding, execution assistance, source retrieval and manuscript drafting, including this revision. No artificial observations were substituted for the downloaded GBD data. Final human review of code, sources, interpretation and all author declarations remains necessary before submission. The tool is not an author and cannot approve the manuscript.

3 Results

3.1 Coverage and overall contrasts

The total burden export contained 36,176 rows and the population export 4,522 age–country–year cells. A further 36,176 annual subtype Number rows and 4,256 endpoint Number/Rate

rows covered both causes and measures. Primary plus secondary reproduced total point estimates throughout. Reconstructed subtype endpoint rates agreed with original rates to within 1.51×10^{-11} per 100,000. These checks establish extraction and arithmetic consistency, not independent validity of GBD estimates.

China's PCOS-related infertility standardized prevalence increased from 333.56 to 605.08 per 100,000, compared with 606.48 to 726.45 in the reference (Table 1; Figure 1). The difference changed from −272.92 to −121.38 and the ratio from 0.550 to 0.833. For endometriosis, rates changed from 63.70 to 53.30 in China and 66.63 to 53.51 in the reference. The 2023 total difference was −0.20; it was −8.76 in 2022. China's endometriosis estimate rose by 23.6% in the last year, so the terminal comparison does not summarize a stable trajectory.

Table 1: Total infertility prevalence in China and the reference population.

		Cases, thousands		Rate per 100,000			
Cause	Year	China	Reference	China	Reference	*D*	Ratio
Endometriosis	1990	192.8	375.6	63.70	66.63	-2.93	0.956
Endometriosis	2023	181.6	457.1	53.30	53.51	-0.20	0.996
PCOS	1990	1083.3	3446.2	333.56	606.48	-272.92	0.550
PCOS	2023	1866.7	6124.2	605.08	726.45	-121.38	0.833

Rates standardized within ages 15–49. Reference pools the other 18 sovereign G20 countries within age. *D*: China minus reference. All cells are modeled point estimates without propagated uncertainty.

3.2 Primary and secondary infertility

In 2023, the PCOS primary-infertility difference was −123.54 per 100,000 (ratio 0.441), whereas the secondary difference was +2.16 (ratio 1.004; Table 2; Figure 2). Their changes since 1990 were +46.40 and +105.14, respectively, summing to +151.54. Thus secondary infertility accounted for 69.4% of the signed total change. The share was 70.5% ending in 2021 and 69.0% ending in 2022. Across country exclusions and restriction to ages 20–49, the 2023 share ranged from 64.2% to 75.4%; fixing reference-country shares at 1990 yielded 78.1%. The exact percentage therefore depends on reference construction, while secondary infertility remained the larger positive component in these checks.

The 2023 PCOS primary difference remained negative under every country exclusion (−201.52 to −101.89), but the secondary difference ranged from −113.63 to +30.72. Near parity of secondary infertility was therefore a main-reference result, not a stable sign conclusion. For endometriosis,

the 2023 primary difference of −7.94 and secondary difference of +7.73 nearly canceled. Both subtype signs survived country exclusions in 2023, but the secondary difference was −0.40 in 2022. Its endpoint dependence limits the interpretation of cancellation.

Table 2: Primary and secondary infertility prevalence at the endpoints.

Cause	Type	Year	China rate	Ref. rate	*D*	Ratio
Endometriosis	Primary	1990	4.92	19.64	-14.72	0.250
Endometriosis	Primary	2023	4.94	12.87	-7.94	0.384
Endometriosis	Secondary	1990	58.78	46.98	11.79	1.251
Endometriosis	Secondary	2023	48.37	40.63	7.73	1.190
PCOS	Primary	1990	44.70	214.64	-169.94	0.208
PCOS	Primary	2023	97.35	220.89	-123.54	0.441
PCOS	Secondary	1990	288.86	391.84	-102.98	0.737
PCOS	Secondary	2023	507.72	505.56	2.16	1.004

Rate units: per 100,000 females; standardization restricted to ages 15–49. Point estimates only. Annual rates were reconstructed from Number and same-release population, with original endpoint Rate checks. Types sum to total before displayed rounding.

3.3 Changing reference composition

For PCOS, Chinese rate changes contributed +271.51 per 100,000 to the changing contrast, reference-country rate changes −208.62, and reference-country share changes +88.65 (Table 3; Figure 3). An additional post hoc annual-chained decomposition gave a share component of +87.11 (57.5%), compared with the endpoint value +88.65 (58.5%). This is a deterministic time-path sensitivity, not uncertainty estimation. The endpoint share component accounted for 58.5% of the +151.54 signed net change; the combined rate components accounted for +62.89. The corresponding share components were +88.11 ending in 2021 and +86.74 ending in 2022. Across country exclusions and the age-range sensitivity, the 2023 composition share ranged from 39.8% to 67.9%. A material reference-composition contribution persisted, although a majority contribution did not hold in every specification.

For primary PCOS infertility, the three components were +52.65, −38.09 and +31.84; for secondary infertility, they were +218.86, −170.53 and +56.81. Thus subtype and reference composition are intersecting dimensions of the same contrast change: the 69.4% subtype share and 58.5% composition share must not be added. Fixing all reference-country shares at 1990 instead yielded a 2023 total difference of −202.48, compared with −121.38 under current pooling. This

fixed-share calculation follows one replacement path and therefore differs from the symmetric component.

Endometriosis's +2.73 net difference change combined Chinese rate changes of -10.39, reference-country rate changes of +13.50 and reference composition of -0.38. Its near-zero net change makes percentage contributions unstable; we report absolute components. Age-by-type component tables for each endpoint accompany Additional file 2.

Table 3: Symmetric decomposition of changes in standardized differences, 1990–2023.

Cause	Type	Net change	China rates	Ref. rates	Ref. shares
Endometriosis	Primary	+6.79	+0.02	+6.69	+0.08
Endometriosis	Secondary	-4.06	-10.41	+6.81	-0.46
Endometriosis	Total	+2.73	-10.39	+13.50	-0.38
PCOS	Primary	+46.40	+52.65	-38.09	+31.84
PCOS	Secondary	+105.14	+218.86	-170.53	+56.81
PCOS	Total	+151.54	+271.51	-208.62	+88.65

All components use per 100,000 females and sum to net change. Reference shares are country population shares within each age. Positive values move China minus reference upward. Percentages of a signed net change are accounting summaries, not causal fractions.

3.4 Age patterns and secondary analyses

Ages 35–39 and 40–44 contributed +31.07 and +33.60 per 100,000, respectively, to the PCOS difference change, together 42.7% of the total. These are standard-weighted contrast contributions, not case counts. The age-by-type table separates Chinese rates, reference rates and reference composition within each cell.

China's PCOS-related infertility count rose by 783,425. A separate case-count decomposition assigned +860,721 to age-specific prevalence, offset by $-68,426$ from female population size and $-8,869$ from age distribution within ages 15–49. This is consistent with increasing modeled prevalence despite demographic offsets; it does not identify its causes. The total PCOS difference remained negative across reference exclusions. The endometriosis total difference changed sign across country exclusions. Rounding the published standard weights within their evaluated precision did not reverse the main 2023 signs. Secondary YLD, trend and benchmark results are retained in Additional file 1. Forecast candidates failed the comparative validation rule, so no long-range forecasts are reported.

3.5 Independent survey: definition and denominator sensitivity

The two diagnosis proportions, four broad-outcome percentages and two published diagnosis standard errors were reproduced at CDC's published precision. These were already published, not novel epidemiological discoveries. Complete-case fertility-service estimates did not all reproduce the report's rounding; the discrepancy was retained rather than forced to agree.

For PCOS, broad fertility problems occurred in 45.16% of diagnosed versus 17.07% of non-diagnosed women: prevalence ratio 2.65 (95% design CI 2.30–3.04). Restricting the broad outcome to current unions gave 2.17 (1.77–2.66); using 12-month infertility within unions gave 1.77 (1.22–2.57). Endometriosis ratios were 1.78 (1.40–2.25), 1.29 (0.95–1.76) and 0.84 (0.50–1.43), respectively (Table 4; Figure 4). The narrow endometriosis interval included one and does not establish protection.

Within the same union domain, the paired broad-minus-12-month difference in diagnosed–non-diagnosed prevalence differences was 17.15 percentage points for PCOS (95% CI 10.24–24.06), and 7.34 for endometriosis (−1.44 to 16.11). The 12-month diagnosed groups had only 29 PCOS and 15 endometriosis events. The interview-age sensitivity did not change the rounded core replication targets or qualitative interpretation.

Table 4: Independent NSFG survey estimates under different outcome and denominator definitions.

Diagnosis	Outcome, domain	Diagnosed (%)	Other (%)	Ratio (95% CI)
PCOS	Broad, all	45.16	17.07	2.65 (2.30, 3.04)
PCOS	Broad, union	41.98	19.32	2.17 (1.77, 2.66)
PCOS	12-month, union	12.63	7.12	1.77 (1.22, 2.57)
Endometriosis	Broad, all	33.00	18.59	1.78 (1.40, 2.25)
Endometriosis	Broad, union	27.11	20.98	1.29 (0.95, 1.76)
Endometriosis	12-month, union	6.49	7.70	0.84 (0.50, 1.43)

Ages 20–49 at interview. Other means no reported diagnosis, not verified absence of disease. Broad denotes impaired fecundity or 12-month infertility. Union denotes current opposite-sex marriage/cohabitation. Intervals are pointwise survey design-based, not GBD uncertainty intervals. Unadjusted exploratory comparisons.

4 Discussion

The principal finding is that subtype composition and reference population composition both affect the interpretation of international infertility contrasts. China's negative PCOS-related

infertility difference narrowed, but a substantial part of that change corresponded to population redistribution among reference countries. At the same time, secondary infertility accounted for most of the total contrast change, while a primary-infertility difference persisted. These results concern how an international comparison is constructed and what its summary conceals. The independent survey additionally demonstrates why outcome and denominator definitions must be aligned before validation is claimed.

This is an incremental contribution to an extensively studied field. Previous work already examined these causes, their primary and secondary infertility attribution, China/global patterns and standard trend and decomposition methods Xu et al. (2025); Lu et al. (2026); Wang et al. (2026); Feng et al. (2025). Our contribution is the joint accounting of subtype and within-age reference-country shares, with sensitivity to country inclusion and endpoint. We do not propose a new decomposition method or claim that GBD 2023 or subtype stratification alone is novel. The value is an interpretable application that changes the substantive reading of apparent convergence.

Age standardization removes changes in the chosen age weights, but it does not fix the mix of countries within an age group. When countries with different infertility rates gain or lose population share, a pooled reference can move even with unchanged country rates. The symmetric identity quantifies this component separately from the rate component and avoids assigning the entire rate–share interaction to one replacement order. The fixed-1990-share sensitivity gives a complementary contrast under a different reference definition. Neither is a preferred target rate or a causal counterfactual.

The subtype results further discourage describing narrowing as improved reproductive health. Both Chinese and reference PCOS rates increased. Secondary infertility's larger contribution does not show that individual women moved from primary to secondary infertility, nor that parity history caused the increase. The GBD categories reflect modeled estimates and survey definitions; diagnosis, fertility intentions, exposure to pregnancy, treatment and underlying disease attribution could all affect interpretation, but were not directly measured. The all-female denominator cannot estimate the risk among couples trying to conceive.

For endometriosis, the final-year near cancellation should be viewed as an unstable contrast. Independent download checks reproduced the source values, yet these checks cannot determine whether the abrupt terminal-year change reflects new evidence, model behavior or an actual

epidemiological shift. Analyses ending in 2021 or 2022 gave a materially different overall picture. This unresolved source-level issue warrants investigation with independent evidence before a substantive convergence claim. Retaining the earlier endpoints and full annual trajectories makes the limitation visible.

Case-count accounting answers a different public-health question. The PCOS increase occurred despite offsets from China's reproductive-age female population size and age composition. The age-specific prevalence component includes any changes embedded in the modeled rates; calling it an epidemiological component does not isolate biological risk, diagnosis or policy. Clinical guidelines support individualized reproductive and broader health assessment Teede et al. (2023); Becker et al. (2022); these ecological comparisons cannot select interventions, estimate treatment benefits or demonstrate service inequity. They can identify patterns that surveillance and independent studies should examine.

The independent US survey supports a measurement argument, not validation of Chinese GBD estimates. Association magnitudes changed when the denominator was restricted and the outcome narrowed. Joint survey covariance avoids incorrectly treating estimates from the same respondents as independent. PCOS retained a positive narrow-outcome association; endometriosis did not show a precise protective or harmful narrow-outcome association. Age, diagnostic access, previous treatment, surgical sterilization, parity and selection into ongoing unions remain uncontrolled explanations. These are descriptive estimand changes, not causal effects.

The two data sources cannot be directly pooled or calibrated: NSFG reports diagnostic history and fertility problems in US households; GBD models cause-specific infertility in all-female population denominators. We have not established whether NSFG evidence contributed to upstream GBD modeling. Independent therefore means separately acquired records and independently coded calculations, not a proven held-out GBD validation sample. The survey does not corroborate the Chinese endometriosis terminal-year jump or numerical GBD composition shares.

The principal GBD limitation remains unavailable joint uncertainty draws. We cannot propagate GBD uncertainty through pooling, standardization, subtype shares or the new decomposition. A 58.5% or 69.4% point estimate should therefore not be read as a precisely established population parameter. Country exclusions test dependence on the reference; they cannot replace draw-based

intervals. Bootstrap resampling of the deliberately selected G20 countries would answer another question and would not recover GBD uncertainty. Conditional trend intervals are confined to the supplement and do not establish cross-country differences.

Other limitations include sparse or heterogeneous upstream evidence, modeled cause allocation, the distinction between impairment and disease prevalence, the economically selected comparison group, and the absence of definitionally matched external validation of GBD. The US survey is a separate measurement-sensitivity analysis. Standard weights were available only as a rounded published reproduction. Several analyses were added after inspection, including endpoint checks and the reference-composition decomposition; their exploratory status requires transparent reporting and future corroboration. No release-to-release comparison was performed, so differences from previous publications cannot be assigned to real changes rather than GBD revision. These constraints limit the strength of inference even after formatting and numerical checks.

5 Conclusions

Within GBD 2023 point estimates, China's apparent convergence toward other G20 countries for PCOS-related infertility partly reflected changing reference-country population shares and concealed contrasting primary and secondary infertility patterns. Endometriosis near parity in 2023 depended strongly on the terminal year. International summaries should distinguish subtype, age and reference composition; these results require propagated uncertainty and independent corroboration before causal or policy conclusions.

List of abbreviations

AAPC, average annual percentage change; ARIMA, autoregressive integrated moving average; ASPR, age-standardized prevalence rate restricted here to ages 15–49; GATHER, Guidelines for Accurate and Transparent Health Estimates Reporting; GBD, Global Burden of Disease; G20, Group of Twenty; IHME, Institute for Health Metrics and Evaluation; MAE, mean absolute error; NSFG, National Survey of Family Growth; OLS, ordinary least squares; PCOS, polycystic ovarian syndrome; YLD, years lived with disability.

Authors' roles

Pending the actual human author list and verified individual contributions. All-author approval remains to be documented.

Acknowledgements

We acknowledge the GBD data resource and the contributions of its data providers and collaborators. Any additional named acknowledgements and required permissions remain for the human authors to verify.

Funding

Pending author-provided funding sources, grant numbers and funder roles. No funding status has been assumed.

Disclosures

Pending verified declarations from all human authors. An absence of competing interests has not been assumed.

Availability of data and materials

The derived datasets supporting the reported results and their analysis code are provided in Additional file 2 and the public GitHub repository PCOS research workspace (2026). Original GBD inputs are obtained from IHME through the Results Tool IHME (2026), subject to its data-use terms and access requirements. Exact dimensions, identifiers, download date, hashes and acquisition specifications are supplied. The original CSVs are retained in the research archive; they are not included in the public additional file. Re-running the full source-to-result pipeline requires obtaining these IHME inputs. The repository contains version history; the submission manifest records the analysis revision. A permanent data/code archive identifier remains to be finalized before submission.

Ethics approval and consent to participate

This study analyzed aggregate GBD estimates and deidentified public-use NSFG records, without access to identifiable records or recruitment of participants. The responsible authors must document the applicable institutional determination before submission. No institutional ethics approval or waiver has been invented for this draft.

Consent for publication

Not applicable: the manuscript contains no individual person's data, images or identifiable details.

Additional files

Additional file 1 (PDF, with editable TeX source). Supplementary methods, secondary case-count and trend analyses, endpoint checks, forecast validation, acquisition audit and GATHER reporting map; supplementary tables and figures.

Additional file 2 (ZIP). Machine-readable derived result tables, executable Python source, configuration, dependencies, acquisition and validation manifests, and a README specifying licensed IHME inputs needed to reproduce the analysis. No synthetic research observations or raw licensed downloads are included; the additional survey outputs are aggregate estimates and design covariance.

References

Becker CM, Bokor A, Heikinheimo O, Horne A, Jansen F, Kiesel L, King K, Kvaskoff M, Nap A, Petersen K, et al. ESHRE guideline: endometriosis. *Hum Reprod Open*. 2022;2022:hoac009. doi:10.1093/hropen/hoac009.

Chen S, Yang X, Jiang S, Lin Y, Huang L, Chen H, Zheng J, Xie Y, Kuang Z. Burden and temporal trends of female-specific cancers in China: a systematic analysis of the 2023 global burden of disease study. *PLOS One*. 2026;21:e0351539. S1 Table: reproduced GBD 2023 standard population. doi:10.1371/journal.pone.0351539.

Feng J, Wu Q, Liang Y, Liang Y, Bin Q. Epidemiological characteristics of infertility, 1990–2021, and 15-year

forecasts: an analysis based on the global burden of disease study 2021. *Reprod Health*. 2025;22:26. doi:10.1186/s12978-025-01966-7.

GBD 2023 Collaborators. Burden of 375 diseases and injuries, risk-attributable burden of 88 risk factors, and healthy life expectancy in 204 countries and territories, including 660 subnational locations, 1990–2023: a systematic analysis for the Global Burden of Disease Study 2023. *Lancet*. 2025. doi:10.1016/S0140-6736(25)01637-X. Includes Supplementary Appendix 1, infertility methods.

Institute for Health Metrics and Evaluation. GBD Results Tool. `https://vizhub.healthdata.org/gbd-results/`. Accessed 25 October 2026.

Lu S, Zhang P, Qian H, Li HM. Analysis and projection of the disease burden of endometriosis and polycystic ovary syndrome-related infertility among women aged 15–49 years in China and globally from 1990 to 2021. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2026;47:1091–1098. In Chinese. doi:10.3760/cma.j.cn112338-20250929-00691.

National Center for Health Statistics. 2022–2023 National Survey of Family Growth public-use female respondent data and documentation. `https://www.cdc.gov/nchs/nsfg/nsfg-2022-2023-puf.htm`. Accessed 25 October 2026.

National Center for Health Statistics. 2022–2023 NSFG female respondent recode specifications. `https://www.cdc.gov/nchs/data/nsfg/recodespecs/2022-2023-NSFG-FemResp-RecodeSpecs.pdf`. Accessed 25 October 2026.

Nugent CN, Chandra A. Infertility and impaired fecundity in women and men in the United States, 2015–2019. *Natl Health Stat Rep*. 2024;(202):1–19. doi:10.15620/cdc/147886.

Nugent CN, Lochner KA. Endometriosis, polycystic ovary syndrome, and uterine fibroids in women ages 20–49: United States, 2022–2023. *NCHS Health E-Stat*. 2026;(121):1–5. doi:10.15620/cdc/252459.

PCOS research workspace. Analysis code, query manifests and derived results for GBD 2023 infertility comparisons. GitHub. `https://github.com/jiabao1314/PCOS`. Accessed 25 October 2026. Exact analysis revision is recorded in the accompanying submission manifest.

Stevens GA, Alkema L, Black RE, Boerma JT, Collins GS, Ezzati M, Grove JT, Hogan DR, Hogan MC, Horton R, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *PLOS Med*. 2016;13:e1002056. doi:10.1371/journal.pmed.1002056.

Teede HJ, Tay CT, Laven JJE, Dokras A, Moran LJ, Piltonen TT, Costello MF, Boivin J, Redman LM, Boyle JA, et al. Recommendations from the 2023 International Evidence-based Guideline for the Assessment and Management of Polycystic Ovary Syndrome. *J Clin Endocrinol Metab*. 2023;108:2447–2469. doi:10.1210/clinem/dgad463.

Wang J, Gan C, Jia Z, Zheng Z, Zhang Y, Liang H. Global estimates of infertility burden and contributions of congenital, endocrine, metabolic and infectious factors. *Mol Med*. 2026;32:192. doi:10.1186/s10020-026-01548-8.

Xu Y, Jia Y, Li Q, Jiang H, Xiong Y, Liang W, Liu X, Huang Y, Zou K, Sun X, et al. Global, regional, and national burden of endometriosis, PCOS, and unexplained infertility and their attribution to infertility, 1990–2021: Global Burden of Disease Study 2021. *J Evid Based Med*. 2025;18:e70100. doi:10.1111/jebm.70100.

Figure legends




Figure 1: Annual total infertility rates and China–reference contrasts. Rates are standardized within ages 15–49. The reference pools population and modeled cases within age across 18 countries. Point trajectories have no propagated GBD uncertainty. The terminal-year endometriosis change is a source-estimate feature requiring independent corroboration.

Alt text: Annual rates and China-minus-reference contrasts. PCOS rates rise while the negative contrast narrows. Endometriosis is endpoint-sensitive.




Figure 2: Primary and secondary infertility trajectories. The same restricted-age standard and annual population-pooled reference are used for both types. Opposing subtype contrasts can cancel in the total. These modeled categories do not track transitions in individual women.

Alt text: Primary and secondary trajectories differ and can cancel in the total contrast.




Figure 3: Rate and reference-composition contributions to contrast changes. Contributions sum to the 1990–2023 change in China minus reference, separately by infertility type. Diamonds denote net changes. Positive components move the signed contrast upward. Reference-country shares vary within age; fixed standard age weights are used throughout. The decomposition was added during manuscript review and is descriptive.

Alt text: Signed rate and country-share components sum to contrast changes. Country composition contributes materially to PCOS.




Figure 4: US survey definition and denominator sensitivity. Prevalence ratios compare reported diagnosis with no reported diagnosis. Broad denotes impaired fecundity or 12-month infertility. Union denotes current opposite-sex marriage/cohabitation. Pointwise 95% design intervals use joint covariance and t_{60}, not GBD posterior draws. Unadjusted exploratory associations; narrow diagnosed events: PCOS 29, endometriosis 15. The narrow endometriosis interval crosses one and does not imply protection.

Alt text: Six survey prevalence ratios with intervals. The narrow endometriosis interval crosses one; no protective association is established.