Changing basis

Practice Quiz, 5 questions

5/5 points (100%)

Congratulations! You passed!

Next Item

1 / 1 point

1.

In this quiz, you will practice changing from the standard basis to a basis consisting of orthogonal vectors.

Given vectors $\mathbf{v} = \begin{bmatrix} 5 \\ -1 \end{bmatrix}$, $\mathbf{b_1} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\mathbf{b_2} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ all written in the standard basis, what is $\mathbf{v}$ in the basis defined by $\mathbf{b_1}$ and $\mathbf{b_2}$? You are given that $\mathbf{b_1}$ and $\mathbf{b_2}$ are orthogonal to each other.

- $\mathbf{v_b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$

Correct

The vector $\mathbf{v}$ is projected onto the two vectors $\mathbf{b_1}$ and $\mathbf{b_2}$.

---

1 / 1 point

2.

Changing basis

Practice Quiz, 5 questions

5/5 points (100%)

Given vectors $\mathbf{v} = \begin{bmatrix} 10 \\ -5 \end{bmatrix}$, $\mathbf{b_1} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$ and $\mathbf{b_2} = \begin{bmatrix} 4 \\ -3 \end{bmatrix}$ all written in the standard basis, what is $\mathbf{v}$ in the basis defined by $\mathbf{b_1}$ and $\mathbf{b_2}$? You are given that $\mathbf{b_1}$ and $\mathbf{b_2}$ are orthogonal to each other.

- $\mathbf{v_b} = \begin{bmatrix} 2 \\ 11 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} 11/5 \\ 2/5 \end{bmatrix}$
- ✓ $\mathbf{v_b} = \begin{bmatrix} 2/5 \\ 11/5 \end{bmatrix}$

Correct

The vector $\mathbf{v}$ is projected onto the two vectors $\mathbf{b_1}$ and $\mathbf{b_2}$.

- $\mathbf{v_b} = \begin{bmatrix} -2/5 \\ 11/5 \end{bmatrix}$

✓ 1 / 1 point

3.

Given vectors $\mathbf{v} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$, $\mathbf{b_1} = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$ and $\mathbf{b_2} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ all written in the standard basis, what is $\mathbf{v}$ in the basis defined by $\mathbf{b_1}$ and $\mathbf{b_2}$? You are given that $\mathbf{b_1}$ and $\mathbf{b_2}$ are orthogonal to each other.

- $\mathbf{v_b} = \begin{bmatrix} 5/4 \\ -5/2 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} -2/5 \\ 5/4 \end{bmatrix}$
- ✓ $\mathbf{v_b} = \begin{bmatrix} -2/5 \\ 4/5 \end{bmatrix}$

Correct

The vector $\mathbf{v}$ is projected onto the two vectors $\mathbf{b_1}$ and $\mathbf{b_2}$.

-

$\mathbf{v} = \begin{bmatrix} 2/5 \\ -4/5 \end{bmatrix}$

5/5 points (100%)

1 / 1 point

4.

Given vectors $\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, $\mathbf{b_1} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{b_2} = \begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}$ and $\mathbf{b_3} = \begin{bmatrix} -1 \\ 2 \\ -5 \end{bmatrix}$ all written in the standard basis, what is $\mathbf{v}$ in the basis defined by $\mathbf{b_1}$, $\mathbf{b_2}$ and $\mathbf{b_3}$? You are given that $\mathbf{b_1}$, $\mathbf{b_2}$ and $\mathbf{b_3}$ are all pairwise orthogonal to each other.

- $\mathbf{v_b} = \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} -3/5 \\ -1/3 \\ -2/15 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} -3/5 \\ -1/3 \\ 2/15 \end{bmatrix}$
- $\mathbf{v_b} = \begin{bmatrix} 3/5 \\ -1/3 \\ -2/15 \end{bmatrix}$

Correct

The vector $\mathbf{v}$ is projected onto the vectors $\mathbf{b_1}, \mathbf{b_2}$ and $\mathbf{b_3}$.

1 / 1 point

5.

Given vectors $\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 3 \end{bmatrix}$, $\mathbf{b_1} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$, $\mathbf{b_2} = \begin{bmatrix} 0 \\ 2 \\ -1 \\ 0 \end{bmatrix}$, $\mathbf{b_3} = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 0 \end{bmatrix}$ and $\mathbf{b_4} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 3 \end{bmatrix}$ all written in the standard basis, what is $\mathbf{v}$ in the basis defined by $\mathbf{b_1}$, $\mathbf{b_2}$, $\mathbf{b_3}$ and $\mathbf{b_4}$? You are given that $\mathbf{b_1}$, $\mathbf{b_2}$, $\mathbf{b_3}$ and $\mathbf{b_4}$ are all pairwise orthogonal to each other.

5/5 points (100%)

$\mathbf{v_b} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}$

$\mathbf{v_b} = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 1 \end{bmatrix}$

Correct

The vector $\mathbf{v}$ is projected onto the vectors $\mathbf{b_1}, \mathbf{b_2}, \mathbf{b_3}$ and $\mathbf{b_4}$.

$\mathbf{v_b} = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix}$

$\mathbf{v_b} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}$