Supplementary Information

Cloudra: Geometric Reasoning and Scientific Evidence in Autonomous Mechanism Exploration

Chengyi Yang[1], Qianxin Zhou[2,†], and Yang Li[1,†]

[1]College of Chemistry and Pharmaceutical Engineering, Changsha University of Science and Technology, Changsha 410114, China
[2]State Key Laboratory of Photocatalysis on Energy and Environment, College of Chemistry, Fuzhou University, Fuzhou 350002, China

[†]Qianxin Zhou and Yang Li contributed equally as co-second authors.

Draft — 31 October 2026

This supplement provides the experimental protocol, layered evaluation, complete task and score tables, expanded diagnostic analyses, literature source register, and machine-readable materials accompanying the manuscript. Standard denotes the same-model backbone comparison; Record and Brief are workflow ablations. Section, table, figure, equation, and page numbers carry the prefix S to distinguish them from the main text.

Contents

S1 Benchmark design and experimental protocol **S2**
S1.1 Task specification and study units . . . S2
S1.2 Workflow configurations and primary comparison . . . S2
S1.3 Allowed calculations and reference definitions . . . S3
S1.4 Physical evidence associated with a mechanism . . . S3
S1.5 Offline organization and publication records . . . S3
S1.6 Paired statistical analysis and time measurements . . . S4

S2 Layered scoring and scientific completion **S4**
S2.1 Composite and reporting layers . . . S4
S2.2 Graph structure and chemical correspondence . . . S4
S2.3 Graded physical self-consistency . . . S4
S2.4 Electronic barriers and numerical accuracy . . . S5
S2.5 Qualitative selectivity and property targets . . . S5
S2.6 Impossible tasks and final scientific claims . . . S5
S2.7 Original delivery, organized evaluation, and score reproduction . . . S6

S3 CloudraBench task catalogue **S6**

S4 Detailed benchmark results and scientific attribution **S9**
S4.1 Complete workflow and tier comparisons . . . S9
S4.2 Runtime delivery and the organized scientific record . . . S11
S4.3 Elapsed research time and accumulated calculation effort . . . S11
S4.4 Scientific-attribution cases . . . S12
S4.5 Impossible-tier task and claim resolution . . . S12
S4.6 Discussion of the supplementary comparisons . . . S13
S4.7 Complete per-task and per-configuration scores . . . S13
S4.8 Expanded task-score matrices . . . S17

S5 Behavioral diagnostics and archived research trajectories **S18**
S5.1 Study materials and local inference settings . . . S18
S5.2 Chemical identity under equivalent representations and answer reversal . . . S19
S5.3 XYZ coordinates and explicitly measured distances . . . S20
S5.4 Target-specific geometric feedback in an E1 operation chain . . . S20
S5.5 Single-token transfer of a reported numerical fact . . . S21
S5.6 Evidence associations under fixed local calculation facts . . . S21
S5.7 From a detected gap to the next research action . . . S21

S5.8 Discussion of the diagnostic scope S22

S6 Literature sample and source-linked capability register **S22**
S6.1 Selection and coding procedure S22
S6.2 Source register and counting variants S23

S7 Data, software, and reproduction materials **S25**
S7.1 Machine-readable task and score records S25
S7.2 Software responsibilities and calculation records S25
S7.3 Reproduction entry points S25
S7.4 Data-package inventory S25

S1 Benchmark design and experimental protocol

S1.1 Task specification and study units

CloudraBench comprises 50 tasks, distributed as 20 Easy, 10 Normal, 10 Hard, and 10 Impossible tasks. Easy tasks cover elementary transformations, conformational changes, and basic execution; Normal tasks require intermediate identification and continuity across elementary steps; Hard tasks emphasize competing pathways and stereochemical identity; Impossible tasks require conclusions appropriate to the capabilities of the allowed electronic-structure methods. A task fixes the reactant and target identities, reaction conditions, total charge, spin multiplicity, requested deliverables, and permitted methods. These specifications are common to the workflow configurations compared on that task. The complete task catalogue accompanies the per-task results.

Two studies provide complementary experimental units (Table S1). The repeated-task study contains three independent runs for each combination of four tasks and four configurations. E1 concerns elimination preorganization; NB1 concerns multistep decarboxylation; HA1 concerns stereochemical ring opening; and IA1 concerns an electronic-structure method limitation. The full-benchmark study contains one retained run for each of the 50 tasks under each configuration. Its primary matched unit is therefore a task represented in all four configurations. The 48 repeated-task runs and the 200 full-benchmark runs are reported as separate studies.

Table S1. Experimental units in the two studies.

Design	Repeated-task study (P1b)	Full-benchmark study (P2)
Task set	E1, NB1, HA1, IA1	All 50 CloudraBench tasks
Configurations	Guided, Standard, Record, Brief-errors	Guided, Standard, Record, Brief-errors
Repetitions	Three per task and configuration	One retained run per task and configuration
Task-level runs	$4 \times 4 \times 3 = 48$	$50 \times 4 = 200$
Elapsed-time unit	Four-task batch at concurrency two; three batches per configuration	Computational effort on the common 15-task subset

S1.2 Workflow configurations and primary comparison

Guided is compared primarily with the actual Standard branch using the same model backbone. Record and Brief-errors isolate changes to scientific acceptance and diagnostic presentation within the available workflow (Table S2). Standard retains access to computational tools while hiding explicit delivery-DAG tools, removing scientific acceptance interception, and providing neutral computational feedback. Guided retains the delivery graph, scientific checks, and detailed diagnostics. Record retains available tools and recording while removing mandatory scientific acceptance. Brief-errors retains guidance and acceptance while shortening diagnostics. The task and method restrictions are not relaxed in an ablation.

Table S2. Operational definition of the four workflow configurations. Brief-errors is abbreviated as Brief in figures and tables; Standard is also labelled Backbone.

Configuration	Delivery-graph interface	Mandatory scientific acceptance	Diagnostic presentation
Guided	Explicit	Retained	Detailed
Standard	Explicit DAG tools hidden	Removed	Neutral computation feedback
Record	Available tools and records retained	Removed	Recording retained
Brief-errors	Retained	Retained	Shortened

The repeated-run archive records the model name as DeepSeek-v4-flash.[1] Model names, campaign identifiers, budgets, stopping reasons, and artifacts are retained with the corresponding runs. Cloudra revision 70daa5a identifies the

implementation described in the main-text methods.[2] This implementation identifier, the experiment records, and the scoring/organization records serve different purposes: software or API versions are taken from the applicable archive rather than inferred from a later installed environment or model alias.

S1.3 Allowed calculations and reference definitions

Easy, Normal, and Hard tasks permit xTB, PM7, and B3LYP-D3(BJ)/def2-SVP.[3–10] Impossible tasks are restricted to xTB and PM7. Native ORCA, Gaussian, and MOPAC records supply the corresponding computational quantities and termination information.[11–14] Each result is associated with the engine, calculation level, species, and task conditions represented in its source record.

The reference answer (Gold) specifies the mechanistic species and elementary steps, stereochemical identities, applicable numerical targets, and explicit negative targets. Numerical activation barriers use electronic energies at a consistent level of theory:

$$\Delta E_e^{\ddagger} = E_{\mathrm{TS},e} - E_{\mathrm{ref},e}. \tag{S1}$$

These benchmark barriers do not include zero-point or thermal corrections. The energy reference is the verified stable minimum of the corresponding species. Declared conformational connections within the same species can share that stable reference; a conformational conversion itself retains its own starting conformer as the reference. Distinct stereochemical substrates and independent channel contact complexes retain separate references. The actual IRC connection and the energy denominator are recorded separately, so a stable reference is not automatically identified with the conformer reached directly by an IRC.

Reference treatment follows the individual task definition. NB1_E2 and NQ5_E2 have no numerical reference under the specified proton-transfer policy. IB9 combines a numerical first-step reference of 5.65 kcal mol^{-1} with a negative second-step target. The qualitative selectivity tasks and numerical tolerance settings are specified in Section S2. Reference status belongs to the corresponding reference record; task availability, presence of a Gold graph, and physical acceptance of a complete reference chain are recorded separately.

S1.4 Physical evidence associated with a mechanism

A delivered mechanism represents species occurrences as nodes and elementary steps as edges. A species minimum requires convergence and a frequency calculation with zero imaginary frequencies. An accessible transition-state step requires a first-order saddle with exactly one imaginary frequency associated with the intended reaction, together with direction evidence linking that saddle to the declared endpoint species.[15, 16] Native normal termination, geometry convergence, frequencies, and pathway connectivity are checked as separate facts.

The direction evidence starts at the same transition state in the two IRC directions. The actual outer frame from each direction is connected through a converged optimization to an endpoint minimum with zero imaginary frequencies and the appropriate chemical identity and stereochemistry. The IRC trajectory itself need not converge all the way to either minimum when this outer-frame–optimization–frequency chain establishes the connection. The archived frames and their calculation lineage are retained. Geometric interval measurements assist interpretation; they do not replace a supported native endpoint chain. For separated products, the required constituent minima are verified; an intact molecular complex is evaluated as the corresponding whole-system species.

These requirements apply to the objects requested by the task. An explicitly negative target is represented as inaccessible with its basis retained. A species-only negative question does not require an invented transition state, process edge, or numerical barrier.

S1.5 Offline organization and publication records

Formal evaluation is conducted after the run artifacts are organized into the common mechanism-graph representation. Organization uses that run's conversations, tool records, reports, geometries, and native calculation outputs to recover graph objects, repair file references, and convert quantities already present in the run. Human organization and field-level sources are recorded, and the same organizational rules apply across configurations. The resulting graph expresses the scientific content of the run; Gold is used for evaluation rather than as a source of additional run conclusions or computational evidence.

The full-benchmark publication table contains 200 runs and all 600 Graph, Self-consistency, and Accuracy component values. Each published composite is aligned with its own existing component records. The 200 released composite scores are retained; the weighted components reproduce them to their four-decimal reporting precision. The publication tables separately preserve whether the run itself generated a root mechanism graph and whether the graph was subsequently organized offline. Thus, the original delivery behavior and the organized scientific score remain separately inspectable quantities.

The repeated-task study retains its historical four-configuration snapshots. The subsequent Standard offline organization is a separately identified update to those records; it is not labelled as a new uniform rescore of all 48 runs. Analyses using the repeated-task archive specify which record supplies the result.

S1.6 Paired statistical analysis and time measurements

For a task i and configuration a, let S_{ia} denote its composite score. Configuration means give each task equal weight within the reported set. The primary paired difference is

$$\overline{\Delta}_{\mathrm{G-S}} = \frac{1}{n}\sum_{i=1}^{n}\left(S_{i,\mathrm{Guided}} - S_{i,\mathrm{Standard}}\right). \tag{S2}$$

The same calculation is used within each tier and for the combined Normal/Hard subset. Weighted component contributions are the mean paired component differences multiplied by 0.1, 0.3, and 0.6, respectively. They add to the composite difference up to the reporting precision of the released scores.

The 95% percentile intervals use 100,000 task-cluster bootstrap samples with seed 20261031. Each resampled task carries its four configuration records together, preserving the paired comparison. The bootstrap unit is a task, not an elementary edge or an individual quantum-chemical calculation.

The repeated-task elapsed-time analysis uses the recorded start and finish of each four-task batch at concurrency two: three batches per configuration, or 12 batches in total. Its unit is batch wall-clock hours. The full-benchmark effort analysis uses the 15 tasks with common effort records across all four configurations, comprising 60 task-level runs. It reports calculation counts and cumulative quantum-chemical job-hours, obtained by summing the individual calculation durations within a run. In the archived Standard/M1 extraction, native-log CPU time is divided by four to obtain the corresponding job-duration estimate. Job-hours are neither core-hours nor a measurement of the full P2 research-run wall-clock duration. These time quantities retain their distinct denominators and units throughout the supplementary tables.

S2 Layered scoring and scientific completion

S2.1 Composite and reporting layers

For a formally organized run, the reported composite combines graph structure G, physical self-consistency C, and accuracy A:

$$S = 0.1G + 0.3C + 0.6A. \tag{S3}$$

The three components are retained separately. Execution status, original graph delivery, completion of requested steps, graded physical evidence, and complete scientific acceptance are also reported separately. A completed process or a partial score does not substitute for the evidence required by a particular scientific claim.

S2.2 Graph structure and chemical correspondence

The graph component measures coverage of the reference mechanism by the delivered mechanism. Transition-state nodes are folded into their elementary-step edges for graph comparison. Species are matched chemically and reaction steps are compared through their endpoints and atom-mapped bond changes. RDKit supplies chemical representations and NetworkX supplies graph representation and matching.[17, 18] Missing reference elements, incorrect chemical identities, or mismatched accessibility declarations contribute edit costs. Additional delivered nodes or edges are not penalized merely for being additional in this component.

The tier-specific graph comparison ignores stereochemistry for Easy and Normal tasks and retains it for Hard tasks. Bound native evidence determines the evidence association of a delivered species. The graph score is not multiplied by the binary self-consistency pass flag: graph coverage and physical self-consistency remain distinct components.

S2.3 Graded physical self-consistency

The self-consistency assessment operates on delivered species and steps, independently of numerical agreement with Gold. A transition-state step receives graded credit for the evidence stages in Table S3. A complete verified chain receives 1.0. When necessary, evaluation searches the same run's native artifacts for the corresponding chain; recovery of a file association retains the same physical acceptance criteria.

Table S3. Graded self-consistency contribution for an accessible transition-state step. The endpoint contribution is 0.25 for one verified side and 0.50 for both sides.

Evidence stage	Contribution
Verified first-order saddle: exactly one imaginary frequency	0.35
IRC starting geometry consistent with the selected saddle	0.15
Native outer-frame–optimization–frequency lineage to both declared endpoint minima	0.50

For graphs containing reaction edges, C is the mean of the delivered edge scores. An edge without a transition-state node requires both endpoint species to be covered. For a species-only graph, C is the mean species-coverage score. Explicit inaccessible targets are exempt from requirements for physically nonexistent stationary points or paths and receive the corresponding self-consistency credit.

The complete self-consistency pass is a separate all-required-elements condition. It checks the required species and steps, including minimum evidence and the full transition-state/IRC chain. Consequently, a full graded component and a binary scientific-acceptance record are retained as separate outputs. Counts of $C = 1$ in the main text and supplementary score tables refer to full component scores; they are not renamed as counts of fully accepted complete mechanisms. Fractional scores identify the attained stages of physical support.

S2.4 Electronic barriers and numerical accuracy

For a numerical reference step, the declared barrier is checked against an energy difference derived from its native calculation evidence. An omitted edge barrier can be obtained from the delivered transition-state and starting-species energies when those quantities and their native records support the conversion. The energy difference is converted from hartree to kcal mol^{-1} using the common conversion factor 627.509474.

For a supported numerical value $\widehat{\Delta E}_e^{\ddagger}$ and reference $\Delta E_{e,\mathrm{Gold}}^{\ddagger}$, the underlying Gaussian score is

$$q_e = \exp\left[-\frac{\left(\widehat{\Delta E}_e^{\ddagger} - \Delta E_{e,\mathrm{Gold}}^{\ddagger}\right)^2}{2\sigma_e^2}\right]. \tag{S4}$$

The released task rubrics use base widths of 3.0 kcal mol^{-1} for Easy and Impossible tasks and 2.0 kcal mol^{-1} for Normal and Hard tasks. For a reference barrier at or below 10.0 kcal mol^{-1}, the reference-comparison width is the larger of the base width and 5.0 kcal mol^{-1}. This small-barrier width applies to comparison with Gold; the declaration-to-log consistency check retains the task's base width.

All reference edges remain in the accuracy denominator, with their recorded weights. A numerical reference edge lacking a matching delivered step, a declared barrier, or a log-derived value receives zero for that edge. Other evidenced steps retain their contributions. Numerical deviations on supported reference edges are aggregated with a maximum penalty fraction of 0.3; missing-step/evidence penalties and incorrect negative conclusions are not protected by this numerical-deviation cap. More explicitly, for total reference-edge weight W, supported numerically graded edges D, and missing or incorrectly delivered targets U, the aggregation takes the form

$$A = 1 - \min\left(\frac{\sum_{e\in D} w_e(1-q_e)}{W},\, 0.3\right) - \frac{\sum_{e\in U} w_e}{W}. \tag{S5}$$

Correct negative targets and accessible edges intentionally lacking a numerical reference contribute their prescribed full-credit terms through W. The latter are not interpreted as numerical barrier predictions.

The evaluator considers the legal transition-state/reference-energy combinations present in the delivered evidence. When those records contain a combination agreeing with Gold but the reported value uses a different supported conformer or energy-reference definition, the documented answer-definition treatment assigns an edge score of 0.9 in place of q_e. A declared barrier that is unsupported by any legal evidence combination, with no reference-consistent alternative in the delivered records, triggers the integrity short circuit: the composite retains only $0.1G$, and the self-consistency and accuracy contributions are zeroed. An omitted step is handled by its missing contribution rather than by this inconsistent-declaration rule.

S2.5 Qualitative selectivity and property targets

HA1, HA6, HB1, and HF1 use a qualitative favored-channel target in the accuracy slot. The delivered mechanism must identify one favored edge corresponding to the reference channel or its chemically matched product. Product identity supplies the chemical correspondence when the delivered channel label differs from the internal reference label. The numerical channel spread is not scored as a quantitative selectivity prediction in these tasks. HL1 instead retains its independently referenced channel barriers and does not use this favored-channel substitution.

When a task specifies species properties, its accuracy assessment compares the declared properties with the corresponding reference targets. Numerical property widths are 0.02 Å for bond lengths, 50 cm^{-1} for frequency quantities, and 2.0 kcal mol^{-1} for energy differences; the specific keyed property determines the width. Specified state-ordering text is compared after case and whitespace normalization. Multiple specified properties contribute their mean score.

S2.6 Impossible tasks and final scientific claims

Impossible-task assessment is based on the final delivered target. Correctly identifying that a requested target is inaccessible or outside the allowed method's capability earns the corresponding negative-target credit. An exploratory calculation followed by a correct negative conclusion is not penalized for having been attempted. An incorrect positive accessibility claim, a purportedly trustworthy barrier/property value for the negative target, or an absent required negative conclusion loses the corresponding credit. Negative endpoint species are treated consistently with their associated negative step. A species-only negative target is assessed without introducing a process edge.

The result must be interpreted at the identity of the requested target. A valid result for another electronic state, species, or pathway does not automatically answer the stated question. The per-task tables and case analyses retain this target association alongside the numerical score.

S2.7 Original delivery, organized evaluation, and score reproduction

Original delivery completeness records whether the run generated its mechanism graph during execution. Organized evaluation records the graph obtained by the formal offline procedure in Section S1. The publication component table preserves each run's original or organized score record and field provenance; existing released scores are reproduced rather than replaced by scores from a later software environment.

The retained reproduction procedure joins the released run table to the corresponding recorded components and reconstructs the aggregate tables and figure data. Its checks cover 200 unique task/configuration records, 600 nonmissing component values, and consistency of the weighted components with the 200 released totals. The maximum absolute difference from the published four-decimal composites is 4.9711×10^{-5}. The scoring formulas above explain the layered evaluation; regenerating publication tables requires the retained component records, while independently reassessing native scientific evidence additionally uses the associated run artifacts and evaluator revision.

S3 CloudraBench task catalogue

CloudraBench comprises 20 Easy, 10 Normal, 10 Hard, and 10 Impossible tasks. Tables S4–S7 list the chemical task, charge and multiplicity, stated conditions, and task-specific research target. The task names and molecular inputs are retained in the accompanying machine-readable catalogue; the original task definitions additionally specify atom mappings, complete prompts, reference graphs, and scoring settings.

Method set A comprises xTB, PM7, and B3LYP-D3(BJ)/def2-SVP and applies to Easy, Normal, and Hard tasks. Method set B comprises xTB and PM7 and applies to Impossible tasks. The research target is evaluated within its assigned method set. Reaction-mechanism tasks request a graph of the investigated species and elementary steps, electronic activation barriers for the steps actually computed, and bound stationary-point and pathway evidence. Stereochemical tasks additionally require identities established from the computed geometries. The rows below state the distinguishing target for each task rather than restating this shared delivery protocol.

For Impossible tasks, the listed quantity or process remains the question posed to the agent. The task evaluates the final conclusion under the allowed methods: an appropriately supported negative conclusion is a valid deliverable, and exploratory calculations need not be presented as trustworthy target properties. Species-property questions do not acquire invented process edges or transition states.

Table S4. Easy-tier task catalogue. Method set A applies to every task in this table.

ID	Chemical task	Stated conditions and task-specific research target
D1	butadiene + ethylene Diels-Alder (parent DA) Charge 0; multiplicity 1.	*Conditions:* gas phase, 487-648 degC kinetics. *Target:* Cycloaddition pathway, stationary points, and electronic activation barrier.
D3	cyclopentadiene dimerization (self Diels-Alder) Charge 0; multiplicity 1.	*Conditions:* gas/liquid phase, slow at room temperature. *Target:* Dimerization pathway, stationary points, and electronic activation barrier.
D4	cyclopentadiene + acrolein Diels-Alder Charge 0; multiplicity 1.	*Conditions:* gas phase or hydrocarbon solvent, heated. *Target:* Cycloaddition pathway, stationary points, and electronic activation barrier.
E1	F- + CH3CH2Cl elimination -> ethylene + HF + Cl- Charge -1; multiplicity 1.	*Conditions:* gas phase. *Target:* Elimination pathway from the pre-reaction complex, stationary points, and electronic activation barrier.
F1	BH4- + formaldehyde -> borane-methoxide adduct [CH3OBH3]- Charge -1; multiplicity 1.	*Conditions:* gas phase. *Target:* Hydride-transfer pathway to the borane–methoxide adduct, stationary points, and electronic barrier.
F4	BH3 + formaldehyde -> methoxyborane via Lewis adduct Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Lewis-adduct-mediated conversion to methoxyborane, with the investigated species and elementary steps.
G2	diimide cis reduction of ethylene Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Reduction pathway, stationary points, and electronic activation barrier.
G3	propylene + ethylene ene reaction (Alder-ene) Charge 0; multiplicity 1.	*Conditions:* gas phase, high temperature. *Target:* Ene-reaction pathway, stationary points, and electronic activation barrier.

Continued on the next page

Table S4 (continued)

ID	Chemical task	Stated conditions and task-specific research target
L1	cyclobutene -> butadiene ring opening Charge 0; multiplicity 1.	*Conditions:* gas phase, 403-448 K. *Target:* Ring-opening pathway, stationary points, and electronic activation barrier.
L2	hexatriene -> cyclohexadiene ring closure Charge 0; multiplicity 1.	*Conditions:* gas phase, 390-434 K. *Target:* Electrocyclic ring-closure pathway, stationary points, and electronic activation barrier.
M1	2,2'-dimethylbiphenyl aryl-aryl rotation Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Aryl–aryl rotational pathway, connected conformers, and rotational barrier.
M14	acrolein s-trans <-> s-cis rotation Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* s-trans/s-cis conformers, their connecting pathway, and rotational barrier.
M15	HONO cis-trans isomerization Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* cis/trans structures, their connecting pathway, and isomerization barrier.
M16	formic acid dehydrogenation HCOOH -> CO2 + H2 Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Dehydrogenation pathway and electronic activation barrier.
M17	methyl isocyanide -> acetonitrile rearrangement Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Isomerization pathway to acetonitrile and electronic activation barrier.
M3	NH3 umbrella inversion Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Inversion pathway, stationary points, and inversion barrier.
M8	cyclohexane chair flip Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Chair-interconversion pathway, relevant conformers, and barrier.
P2	cyclopentadiene degenerate H shift Charge 0; multiplicity 1.	*Conditions:* gas phase or solution, fast at room temperature. *Target:* Degenerate hydrogen-shift pathway, stationary points, and electronic activation barrier.
P3	allyl vinyl ether Claisen rearrangement Charge 0; multiplicity 1.	*Conditions:* gas phase, ~200 degC. *Target:* Claisen-rearrangement pathway, stationary points, and electronic activation barrier.
X1	beta-propiolactone -> ethylene + CO2 Charge 0; multiplicity 1.	*Conditions:* gas phase, ~350 degC. *Target:* Decomposition pathway to ethylene and carbon dioxide, with stationary points and barrier.

Table S5. Normal-tier task catalogue. Method set A applies to every task in this table.

ID	Chemical task	Stated conditions and task-specific research target
NB1	malonic acid thermal decarboxylation: malonic acid -> acetic acid + CO2 Charge 0; multiplicity 1.	*Conditions:* gas-phase thermolysis. *Target:* Identify intermediates and preserve elementary-step continuity through decarboxylation to the final products.
NI1	imine formation: formaldehyde + methylamine -> N-methylmethanimine + water Charge 0; multiplicity 1.	*Conditions:* gas phase; optimal around pH 4-6 (bell-shaped rate profile). *Target:* Identify the addition/condensation intermediates and connecting elementary steps leading to the imine and water.
NI5	imine formation (aqueous): formaldehyde + methylamine -> N-methylmethanimine + water Charge 0; multiplicity 1.	*Conditions:* aqueous solution, weakly acidic (pH 4-6, bell-shaped rate profile). *Target:* Resolve the aqueous imine-formation network and relevant proton-transfer steps under the stated conditions.
NQ1	aromatic Claisen rearrangement: allyl phenyl ether -> 2-allylphenol Charge 0; multiplicity 1.	*Conditions:* gas phase; thermal, ~200 degC. *Target:* Resolve rearrangement and subsequent conversion to 2-allylphenol, retaining the intermediate network.
NQ5	oxy-Cope rearrangement: 1,5-hexadien-3-ol -> hex-5-enal ([3,3] then enol-keto tautomerization) Charge 0; multiplicity 1.	*Conditions:* gas phase, heated. *Target:* Resolve the [3,3] rearrangement and enol–keto tautomerization as the investigated network.

Continued on the next page

Table S5 (continued)

ID	Chemical task	Stated conditions and task-specific research target
NR1	formamide dehydration: formamide -> hydrogen cyanide + water Charge 0; multiplicity 1.	*Conditions:* Gas phase; isolated neutral, closed-shell formamide; no added water, acid, or catalyst. *Target:* Investigate formimidic-acid intermediates and conformational transitions in the neutral dehydration network.
NU3	H3P=CH2 ylide + formaldehyde -> ethylene + H3P=O Charge 0; multiplicity 1.	*Conditions:* salt-free model conditions (gas-phase ylide, closest to model chemistry). *Target:* Resolve the salt-free model Wittig-reaction network to ethylene and phosphine oxide.
NU5	Passerini reaction: formaldehyde + formic acid + methyl isocyanide -> N-methyl-alpha-formyloxyacetamide Charge 0; multiplicity 1.	*Conditions:* dichloromethane (DCM), room temperature, high concentration. *Target:* Resolve the three-component Passerini-reaction network to the specified alpha-acyloxy amide.
NU6	Staudinger beta-lactam synthesis: ketene + methanimine -> 2-azetidinone Charge 0; multiplicity 1.	*Conditions:* ketene generated in situ; inert solvent. *Target:* Identify the ketene–imine addition and beta-lactam-forming network, retaining any identified intermediate.
NY3	isocyanic acid hydrolysis: HNCO + one water -> ammonia + carbon dioxide Charge 0; multiplicity 1.	*Conditions:* Gas phase; neutral closed-shell system with exactly one water molecule; no added solvent, acid, counterion, or catalyst. *Target:* Investigate carbamic-acid intermediates and connecting conformers without presupposing a fixed number of steps.

Table S6. Hard-tier task catalogue. Method set A applies to every task in this table.

ID	Chemical task	Stated conditions and task-specific research target
HA1	3-methylcyclobutene ring opening: determine the stereochemical outcome Charge 0; multiplicity 1.	*Conditions:* gas phase or inert solvent, ~150-200 degC, thermal. *Target:* Determine stereochemical products and channel identities; report channel barriers and the requested relative preference.
HA6	trans-3,4-dimethylcyclobutene ring opening: determine the stereochemical outcome Charge 0; multiplicity 1.	*Conditions:* gas phase or inert solvent, heating, thermal. *Target:* Determine stereochemical products and channel identities; report channel barriers and the requested relative preference.
HA8	(2E,4Z,6E)-octa-2,4,6-triene thermal 6-pi electrocyclization: determine the stereochemical outcome Charge 0; multiplicity 1.	*Conditions:* Gas phase; first thermal ground-state electrocyclization only; no subsequent hydrogen shifts or photochemistry. *Target:* Resolve the product stereochemistry of the first thermal electrocyclization; report investigated channel identities and barriers.
HB1	cyclopentadiene + maleic anhydride Diels-Alder: determine endo/exo outcome Charge 0; multiplicity 1.	*Conditions:* benzene, room temperature or mild heating. *Target:* Determine endo/exo products, channel barriers, and relative preference.
HB10	cyclopentadiene + acrylonitrile Diels-Alder reaction: determine both stereochemical pathways Charge 0; multiplicity 1.	*Conditions:* Gas phase; concerted addition; common reactant-complex reference for the two channels. *Target:* Compute both endo/exo pathways using a common reactant-complex reference; report identities and barriers.
HD2	cyclohexene oxide + hydroxide: determine the stereochemical outcome of the epoxide opening Charge -1; multiplicity 1.	*Conditions:* aqueous NaOH, heat. *Target:* Determine the stereochemistry of the epoxide-opening product from the computed structures.
HF1	cyclohex-2-en-1-ol peracid epoxidation: determine the stereochemical outcome Charge 0; multiplicity 1.	*Conditions:* performic acid, CH2Cl2, 0-25 degC. *Target:* Determine the two facial pathways, product stereochemistry, barriers, and relative preference.
HH1	1,1'-binaphthyl thermal racemization Charge 0; multiplicity 1.	*Conditions:* thermal racemization, ~50 degC, DMF. *Target:* Characterize the thermal racemization pathway, stereochemical endpoints, and barrier.

Continued on the next page

Table S6 (continued)

ID	Chemical task	Stated conditions and task-specific research target
HI1	Cope elimination of threo/erythro N,N-dimethyl-3-phenyl-2-butylamine oxide: determine the stereochemical mapping Charge 0; multiplicity 1.	*Conditions:* Gas phase; isolated neutral singlet amine oxide. Treat threo and erythro substrates as two separate systems and retain explicit stereochemical SMILES. *Target:* Compute threo and erythro substrates separately; establish substrate CIP assignments and alkene E/Z identity from 3D geometries. Use each substrate's own minimum for its barrier.
HL1	formonitrile oxide + norbornene [3+2] cycloaddition: determine both facial pathways Charge 0; multiplicity 1.	*Conditions:* Gas phase; neutral singlet formonitrile oxide and unsubstituted norbornene. Compare approach from both faces relative to the methano bridge. *Target:* Compute both facial approaches relative to the methano bridge and assign products from 3D geometries. Use each channel's own stable contact minimum; report contact-conformer energies separately.

Table S7. Impossible-tier task catalogue. Method set B applies to every task in this table.

ID	Chemical task	Stated conditions and task-specific research target
IA1	cyclobutadiene D4h automerization Charge 0; multiplicity 1.	*Conditions:* gas phase, low temperature (matrix isolation regime). *Target:* Thermal automerization of rectangular cyclobutadiene through the square transition structure: mechanism and activation barrier.
IA5	cyclobutene disrotatory ring opening Charge 0; multiplicity 1.	*Conditions:* gas phase, thermal. *Target:* Thermal disrotatory ring opening of cyclobutene to butadiene: mechanism and activation barrier.
IA7	Bergman cyclization of (Z)-hex-3-ene-1,5-diyne Charge 0; multiplicity 1.	*Conditions:* gas phase, thermal. *Target:* Bergman cyclization to p-benzyne: mechanism and activation barrier.
IA9	1,2-dioxetane thermolysis to two formaldehyde Charge 0; multiplicity 1.	*Conditions:* gas phase, thermal. *Target:* Thermolysis of 1,2-dioxetane to two formaldehyde molecules: mechanism and activation barrier.
IB9	diazomethane and ethylene to cyclopropane via 1-pyrazoline Charge 0; multiplicity 1.	*Conditions:* gas-phase thermolysis. *Target:* Conversion of diazomethane and ethylene to cyclopropane and nitrogen through 1-pyrazoline.
IC3	phenylnitrene open-shell singlet state Charge 0; multiplicity 3.	*Conditions:* gas phase. *Target:* Singlet–triplet energy gap and the electronic character of the lowest phenylnitrene singlet state.
ID2	anomalous bonding in dioxygen difluoride (FOOF) Charge 0; multiplicity 1.	*Conditions:* gas phase, low temperature. *Target:* Equilibrium F–O and O–O distances of FOOF and its F–O bond-dissociation energy.
IE2	nitrogen triple-bond dissociation curve Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Ground-state N2 dissociation curve and bond-dissociation energy.
IF3	triangulene triplet ground state and singlet-triplet gap Charge 0; multiplicity 3.	*Conditions:* gas phase (surface-synthesis analogue). *Target:* Ground-state spin multiplicity and singlet–triplet gap of triangulene.
IG8	chromium dimer ground-state dissociation curve Charge 0; multiplicity 1.	*Conditions:* gas phase. *Target:* Ground-state Cr2 potential-energy curve and bond-dissociation energy.

S4 Detailed benchmark results and scientific attribution

S4.1 Complete workflow and tier comparisons

The publication dataset contains 200 task runs: one retained run for each of 50 tasks under Guided, Standard, Record, and Brief. Each run has a composite score and all three component scores. Table S8 reports the means without pooling away the task tiers. Standard is the actual comparison branch using the same model backbone. Record and Brief vary the workflow's persistent acceptance requirements and diagnostic feedback, respectively.

Table S9 gives all six pairwise comparisons on the same 50 tasks. Guided–Standard has a mean difference of 0.186302, a median difference of 0.107950, and 35 task wins, eight losses, and seven ties. The 95% task-bootstrap interval is [0.100268, 0.275854]. Within the Normal tier, the means are 0.737020 and 0.573050, with a difference of 0.163970 [0.036520, 0.292230] and seven wins in ten tasks. Within Hard, the means are 0.957650 and 0.688750; the difference is 0.268900

Table S8. Composite and component means for each workflow and task tier.

Tier	Workflow	n	Composite	Graph	SC	Accuracy
All	Guided	50	0.7803	0.8788	0.9233	0.6924
	Standard	50	0.5940	0.7837	0.5303	0.5943
	Record	50	0.7044	0.7876	0.7177	0.6839
	Brief	50	0.7415	0.8430	0.8468	0.6720
Easy	Guided	20	0.8809	0.9620	0.9250	0.8454
	Standard	20	0.7305	0.9007	0.5858	0.7744
	Record	20	0.9443	0.9576	0.9050	0.9618
	Brief	20	0.9054	0.9523	0.9125	0.8940
Normal	Guided	10	0.7370	0.8589	0.9021	0.6342
	Standard	10	0.5731	0.7607	0.5217	0.5675
	Record	10	0.5287	0.6786	0.5783	0.4789
	Brief	10	0.5857	0.7472	0.7740	0.4646
Hard	Guided	10	0.9577	0.9107	0.9542	0.9672
	Standard	10	0.6887	0.8481	0.5033	0.7549
	Record	10	0.8723	0.9243	0.6650	0.9672
	Brief	10	0.8282	0.9286	0.7767	0.8372
Impossible	Guided	10	0.4451	0.7006	0.9100	0.1700
	Standard	10	0.2473	0.5083	0.4550	0.1000
	Record	10	0.2325	0.4200	0.5350	0.0500
	Brief	10	0.4829	0.6345	0.8583	0.2700

SC denotes the graded self-consistency component. Every mean is calculated from the complete publication table. The overall tier weights are 20/50 for Easy and 10/50 for each other tier.

[0.121300, 0.445930], with Guided higher in all ten task pairs. Easy has means of 0.880940 and 0.730485, with 11 wins, five losses, and four ties; its paired median difference is 0.020850. Impossible has means of 0.445060 and 0.247330 and is examined at the level of the requested target below.

Table S9. All pairwise composite-score comparisons across the 50 tasks.

Comparison	Mean difference	95% interval	W/L/T
Guided–Record	+0.0759	[+0.0121, +0.1438]	27/8/15
Guided–Standard	+0.1863	[+0.1003, +0.2759]	35/8/7
Guided–Brief	+0.0388	[-0.0317, +0.1076]	26/10/14
Record–Standard	+0.1104	[+0.0167, +0.2062]	28/13/9
Record–Brief	-0.0371	[-0.0999, +0.0258]	15/23/12
Standard–Brief	-0.1475	[-0.2474, -0.0494]	12/33/5

The difference is the first workflow minus the second. W/L/T gives the first workflow's task wins, losses, and ties. Percentile task-bootstrap intervals use 100,000 resamples and seed 20261031; the four configurations of each sampled task remain together.

The component decomposition in Table S10 expresses each contribution in composite-score units. Across all tasks, the self-consistency contribution is +0.117875 and the accuracy contribution +0.058916. In Normal and Hard together, the composite difference is +0.216435, including +0.124688 from self-consistency and +0.083712 from accuracy. Hard's accuracy contribution, +0.127391, accompanies a +0.135250 self-consistency contribution. These quantities describe the requested comparative study at two levels: the answer reached and the evidence supporting its attribution.

Table S10. Weighted decomposition of the Guided–Standard composite difference.

Scope	n	Composite	$0.1\Delta G$	$0.3\Delta SC$	$0.6\Delta A$
All	50	+0.186302	+0.009513	+0.117875	+0.058916
Easy	20	+0.150455	+0.006130	+0.101750	+0.042579
Normal	10	+0.163970	+0.009823	+0.114125	+0.040033
Hard	10	+0.268900	+0.006259	+0.135250	+0.127391
Impossible	10	+0.197730	+0.019226	+0.136500	+0.042000
Normal + Hard	20	+0.216435	+0.008041	+0.124688	+0.083712

Component contributions are calculated from the recorded unrounded components. Composite differences retain released composite scores, so the final displayed digits can differ by score rounding.

The ablations give additional resolution. Guided–Record averages +0.075892 [0.012134, 0.143818] across all tasks. In Normal and Hard together, its +0.146845 composite difference decomposes into +0.008338 from graph, +0.091938 from self-consistency, and +0.046574 from accuracy. Guided–Brief averages +0.038816 [-0.031704, 0.107648], with mean

differences of +0.151350 on Normal and +0.129470 on Hard. The task-level records retain both the general patterns and the crossed outcomes: NB1 has composite scores of 0.9628, 0.9128, and 0.9900 for Guided, Standard, and Brief; Guided and Brief both have self-consistency 1. NI1 has Guided and Standard composites of 0.9958 and 0.6196, with self-consistency 1 and 0.35. NQ5 favors Record over Guided, while NU3 favors Brief. Multistep performance is therefore resolved through the individual pathways and their evidence relationships.

S4.2 Runtime delivery and the organized scientific record

During execution, Guided, Record, and Brief each produced 50 root mechanism graphs. Standard produced 43. Its remaining seven studies—F1, HA1, HF1, IA9, M17, NR1, and P3—were organized offline from their own conversations, reports, tool results, and calculation artifacts. The resulting analysis contains 50 organized graphs per workflow and 200 complete score profiles. Offline organization follows the evaluation procedure described in the methods and retains the source of each recovered field.

The seven Standard studies have a mean organized composite of 0.153743, compared with 0.828529 for Guided on the same tasks. Their contribution is 50.7% of the total Guided–Standard score difference. The scientific content retained in a run and the production of its final root artifact are both visible in these records. The self-consistency distribution in Table S11 further resolves how much evidence was assembled for the 40 non-Impossible tasks. Guided has 24 full-score and 16 partial-score records; Standard has seven full-score, 29 partial-score, and four zero-score records.

Table S11. Self-consistency distribution over all 40 non-Impossible tasks.

Workflow	$SC = 1$	$0 < SC < 1$	$SC = 0$	Mean SC
Guided	24	16	0	0.9266
Standard	7	29	4	0.5492
Record	16	24	0	0.7633
Brief	20	20	0	0.8439

This table describes the graded component for the positive-chemistry tiers. The separate scientific-acceptance record is evaluated from the complete task contract.

S4.3 Elapsed research time and accumulated calculation effort

Table S12 preserves every batch observation from the 48-run repeated-task experiment. A batch contains the same four tasks at a concurrency of two, and its elapsed time spans the batch's actual start and finish. Guided's mean is 4.4514 h, Standard's 3.9577 h, Record's 3.8942 h, and Brief's 4.8273 h. Relative to Standard, Guided's observed mean is 12.5% longer; relative to Brief, it is 7.8% shorter. These observations include waiting, calculation execution, and continuation of the research after results return.

Table S12. All 12 elapsed-time observations in the repeated-task study, in hours.

Workflow	Batch 1	Batch 2	Batch 3	Mean
Guided	4.2383	3.8669	5.2489	4.4514
Standard	5.3381	2.5044	4.0306	3.9577
Record	5.4189	2.6203	3.6433	3.8942
Brief	7.2689	3.6475	3.5656	4.8273

Each observation is a four-task batch (E1, NB1, HA1, IA1) at concurrency two. Three batches per workflow contain 48 task runs in total. Batch completion times include the original research lifecycle and outcomes.

The P2 effort comparison uses the common 15-task set D3, M1, M8, NB1, NQ5, NR1, NU3, NU5, NU6, HA1, HA6, HB1, HB10, HD2, and HF1. This yields 60 run-level effort records, with all four configurations represented for each task. Table S13 gives the total and mean job-hours, job counts, terminal outcomes, and corresponding final scores. Recorded ledger durations are summed across quantum-chemical jobs. For Standard/M1, the archived effort reconstruction uses native-log CPU time divided by four, as specified in the source record. The measurement unit throughout this comparison is accumulated job-hours.

Table S13. Recorded quantum-chemical effort on the 15 tasks shared by all configurations.

Workflow	Job-h total	Job-h/task	Jobs	Failed	Cancelled	Mean C
Guided	166.19	11.079	789	114	94	0.8124
Standard	113.32	7.555	419	45	50	0.6495
Record	118.16	7.877	431	62	38	0.7360
Brief	233.85	15.590	883	167	120	0.7273

Each workflow contributes 15 task runs. Job-hours sum recorded process durations. Failed and cancelled jobs are separate terminal categories; job counts do not measure scientific acceptance.

Guided's mean of 11.0793 job-hours accompanies a mean composite of 0.812413. Relative to Standard, it uses 46.7% more accumulated job time and gains 0.162933 in mean composite score. Relative to Brief, it uses 28.9% less accumulated job time and gains 0.085080. On seven of the 15 paired tasks, Guided has both a higher score and lower accumulated job time than Brief. The job counts show how this expenditure was made: Guided records 789 jobs, including 114 failures and 94 cancellations; Brief records 883 jobs, including 167 failures and 120 cancellations. In the wider 63-run effort archive, NU3, NU5, NU6, and NR1 account for 163.5 of 673.82 job-hours across their 16 runs. Their trajectories contain repeated attempts to establish the required network and support examination of the concrete structures and pathways reached by those attempts.

S4.4 Scientific-attribution cases

Table S14 gives every score component for the cases below. Each comparison follows the chemical object to which the numerical or path evidence was attached.

Table S14. Complete score profiles for the four scientific-attribution cases.

Task	Workflow	Composite	Graph	SC	Accuracy
HA8	Guided	0.9625	1.0000	0.8750	1.0000
	Standard	0.6250	1.0000	0.3500	0.7000
HH1	Guided	0.9838	1.0000	1.0000	0.9731
	Standard	0.8338	1.0000	0.5000	0.9731
HI1	Guided	0.9999	0.9992	1.0000	1.0000
	Standard	0.7032	0.3325	0.2333	1.0000
NU6	Guided	0.6719	1.0000	0.9375	0.4844
	Standard	0.6892	0.8167	0.6250	0.7000

These profiles retain the distinct contributions of pathway coverage, evidence support, and agreement with the requested answer.

HA8: stable reactant reference and reactive conformer. Guided reports 29.46 kcal/mol and Standard 20.43 kcal/mol, against a reference of 29.48 kcal/mol. The 9.03 kcal/mol difference is traced to their energy references: the lowest all-s-trans reactant minimum in Guided and a reactive cisoid conformer in Standard. The stable-species energy reference specifies the activation-energy denominator, while the IRC identifies the local structures connected by a particular saddle. Maintaining both associations allows a path reached from a reactive conformer to be interpreted relative to the specified reactant species.

HH1: a reference-consistent number followed by evidence closure. Both workflows report a binaphthyl racemization barrier of 23.48 kcal/mol; the reference is 23.01 kcal/mol. Their accuracy components are 0.973067 and 0.973064 and both graph components are 1. Guided has self-consistency 1, whereas Standard has 0.5. Their composites are consequently 0.9838 and 0.8338. Standard's archive contains candidate channels and IRC inputs; Guided's delivered evidence completes the saddle-to-IRC-to-endpoint associations required by the step. The pair locates a 0.1500 composite difference after closely matching numerical answers have already been obtained.

HI1: numerical agreement and coverage of the supported study. Guided and Standard both have accuracy approximately 1. Their graph components are 0.999176 and 0.332509, and self-consistency components are 1 and 0.233333. The corresponding composites are 0.9999 and 0.7032. Here, evidence closure and delivered coverage jointly determine how much of the requested study accompanies the numerical answer.

NU6: the optimized saddle and the intended second step. The Guided record gives $E_{\mathrm{TS}} = -247.030042392$ Ha for its concerted saddle region and $E_{\mathrm{betaine}} = -247.060060699$ Ha. Using 627.5094740631 kcal mol^{-1} Ha^{-1} gives

$$[E_{\mathrm{TS}} - E_{\mathrm{betaine}}]\ 627.5094740631 = 18.836772\ \mathrm{kcal\,mol^{-1}}. \tag{S6}$$

Standard's archived second-step ring-closure value is 18.837 kcal/mol. Guided retains the target betaine-to-product ring-closure saddle as unresolved and describes the computed energy cost in relation to the concerted saddle region. Its attempts from interpolated betaine geometries and constrained-scan peaks repeatedly return toward the direct reactant-to-product saddle region. Following the optimized structure and its actual endpoints is therefore necessary to assign the energy difference to the step that was established.

S4.5 Impossible-tier task and claim resolution

Table S15 resolves Brief's mean composite of 0.482940 and Guided's 0.445060 into all ten task differences. Brief is higher on three tasks, Guided on five, and two tie. The Brief-minus-Guided mean is +0.037880, with the existing task-bootstrap interval $[-0.188220, 0.283970]$. Its weighted contributions are -0.006607 from graph, -0.015500 from self-consistency,

and +0.060000 from accuracy. IA1 and IA9 provide the two large positive differences, while IG8 has the opposite accuracy ordering. Across the seven tasks with equal accuracy components in these two workflows, Brief-minus-Guided averages -0.050000.

Table S15. Task-resolved comparison of Brief and Guided in the Impossible tier.

Task	Guided C	Brief C	Brief − Guided	Guided A	Brief A
IA1	0.1717	1.0000	+0.8283	0.0	1.0
IA5	0.4000	0.3667	-0.0333	0.0	0.0
IA7	0.2583	0.2583	+0.0000	0.0	0.0
IA9	0.3667	0.9333	+0.5666	0.0	1.0
IB9	0.8200	0.7700	-0.0500	0.7	0.7
IC3	0.4000	0.4000	+0.0000	0.0	0.0
ID2	0.3339	0.1839	-0.1500	0.0	0.0
IE2	0.3333	0.1833	-0.1500	0.0	0.0
IF3	0.3667	0.4000	+0.0333	0.0	0.0
IG8	1.0000	0.3339	-0.6661	1.0	0.0

C denotes the composite and A the accuracy component for the task-defined numerical or negative target. All values are retained from the publication score table.

Brief reaches accuracy 1 for IA1, the square-geometry cyclobutadiene automerization target, and IA9, dioxetane thermolysis. Guided reaches accuracy 1 for IG8, the chromium-dimer target. Both have accuracy 0.7 in IB9, which contains positive and negative target components. Across all 40 Impossible runs, four records have accuracy 1 and 14 have self-consistency 1 with accuracy 0. These component combinations are reported together with the identity of the requested state, channel, or property.

The preserved Brief/IC3 graph identifies the requested lowest open-shell singlet $^1A_2$ as inaccessible under the permitted route, marks its state-change edge accordingly, and withholds the requested singlet–triplet gap. It retains a separately characterized closed-shell $^1A_1$ branch. The PM7 value of +45.9 kcal/mol and GFN2-xTB value of -1.7 kcal/mol are described as method-dependent exploratory relative energies and are explicitly excluded from the requested gap. The graph records that the restricted calculation with a mixed initial guess does not test the requested unrestricted electronic state. The released profile remains Graph/SC/Accuracy = 1/1/0 and composite 0.4.

The Record/IA5 archive similarly distinguishes the requested disrotatory ring opening from an allowed conrotatory path. It marks the requested disrotatory edge inaccessible and records the positive conrotatory path separately at 33.57 kcal/mol. The archived adjudication documents that the matcher selected the positive path sharing the endpoints, leaving the accuracy component at 0. These records preserve both the delivered scientific claims and the values used in the publication analysis.

S4.6 Discussion of the supplementary comparisons

The Normal and Hard comparisons resolve the framework's contribution through relations that must survive across many local decisions: species and conformer identity, reaction-channel identity, the reference defining a reported energy, and the association of each result with the claim it supports. Generating a plausible next action from a current language context does not itself maintain this global state. A persistent evidence graph makes these accumulated relations available for subsequent reasoning and revision, complementing work on external planning and tool feedback.[19–21] The task cases further distinguish repairing an incomplete local proof from reconsidering which scientific question the current calculation can answer.

The P2 task-bootstrap intervals describe variation across the sampled tasks, since there is one retained run per task and configuration. The Impossible rubric targets the specified allowed-method contract; its negative-target self-consistency convention is distinct from a positive TS–IRC–endpoint chain. IA5 and IC3 show why score interpretation also requires channel and electronic-state identities. Their released scores remain unchanged. The Brief comparison is consistent with task-dependent effects of diagnostic content, while the final outcomes alone do not identify a causal effect of diagnostic length. The NU6 energy fingerprint identifies an assignment question; native geometry and pathway evidence are the independent basis for establishing whether two optimized structures represent the same saddle. Timing results retain their measured units: pilot batch elapsed time and P2 accumulated job-hours. Separating queueing, agent-active time, and end-to-end P2 latency is a further experimental question.

S4.7 Complete per-task and per-configuration scores

Tables S16–S19 report all 200 runs and all four reported quantities, comprising 200 composite scores and 600 component values. Standard is the primary comparison using the same model backbone; Record and Brief are workflow ablations. Every task has one result for each configuration.

Composite scores are displayed to four decimal places and component scores to six decimal places. The accompanying `cloudrabench_200_run_scores.csv` retains the original full-precision numeric strings. Accordingly, recomputing the weighted composite from the displayed component values can differ from the separately rounded composite in the last

shown digit. The printed scientific self-consistency score is the graded component, distinct from the separate all-or-none scientific acceptance decision.

Table S16. Complete Easy-tier results for all four configurations. S: composite score; G: graph structure; SC: scientific self-consistency score; A: accuracy.

Task	Configuration	S	G	SC	A
D1	Guided	0.8200	1.000000	1.000000	0.700000
D1	Standard	0.8291	0.791667	0.500000	0.999960
D1	Record	0.7992	0.791667	1.000000	0.700000
D1	Brief	0.7450	1.000000	0.750000	0.700000
D3	Guided	0.7450	1.000000	0.750000	0.700000
D3	Standard	0.9247	1.000000	0.750000	0.999496
D3	Record	0.7940	0.944444	0.350000	0.990939
D3	Brief	0.7283	0.833333	0.750000	0.700000
D4	Guided	0.9250	1.000000	0.750000	0.999982
D4	Standard	0.8500	1.000000	0.500000	0.999982
D4	Record	0.9104	1.000000	0.750000	0.975699
D4	Brief	0.9049	0.944444	0.750000	0.975698
E1	Guided	1.0000	1.000000	1.000000	0.999972
E1	Standard	0.7828	0.777778	0.350000	0.999973
E1	Record	1.0000	1.000000	1.000000	0.999973
E1	Brief	1.0000	1.000000	1.000000	0.999973
F1	Guided	0.9417	0.916667	0.833333	1.000000
F1	Standard	0.0333	0.333333	0.000000	0.000000
F1	Record	0.9917	0.916667	1.000000	1.000000
F1	Brief	0.9117	0.916667	0.750000	0.991654
F4	Guided	0.3601	0.600902	1.000000	0.000000
F4	Standard	0.9083	0.833333	0.750000	0.999999
F4	Record	0.9167	0.916667	0.750000	0.999999
F4	Brief	0.3601	0.600902	1.000000	0.000000
G2	Guided	0.2389	0.888889	0.500000	0.000000
G2	Standard	0.7769	1.000000	0.500000	0.878159
G2	Record	0.7769	1.000000	0.500000	0.878161
G2	Brief	0.8500	1.000000	0.500000	0.999986
G3	Guided	0.9832	0.833333	1.000000	0.999830
G3	Standard	0.1417	0.666667	0.250000	0.000000
G3	Record	0.9832	0.833333	1.000000	0.999830
G3	Brief	0.9749	0.750000	1.000000	0.999833
L1	Guided	0.9999	1.000000	1.000000	0.999899
L1	Standard	0.9749	1.000000	0.916667	0.999900
L1	Record	0.9999	1.000000	1.000000	0.999900
L1	Brief	0.9999	1.000000	1.000000	0.999899
L2	Guided	0.7450	1.000000	0.750000	0.700000
L2	Standard	0.6700	1.000000	0.500000	0.700000
L2	Record	0.8549	1.000000	0.750000	0.883231
L2	Brief	0.8799	1.000000	0.833333	0.883239
M1	Guided	0.9478	1.000000	1.000000	0.913062
M1	Standard	0.7526	1.000000	0.350000	0.912639
M1	Record	0.9476	1.000000	1.000000	0.912639
M1	Brief	0.9476	1.000000	1.000000	0.912638
M14	Guided	1.0000	1.000000	1.000000	1.000000
M14	Standard	0.8050	1.000000	0.350000	1.000000
M14	Record	1.0000	1.000000	1.000000	1.000000
M14	Brief	1.0000	1.000000	1.000000	1.000000
M15	Guided	1.0000	1.000000	1.000000	0.999999
M15	Standard	1.0000	1.000000	1.000000	0.999999
M15	Record	1.0000	1.000000	1.000000	0.999983
M15	Brief	1.0000	1.000000	1.000000	0.999999

Continued on the next page

Table S16 (continued)

Task	Configuration	S	G	SC	A
M16	Guided	0.9748	1.000000	0.916667	0.999691
M16	Standard	0.9998	1.000000	1.000000	0.999692
M16	Record	0.9915	0.916667	1.000000	0.999694
M16	Brief	0.9998	1.000000	1.000000	0.999692
M17	Guided	0.9999	1.000000	1.000000	0.999914
M17	Standard	0.0778	0.777778	0.000000	0.000000
M17	Record	0.9999	1.000000	1.000000	0.999914
M17	Brief	0.9999	1.000000	1.000000	0.999914
M3	Guided	1.0000	1.000000	1.000000	0.999996
M3	Standard	1.0000	1.000000	1.000000	0.999996
M3	Record	1.0000	1.000000	1.000000	0.999996
M3	Brief	1.0000	1.000000	1.000000	0.999996
M8	Guided	0.9996	1.000000	1.000000	0.999335
M8	Standard	0.9996	1.000000	1.000000	0.999339
M8	Record	0.9996	1.000000	1.000000	0.999294
M8	Brief	0.9996	1.000000	1.000000	0.999294
P2	Guided	1.0000	1.000000	1.000000	0.999948
P2	Standard	0.9833	0.833333	1.000000	0.999947
P2	Record	0.9833	0.833333	1.000000	0.999947
P2	Brief	0.9750	1.000000	0.916667	0.999947
P3	Guided	0.9380	1.000000	1.000000	0.896739
P3	Standard	0.1000	1.000000	0.000000	0.000000
P3	Record	0.9380	1.000000	1.000000	0.896735
P3	Brief	0.8309	1.000000	1.000000	0.718109
X1	Guided	0.9999	1.000000	1.000000	0.999916
X1	Standard	0.9999	1.000000	1.000000	0.999914
X1	Record	0.9999	1.000000	1.000000	0.999916
X1	Brief	0.9999	1.000000	1.000000	0.999914

Table S17. Complete Normal-tier results for all four configurations. S: composite score; G: graph structure; SC: scientific self-consistency score; A: accuracy.

Task	Configuration	S	G	SC	A
NB1	Guided	0.9628	0.628571	1.000000	0.999972
NB1	Standard	0.9128	0.628571	0.833333	0.999970
NB1	Record	0.8625	0.600000	0.675000	0.999972
NB1	Brief	0.9900	0.900000	1.000000	0.999972
NI1	Guided	0.9958	1.000000	1.000000	0.993046
NI1	Standard	0.6196	0.946119	0.350000	0.700000
NI1	Record	0.7767	0.816667	0.916667	0.700000
NI1	Brief	0.9889	1.000000	1.000000	0.981480
NI5	Guided	0.7715	0.928571	1.000000	0.631094
NI5	Standard	0.3136	0.571429	0.675000	0.089878
NI5	Record	0.1542	0.442375	0.366667	0.000000
NI5	Brief	0.3571	0.571429	1.000000	0.000000
NQ1	Guided	0.7395	0.945455	0.750000	0.700000
NQ1	Standard	0.9040	0.797980	0.750000	0.998691
NQ1	Record	0.7131	0.680808	0.750000	0.700000
NQ1	Brief	0.1731	0.680808	0.350000	0.000000
NQ5	Guided	0.7075	1.000000	0.625000	0.700000
NQ5	Standard	0.7510	0.920000	0.550000	0.823369
NQ5	Record	0.8725	1.000000	0.625000	0.974999
NQ5	Brief	0.7810	1.000000	0.625000	0.822502
NR1	Guided	0.6167	0.666667	0.833333	0.500000
NR1	Standard	0.1643	0.592593	0.350000	0.000002
NR1	Record	0.5378	0.777778	0.533333	0.500000

Continued on the next page

Table S17 (continued)

Task	Configuration	S	G	SC	A
NR1	Brief	0.6142	0.666667	0.875000	0.475000
NU3	Guided	0.5527	0.776832	0.916667	0.333330
NU3	Standard	0.3621	0.571429	0.350000	0.333330
NU3	Record	0.4071	0.571429	0.500000	0.333330
NU3	Brief	0.6829	0.828571	0.666667	0.666657
NU5	Guided	0.3518	0.642857	0.958333	0.000000
NU5	Standard	0.2654	0.904762	0.116667	0.233136
NU5	Record	0.2966	0.779762	0.233333	0.247741
NU5	Brief	0.1983	0.662546	0.440000	0.000000
NU6	Guided	0.6719	1.000000	0.937500	0.484393
NU6	Standard	0.6892	0.816667	0.625000	0.700000
NU6	Record	0.1450	0.400000	0.350000	0.000000
NU6	Brief	0.3400	0.400000	1.000000	0.000000
NY3	Guided	1.0000	1.000000	1.000000	0.999998
NY3	Standard	0.7485	0.857143	0.616667	0.796239
NY3	Record	0.5217	0.716855	0.833333	0.333333
NY3	Brief	0.7312	0.761905	0.783333	0.700000

Table S18. Complete Hard-tier results for all four configurations. S: composite score; G: graph structure; SC: scientific self-consistency score; A: accuracy.

Task	Configuration	S	G	SC	A
HA1	Guided	0.9625	1.000000	0.875000	1.000000
HA1	Standard	0.1000	1.000000	0.000000	0.000000
HA1	Record	0.9250	1.000000	0.750000	1.000000
HA1	Brief	0.9250	1.000000	0.750000	1.000000
HA6	Guided	0.9750	1.000000	0.916667	1.000000
HA6	Standard	0.9625	1.000000	0.875000	1.000000
HA6	Record	0.9750	1.000000	0.916667	1.000000
HA6	Brief	0.9750	1.000000	0.916667	1.000000
HA8	Guided	0.9625	1.000000	0.875000	0.999961
HA8	Standard	0.6250	1.000000	0.350000	0.700000
HA8	Record	0.9250	1.000000	0.750000	0.999961
HA8	Brief	0.7450	1.000000	0.750000	0.700000
HB1	Guided	0.9325	0.700000	0.875000	1.000000
HB1	Standard	0.8250	0.750000	0.500000	1.000000
HB1	Record	0.9700	0.700000	1.000000	1.000000
HB1	Brief	0.9800	0.800000	1.000000	1.000000
HB10	Guided	0.9995	1.000000	1.000000	0.999225
HB10	Standard	0.9928	0.932344	1.000000	0.999217
HB10	Record	0.8044	0.999011	0.350000	0.999226
HB10	Brief	0.7845	0.800000	0.350000	0.999239
HD2	Guided	0.7867	0.666667	1.000000	0.700000
HD2	Standard	0.6977	0.666667	0.350000	0.876795
HD2	Record	0.5950	1.000000	0.250000	0.700000
HD2	Brief	0.6699	0.999176	0.500000	0.700000
HF1	Guided	0.9742	0.741868	1.000000	1.000000
HF1	Standard	0.3425	0.799505	0.875000	0.000000
HF1	Record	0.9072	0.821728	0.750000	1.000000
HF1	Brief	0.2937	0.687405	0.750000	0.000000
HH1	Guided	0.9838	1.000000	1.000000	0.973067
HH1	Standard	0.8338	1.000000	0.500000	0.973064
HH1	Record	0.7888	1.000000	0.350000	0.973066
HH1	Brief	0.9838	1.000000	1.000000	0.973066
HI1	Guided	0.9999	0.999176	1.000000	0.999998
HI1	Standard	0.7032	0.332509	0.233333	0.999998

Continued on the next page

Table S18 (continued)

Task	Configuration	S	G	SC	A
HI1	Record	0.9350	0.999588	0.783333	0.999998
HI1	Brief	0.9999	0.999176	1.000000	0.999998
HL1	Guided	0.9999	0.999176	1.000000	0.999999
HL1	Standard	0.8050	1.000000	0.350000	0.999999
HL1	Record	0.8972	0.722222	0.750000	0.999995
HL1	Brief	0.9250	1.000000	0.750000	0.999999

Table S19. Complete Impossible-tier results for all four configurations. S: composite score; G: graph structure; SC: scientific self-consistency score; A: accuracy.

Task	Configuration	S	G	SC	A
IA1	Guided	0.1717	0.666667	0.350000	0.000000
IA1	Standard	0.1717	0.666667	0.350000	0.000000
IA1	Record	0.1717	0.666667	0.350000	0.000000
IA1	Brief	1.0000	1.000000	1.000000	1.000000
IA5	Guided	0.4000	1.000000	1.000000	0.000000
IA5	Standard	0.1417	0.666667	0.250000	0.000000
IA5	Record	0.4000	1.000000	1.000000	0.000000
IA5	Brief	0.3667	0.666667	1.000000	0.000000
IA7	Guided	0.2583	0.333333	0.750000	0.000000
IA7	Standard	0.1383	0.333333	0.350000	0.000000
IA7	Record	0.2583	0.333333	0.750000	0.000000
IA7	Brief	0.2583	0.333333	0.750000	0.000000
IA9	Guided	0.3667	0.666667	1.000000	0.000000
IA9	Standard	0.2583	0.333333	0.750000	0.000000
IA9	Record	0.3667	0.666667	1.000000	0.000000
IA9	Brief	0.9333	0.333333	1.000000	1.000000
IB9	Guided	0.8200	1.000000	1.000000	0.700000
IB9	Standard	0.1800	0.750000	0.350000	0.000000
IB9	Record	0.5450	0.200000	0.750000	0.500000
IB9	Brief	0.7700	1.000000	0.833333	0.700000
IC3	Guided	0.4000	1.000000	1.000000	0.000000
IC3	Standard	0.1833	0.333333	0.500000	0.000000
IC3	Record	0.2167	0.666667	0.500000	0.000000
IC3	Brief	0.4000	1.000000	1.000000	0.000000
ID2	Guided	0.3339	0.339244	1.000000	0.000000
ID2	Standard	0.0000	0.000000	0.000000	0.000000
ID2	Record	0.0000	0.000000	0.000000	0.000000
ID2	Brief	0.1839	0.339244	0.500000	0.000000
IE2	Guided	0.3333	0.333333	1.000000	0.000000
IE2	Standard	0.0333	0.333333	0.000000	0.000000
IE2	Record	0.0333	0.333333	0.000000	0.000000
IE2	Brief	0.1833	0.333333	0.500000	0.000000
IF3	Guided	0.3667	0.666667	1.000000	0.000000
IF3	Standard	0.3667	0.666667	1.000000	0.000000
IF3	Record	0.3333	0.333333	1.000000	0.000000
IF3	Brief	0.4000	1.000000	1.000000	0.000000
IG8	Guided	1.0000	1.000000	1.000000	1.000000
IG8	Standard	1.0000	1.000000	1.000000	1.000000
IG8	Record	0.0000	0.000000	0.000000	0.000000
IG8	Brief	0.3339	0.339329	1.000000	0.000000

S4.8 Expanded task-score matrices

The two vector panels below expand the complete per-task score view for reading alongside the exact values in Tables S16–S19. Both use the same color scale and configuration order as the manuscript.



Figure S1. All Easy and Normal task scores. Each task has four configuration rows and four score layers: composite, graph, scientific self-consistency, and accuracy. Cells are rounded to two decimals; the preceding tables and companion CSV retain greater precision. Backbone denotes Standard.

S5 Behavioral diagnostics and archived research trajectories

S5.1 Study materials and local inference settings

The behavioral studies combine archived Cloudra research trajectories with controlled local judgments about the chemical objects, measured quantities, and evidence associations appearing in those trajectories. Historical research events retain their original statements, inputs, returned artifacts, and subsequent actions. Local probes supply frozen questions derived from those events and record the model's answer under specified information and answer-order conditions. The supplementary data retain each question, expected semantic answer, actual answer, request configuration, and source identifier.

The local deployment was archived as `qwen3.8:27b`, using a GGUF Q4_K_M quantization with a reported parameter size of 27.3B. The deployment digest is

`22130167c4c20e20c7b71454612966ca8e8171e9b3cc8ab6ce8aa6cbfec79643.`

The local API metadata reports a maximum model context of 262,144 tokens. The short diagnostic requests use an 8,192-token context. Temperature is 0, seed 42, top-k 0, top-p 1, minimum-p 0, repeat penalty 1, presence penalty 0, and draft prediction count 0. The initial identity and evidence-binding probes use the archived explicit ChatML prompt rendering, an empty thinking block, thinking disabled, and a maximum of eight generated tokens; top-20 token log probabilities are retained. The native-chat controls use separate system and user messages, the deployment's native rendering, and a maximum of 1,024 generated tokens, with the thinking setting specified per request. In every choice task, the saved final answer is evaluated as an exact permitted uppercase choice letter. Semantic answer identity is recorded separately from the displayed letter.




Figure S2. All Hard and Impossible task scores, with the same layer and configuration ordering as Figure S1. All 80 runs have complete composite and component values.

Table S20. Experimental units and crossed conditions of the local diagnostic studies.

Study	Frozen inputs and experimental factors	Calls
Chemical identity	16 pairs × 2 representations × 2 answer orders	64
Distance judgment	4 geometries × 2 information conditions × 2 answer orders × 2 thinking settings	32
Target-angle feedback	2 information conditions; 3 cyclic next-action permutations and 2 sufficiency permutations per condition	10
Evidence associations	4 chains × 2 association conditions × 2 presentations × 2 answer orders	32
Native diagnostic follow-up	20 identity inputs and 16 association inputs, each under 2 thinking settings	72
Outstanding requirements	3 state questions × 2 context conditions × 2 answer orders	12

Call counts retain the actual experimental units. Equivalent representations, answer-order variants, and information conditions are paired versions of the same source object. The single expanded-budget distance follow-up is additional to the 32-call distance study. The activation replacement experiment is described separately below.

S5.2 Chemical identity under equivalent representations and answer reversal

Sixteen molecular-identity pairs were derived from four task systems. They cover identical identities, same-formula connectivity changes, connectivity and formal-charge relocation, double-bond stereochemistry, and protonation changes. For each pair, the original notation and an equivalent rewriting were crossed with two answer orders. In the forward order, A means the same represented chemical identity and B means different represented chemical identities; in the swapped order these meanings exchange. The questions instruct the model to preserve connectivity, bond orders, formal charges, and specified stereochemistry while treating traversal order, ring digits, and disconnected-component order as notation.

Across the resulting 64 judgments, 58 answers were correct. At fixed identity pair and answer order, equivalent rewriting changed the semantic answer in four of 32 comparisons. At fixed pair and representation, reversing answer order changed the semantic answer in six of 32 comparisons. All six order-sensitive cases selected A in both displayed orders. The same-formula example `CC1CC=C1` versus `C=C1CCC1` places the double bond within and outside the ring, respectively. The saved model answer is A in all four representation/order conditions, although the chemical meaning of A changes with the answer order. This design isolates persistent chemical identity from a particular string traversal or preferred answer position.

The identical-string positive conditions provide a baseline for matching a representation to itself. Equivalent rewritings

test whether that judgment survives a notation change. Different-identity controls ask whether molecular formula agreement can coexist with an actual change of connectivity or stereochemistry. The complete frozen strings and semantic labels are retained with the diagnostic data, allowing these contrasts to be reconstructed for every pair.

S5.3 XYZ coordinates and explicitly measured distances

Four NB1 geometries were selected from the archived research trajectories. The local question asks which of two specified atom pairs has the shorter Euclidean distance. The original-XYZ condition supplies the complete geometry in its saved atom order. The measured-distance condition supplies program-calculated values for the same atom pairs. Each condition appears in both answer orders and in direct-answer and thinking modes, yielding 32 native chat calls. Table S21 gives the measured values used to construct the paired conditions; all atom indices are one-based.

Table S21. Distance comparisons derived from the four archived NB1 geometries.

Source workflow	Pair 1	d_1 (Å)	Pair 2	d_2 (Å)
Guided	C2–C4	2.029	O3–H8	1.248
Brief	C4–H11	2.197	O7–H11	0.981
Brief	C4–H11	1.388	O7–H11	1.362
Brief	C4–H11	2.386	O7–H11	0.968

Table S22. Complete outcomes of the native distance-judgment controls.

Thinking	Information	Calls	Correct	Valid wrong	Budget-truncated
Off	XYZ coordinates	8	4	4	0
Off	Measured distances	8	8	0	0
On	XYZ coordinates	8	7	0	1
On	Measured distances	8	8	0	0

A budget-truncated generation reached the 1,024-token ceiling with no final choice. It remains in the eight-input denominator. A separate 2,048-token follow-up of that one input produced the correct choice after 1,353 generated tokens.

All eight direct-answer XYZ responses selected A, giving four correct answers and no semantic agreement across the four answer-reversed source pairs. Providing the measured distances gave eight correct answers in each thinking setting and preserved the semantic answer under reversal. With thinking enabled, median generated-token counts over the eight matched inputs were 807.5 for XYZ and 86 for measured distances. The archived generated-token count includes thinking tokens where enabled. Supplying the measurements makes the relevant physical relation explicit and removes its repeated numerical reconstruction from the subsequent choice. This allocation complements the use of program-executed numerical operations in program-aided language models.[22]

S5.4 Target-specific geometric feedback in an E1 operation chain

The E1 source trajectory specifies the transferring hydrogen H7 in the H7–C2–C3–Cl4 dihedral and seeks an anti-periplanar arrangement near $180°$. Rotation of the C3-side fragment about C2–C3 by $+120°$ and $-120°$ produces candidates A and B, while the C3–Cl4 distance is set to 2.35 Å. The default structural report describes a dihedral involving H5. The subsequent explicit H7 measurement identifies the quantity required by the current research objective.

Table S23. Dihedral measurements and circular residuals for the saved E1 candidates.

Candidate	Atom quadruplet	Signed angle (°)	Distance to 180°
A	H5–C2–C3–Cl4	−55.7831	124.2169
A	H7–C2–C3–Cl4	59.3752	120.6248
B	H5–C2–C3–Cl4	64.2169	115.7831
B	H7–C2–C3–Cl4	179.3752	0.6248

The Newman projections in the manuscript are drawn from the saved candidate coordinates by projection normal to C2→C3, with projected bond directions normalized for display. Table S23 retains the measured signed angles and circular distances to the target. Candidate B is subsequently supplied as the transition-state guess in the actual trajectory. Its submission identifies the H7-based anti-periplanar arrangement and the stretched C3–Cl4 bond as the geometric rationale.

Ten local probes then separate the available measurement from the action question. In the H5-only condition, the H7 values and XYZ geometry are withheld. In the H5-plus-H7 condition, the measured target values are supplied. For next-action questions, all three cyclic option permutations switch from requesting the H7 measurement to selecting candidate B when the target values are added. Two paired information-sufficiency tests likewise switch from requesting additional target information to declaring the comparison sufficient. The facts used by these probes are measurements from the saved geometries. The controlled intervention is which measured quantities are supplied to the decision.

S5.5 Single-token transfer of a reported numerical fact

The activation experiment uses the same archived local quantized deployment and a short reported-distance comparison.[23] One displayed number is changed from 2.029 to 1.029 Å, while the second remains 1.248 Å. This reverses the correct comparison. The clean and edited prompts each contain 124 tokens; their token IDs differ only at global position 53, corresponding to the initial digit of the changed measurement. The original-fact answer is B in the forward option order and A in the reversed order.

For each option order, the residual-stream state at position 53 is captured from the original prompt and transferred into the edited prompt at zero-based layers 7, 15, 31, and 47. The four layers are fixed in the experiment manifest. The readout is the full-vocabulary softmax probability mass assigned to the original-fact answer, summing its bare-letter and leading-space spellings. Table S24 preserves the clean, edited, and transferred-state values.

Table S24. Probability of the original-fact answer under single-position activation replacement.

Input / intervention	Forward: original answer B	Reversed: original answer A
Original input	0.954032	0.996932
Edited input	0.008694	0.073626
Transfer at layer 7	0.029099	0.955908
Transfer at layer 15	0.015033	0.717660
Transfer at layer 31	0.007961	0.072117
Transfer at layer 47	0.008763	0.069810

The readout uses the full vocabulary, with bare and leading-space answer tokens combined. It is distinct from an A/B-renormalized probability or a proportion of correct task runs.

Two controls are executed in both option orders. Self-replacement transfers the edited prompt's own state at the changed fact token; the prefix control transfers at unchanged token position 0. Both produce exactly equal full-vocabulary logits to the edited baseline, with a maximum absolute logit difference of 0. In the reversed order, layer-7 and layer-15 transfers restore substantial probability to the original-fact answer. In the forward order, the corresponding changes are smaller and the edited answer remains preferred. The result connects a localized representation of a supplied measurement to the subsequent choice while retaining the answer-order asymmetry in the numerical record.

S5.6 Evidence associations under fixed local calculation facts

Four source evidence chains from three tasks are used to construct paired coherent and altered-association records. The two NB1 chains share one source run. Each record appears in a flat list and an explicitly linked representation, with both answer orders. This yields $4 \times 2 \times 2 \times 2 = 32$ inputs. Each altered record changes a single association: the geometry used by the saddle-frequency calculation, the IRC starting geometry, the optimization input for a terminal IRC frame, or the identity of the target species. Local normal-termination flags, convergence flags, energies, and imaginary-frequency values remain fixed.

The question asks whether the supplied associations meet the specified support requirements for the indicated reaction step. Linked presentations give the optimized saddle, frequency geometry, IRC start, terminal frames, endpoint optimization/frequency chain, and declared species as explicit references. The model answers all 16 linked inputs correctly and 14 of 16 flat inputs correctly. Both flat-presentation errors occur in altered records with reversed answer options, in E1 and the second NB1 step. The paired records show how the scientific role of a calculation changes when its association changes, even though its local numerical values are unchanged. Explicit relational encoding gives the model direct access to this organization.[24]

A subsequent native-chat follow-up uses the two source chains implicated by these errors and retains every coherent/altered, flat/linked, and option-order variant, giving 16 association questions per thinking mode. The same follow-up also retains all four variants of five identity pairs implicated by the first-stage identity results, giving 20 identity questions per mode. Table S25 records every outcome category in this 72-call follow-up. Under native chat with thinking disabled, the selected identity and association counts remain 14/20 and 14/16 correct. With thinking enabled, association judgments reach 16/16; the identity family has eight correct final choices and 12 budget-truncated generations at the common 1,024-token maximum.

S5.7 From a detected gap to the next research action

The historical HA1 and NB1 examples provide the corresponding research-level sequence. In a Brief HA1 trajectory, compact diagnostics were followed by changes to the source-minimum reference while the IRC endpoint association remained unresolved. In a Guided NB1 trajectory, a frequency result with zero imaginary modes was reported as a normal-mode defect. The Coordinator then requested changes to the transition-state guess and optimization constraints, and the subsequent calculation used the revised parameters. Each source sequence retains the diagnostic, the next scientific instruction, and the resulting computational request. This follows the relation between feedback and a concrete revision rather than using a tool-success flag as the outcome.[20]

Table S25. Complete outcomes for the selected native diagnostic follow-up.

Rendering / mode	Family	Calls	Correct	Valid wrong	Budget-truncated
First-stage raw	Identity	20	14	6	0
First-stage raw	Association	16	14	2	0
Native, thinking off	Identity	20	14	6	0
Native, thinking off	Association	16	14	2	0
Native, thinking on	Identity	20	8	0	12
Native, thinking on	Association	16	16	0	0

The first-stage rows are the selected records from the existing 96-input identity/association study. The four native rows contain 72 additional calls. All selected base cases retain their complete frozen variants; budget-truncated inputs remain in the denominators.

The IA1 completion sequence addresses another decision. Its delivery report has `complete=false`, an unresolved bidirectional short-IRC requirement, and no formal waiver. The Coordinator nevertheless requests a complete stop after earlier affirmative comments about intermediate optimization. The completion check returns the remaining IRC requirement to the active research context. The next stop request explicitly changes its disposition to partial. The sequence records the actual claim change and the outstanding path requirement.

The local state-reading study extracts three questions from this IA1 source: whether the latest report declares completion, whether a formal short-IRC waiver exists, and whether the evidence gap should remain outstanding. Each is asked in a report-only context and a quoted 12-message context, with both answer orders. All 12 judgments agree with the frozen state labels. All six option-reversed pairs retain the same semantic answer. The expected answer is derived from the explicit report fields and waiver list, providing a persistent reference for subsequent completion decisions.

S5.8 Discussion of the diagnostic scope

These experiments resolve distinct operations in the research process: maintaining identity across representations, obtaining a spatial relation, matching feedback to the atoms named in an objective, transmitting a reported fact, connecting calculation artifacts, and reading outstanding scientific requirements. The local measurements identify where explicit representations change the model's decisions. Their interpretation is specific to the archived local deployment, frozen sources, and experimental contrasts. The native follow-up selects sources after inspection of first-stage errors, while retaining complete variants; its counts describe that selected set. Native thinking also changes conditioning and consumed generation, so a common maximum output budget does not imply matched computation.

The activation replacement establishes a local effect on a reported numerical comparison in the quantized Qwen deployment. Its readout does not identify an internal mechanism of the original DeepSeek research runs or a direct measurement of chemical validity. The E1 historical transition to candidate B is observational, with an intervening Supervisor/workspace input before the next model action. The separate local feedback probes isolate supplied target measurements under the curated comparisons.

The completed IA1 state-reading result covers the report-only and 12-message conditions. An additional full 306-message history request contained 242,803 counted input tokens in a 262,144-token context and reached the 3,600-s HTTP waiting limit without a saved answer. Only the first of six planned full-history inputs was attempted; the other five were not attempted. This execution status is retained separately from the 12 completed short-context judgments. Across the diagnostics, generated-answer accuracy, execution status, evidence-association agreement, and scientific acceptance remain distinct observations.

S6 Literature sample and source-linked capability register

S6.1 Selection and coding procedure

The literature sample retained 85 separately described chemical AI agent systems, architectures, or workflows in the 27 October 2026 source snapshot. Inclusion required an LLM to participate in planning, tool use, input/code modification, or feedback-driven decisions in an executed molecular, atomistic, or molecular coarse-grained physical calculation. The unit was a described system or workflow: journal/preprint versions and aliases were reconciled to the same record. The register retains the original sources, evidence locations, and descriptions used for coding.

Records were assigned according to the physical objects demonstrated in their worked examples: materials, crystals, and surfaces (39); finite molecules and complexes (23); liquids, soft matter, and biomolecules (12); or demonstrations spanning multiple object classes (11). Periodicity alone did not determine the class: a protein, liquid, or polymer simulation box remained associated with its actual physical object. Cloudra itself is shown separately in the manuscript figures and is not added to the 85-record denominator.

Task codes denote demonstrated activities: T, structure construction/editing; E, separately demonstrated static energy, electronic structure, or material-property calculations; O, geometry optimization/relaxation; F, vibrations/frequencies and associated thermochemistry; R, explicit reaction paths or transition-state searches, including solid-state NEB; D, molecular or quantum dynamics. A record can carry multiple task codes. The respective code counts are 61, 61, 53, 22, 11, and 37. Empty codes are unencoded activities in this source sample. Coarse-grained normal modes and MD-derived free-energy

profiles retain their specific physical meaning in the evidence register.

S6.2 Source register and counting variants

Table S26 links every system to its primary article or official project record and to additional evidence sources where applicable. The machine-readable files `companion_data/literature/systems.json` and `companion_data/literature/source_index.csv` preserve the exact source locations and supporting descriptions for each assigned code. They also retain the demonstrated targets, boundary conditions, aliases, family assignments, and access/source qualifications. A crosswalk to the 49-system review used during source discovery is supplied with the screening and search records.

The main manuscript uses separately described systems/workflows. Grouping the specified El Agente, MDAgent, MatAgent/ChatMat, and DynaMate/DynaMate2 families gives 75 records (37 materials/surfaces, 17 molecules/complexes, 11 soft/biomolecular, and 10 multiple-class records). Excluding the three official-demonstration-only systems gives 82 records (38, 21, 12, 11); additionally excluding MLP Copilot gives 81 (37, 21, 12, 11). These counting variants are retained with their definitions in the source data.

Table S26. Complete source-linked register of the 85 chemical AI agent systems/workflows used in the literature sample. Task codes are defined in the text. L01–L85 link to the primary records; lettered links give additional sources.

No.	System/workflow	Demonstrated object class	Task codes	Original sources
1	Buehler molecular-DFT multi-agent demonstration	Molecules/complexes	T E	L01, a
2	ProtAgents	Soft/biomolecular	T F	L02, a
3	LLaMP	Materials/surfaces	T D	L03
4	ChemReasoner	Materials/surfaces	E O T	L04
5	AtomAgents	Materials/surfaces	T E O R	L05
6	Adsorb-Agent	Materials/surfaces	T E O	L06
7	LangSim	Materials/surfaces	T E O	L07, a
8	MDCrow	Soft/biomolecular	D	L08
9	MDAgent	Materials/surfaces	D	L09
10	El Agente Q	Molecules/complexes	T E O F	L10
11	Aitomia	Molecules/complexes	T E O F	L11
12	DynaMate (Mendible-Barreto et al.)	Multiple classes	T O D	L12, a, b
13	MatAgent (Lv et al.)	Molecules/complexes	T E	L13
14	ChemGraph	Molecules/complexes	T E O F	L14
15	TopoMAS	Materials/surfaces	T E	L15
16	NAMD-Agent	Soft/biomolecular	T D	L16, a
17	DREAMS	Materials/surfaces	T E O	L17
18	SparksMatter	Materials/surfaces	T E O	L18
19	AutoSolvate-Agent	Molecules/complexes	T E O F	L19, a, b
20	VASPilot	Materials/surfaces	E O	L20
21	ChemAgents molecular simulation workflow	Molecules/complexes	F D	L21, a
22	AMLP	Materials/surfaces	E O D	L22
23	ToPolyAgent	Soft/biomolecular	T D	L23, a
24	GENIUS	Materials/surfaces	E O	L24, a
25	LAMMPS-Agents	Materials/surfaces	T E O F D	L25, a, b
26	DynaMate (protein–ligand)	Soft/biomolecular	T O D	L26, a
27	AGAPI-Agents	Materials/surfaces	T E O	L27
28	MASTER	Materials/surfaces	T E O	L28
29	El Agente Cuántico	Molecules/complexes	E	L29, a
30	VASP Agent	Materials/surfaces	E O R	L30
31	Masgent	Materials/surfaces	T E O	L31
32	CASCADE	Materials/surfaces	E O	L32
33	MDAgent2-RUNTIME	Materials/surfaces	D	L33, a
34	CatMaster	Materials/surfaces	T E O F R	L34

Continued on the next page

Table S26 (continued)

No.	System/workflow	Demonstrated object class	Task codes	Original sources
35	QUASAR	Multiple classes	T E O D	L35
36	ChatMat	Molecules/complexes	T E	L36, a, b, c
37	El Agente Estructural	Molecules/complexes	T O	L37
38	El Agente Quntur	Molecules/complexes	T E O F R	L38
39	El Agente Gráfico	Multiple classes	T E O F	L39, a
40	El Agente Sólido	Materials/surfaces	T E O F	L40
41	Catalyst-Agent	Materials/surfaces	T E O	L41
42	TritonDFT	Materials/surfaces	E O F	L42
43	Chelatron	Molecules/complexes	T E	L43
44	Electrolyte Agent / Reasoning-to-Simulation	Soft/biomolecular	T D	L44, a
45	MatSciAgent	Materials/surfaces	T D	L45
46	OpenClaw computational-chemistry skills workflow	Molecules/complexes	T O D	L46
47	Protomia	Molecules/complexes	T E F	L47, a
48	SimMOF	Materials/surfaces	T E O D	L48
49	PolyJarvis	Soft/biomolecular	T D	L49
50	FermiLink	Multiple classes	E D	L50
51	MatClaw	Materials/surfaces	E D	L51
52	El Agente Forjador	Molecules/complexes	T E O F D	L52
53	MDAgent (biomolecular, 2026)	Soft/biomolecular	T O D	L53, a
54	VirtualLab_CC	Molecules/complexes	T E O F	L54, a, b
55	XDFT	Materials/surfaces	T E	L55
56	CatGo	Multiple classes	T E O F R	L56, a, b, c
57	TSAgent	Materials/surfaces	E O F R	L57
58	Lang2MLIP	Materials/surfaces	T E D	L58
59	AtomisticSkills	Multiple classes	T E O D	L59
60	DynaMate2	Soft/biomolecular	T D	L60, a
61	ChatMOSP	Materials/surfaces	T	L61
62	AutoDFT	Multiple classes	E O F D	L62
63	MLIPilot	Multiple classes	E O D	L63
64	CatDT	Materials/surfaces	T E O F R	L64
65	Paimon	Multiple classes	T O D	L65
66	MDForge	Soft/biomolecular	D	L66
67	URSA LAMMPS agent	Materials/surfaces	T E D	L67, a
68	AdsMind	Materials/surfaces	T E O	L68
69	LADeQ	Molecules/complexes	E	L69
70	MutexaGPT	Soft/biomolecular	T D	L70, a
71	Agent-MD	Materials/surfaces	D	L71
72	CGMas	Soft/biomolecular	T O D	L72
73	matmcp	Materials/surfaces	T E O	L73, a, b
74	Evolving TS workflows	Molecules/complexes	O F R	L74
75	ARCHE	Molecules/complexes	T E O F R	L75
76	Multireference QC agent	Molecules/complexes	E	L76
77	El Agente Potente	Multiple classes	T E O F R D	L77
78	PVmatAgent	Materials/surfaces	T E O	L78, a, b
79	Agent-E2MD	Materials/surfaces	E O D	L79
80	SQD-Agent	Molecules/complexes	E	L80

Continued on the next page

Table S26 (continued)

No.	System/workflow	Demonstrated object class	Task codes	Original sources
81	QAOS / Quantum Agentic Operating System	Molecules/complexes	T E O	L81, a
82	MLP Copilot	Materials/surfaces	E D	L82, a
83	Magnolia	Multiple classes	T E O D	L83
84	GBClaw	Materials/surfaces	T E O R	L84, a
85	PULSES	Molecules/complexes	T E O D	L85, a

S7 Data, software, and reproduction materials

S7.1 Machine-readable task and score records

The supplementary source package contains `companion_data/cloudrabench_50_task_catalogue.csv`, the 50 formal task JSON files in `companion_data/task_definitions/`, and `companion_data/cloudrabench_200_run_scores.csv`. The task JSON files preserve the molecular inputs, task text, permitted methods, Gold graphs, and scoring settings. The score table retains full source precision for all 200 composites and all 600 components. A provenance manifest records file hashes and the displayed precision used by the printed tables.

The P2 publication-source package retains the recorded component sources, paired summaries, Brief/Impossible analyses, timing extractions, and field-level provenance used by the manuscript figures. The reproducible organization script assembles the 200-run publication table from these retained records; figure scripts derive their panels from the corresponding tables. This reproduction does not launch new electronic-structure or agent calculations.

S7.2 Software responsibilities and calculation records

Cloudra revision 70daa5a identifies the implementation described in the main-text methods.[2] RDKit and ETKDG supply molecular representations and coordinate embedding; MMFF94 supplies force-field relaxation.[17, 25–27] NumPy supplies numerical array/linear-algebra operations, NetworkX supplies graph representation and matching, and cclib works with engine-specific readers to extract quantum-chemical quantities.[18, 28, 29] ASE supplies the NEB/MDMin implementation used in standard IDPP, and Adam is used in the lightweight interpolation path.[30–33] Task-name semantic projections use BGE-M3 embeddings and scikit-learn PCA.[34, 35] Publication tables are assembled with pandas, and the quantitative figures are drawn with Matplotlib.[36, 37]

Run-specific records identify the actual engine, model/API name, charge, spin, method, basis, and keywords used in the applicable calculation or request. Dependency lower bounds in the software distribution are installation requirements; they are not substituted for archived experiment versions. Native calculation outputs retain stationary-point energies, frequencies, normal modes, IRC geometries, and endpoint optimization/frequency lineage. The numerical barrier convention is the electronic-energy difference in Equation S1.

S7.3 Reproduction entry points

Within the accompanying Cloudra repository, the following commands reconstruct the retained P2 publication tables and figures:

```
python scripts/benchmark_p2/prepare_publication_data.py
python scripts/benchmark_p2/plot_benchmark_results.py
python scripts/benchmark_p2/plot_speed_delivery.py
python scripts/benchmark_p2/plot_imp.py
```

The companion-data archive preserves the publication reproduction tree under `companion_data/publication_reproduction/`; running the four commands from that directory retains the same relative file layout. The independent timing extractions and frozen behavioral requests/responses are included. The activation archive includes the original prompts, intervention commands, captured states, and full-vocabulary logits, allowing the reported probability readouts and control equalities to be reconstructed without new model inference. Scripts for replay and analysis are retained under `companion_data/behavioral_scripts/`. No model weights are included. The literature directory retains the complete register, screening records, and search history used for the manuscript sample. Editable TeX files, vector figures, and the bibliography are included with this SI so that scientific content, printed values, and source records remain connected during subsequent revision.

S7.4 Data-package inventory

The data archive is distributed separately from the compact Overleaf typesetting project. Unpack it beside the SI source to obtain the referenced `companion_data/` directory. Its subdirectories contain the following materials:

- `task_definitions/`: all 50 original task definitions, including molecular inputs and rubrics.

- `publication_reproduction/`: publication input records, field provenance, bootstrap summaries, timing records, and table/figure scripts.
- `behavioral/`: frozen input sets, actual request/response records, deployment snapshots, geometric/evidence source cases, and activation readouts.
- `behavioral_scripts/`: the corresponding intervention and diagnostic code.
- `literature/`: all 85 source-linked systems and the coding/search records.

A file-level SHA-256 inventory accompanies the archive. Source data are retained at their recorded precision; the SI introduces no new agent or quantum-chemical calculations.

References

[1] DeepSeek-AI. DeepSeek V4 Model Card, 2026. URL `https://fe-static.deepseek.com/chat/transparency/deepseek-V4-model-card-EN.pdf`.

[2] Cloudra. Cloudra source code, 2026. URL `https://github.com/yyywaa/Cloudra-mechanism-agent/tree/70daa5a`. Revision 70daa5a, 26 October 2026.

[3] Christoph Bannwarth, Sebastian Ehlert, and Stefan Grimme. GFN2-xTB—an accurate and broadly parametrized self-consistent tight-binding quantum chemical method with multipole electrostatics and density-dependent dispersion contributions. *Journal of Chemical Theory and Computation*, 15(3):1652–1671, 2019. doi: 10.1021/acs.jctc.8b01176.

[4] James J. P. Stewart. Optimization of parameters for semiempirical methods VI: more modifications to the NDDO approximations and re-optimization of parameters. *Journal of Molecular Modeling*, 19(1):1–32, 2013. doi: 10.1007/s00894-012-1667-x.

[5] Axel D. Becke. Density-functional thermochemistry. III. the role of exact exchange. *The Journal of Chemical Physics*, 98(7): 5648–5652, 1993. doi: 10.1063/1.464913.

[6] Chengteh Lee, Weitao Yang, and Robert G. Parr. Development of the Colle–Salvetti correlation-energy formula into a functional of the electron density. *Physical Review B*, 37(2):785–789, 1988. doi: 10.1103/PhysRevB.37.785.

[7] P. J. Stephens, F. J. Devlin, C. F. Chabalowski, and M. J. Frisch. Ab initio calculation of vibrational absorption and circular dichroism spectra using density functional force fields. *The Journal of Physical Chemistry*, 98(45):11623–11627, 1994. doi: 10.1021/j100096a001.

[8] Stefan Grimme, Jens Antony, Stephan Ehrlich, and Helge Krieg. A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H–Pu. *The Journal of Chemical Physics*, 132(15):154104, 2010. doi: 10.1063/1.3382344.

[9] Stefan Grimme, Stephan Ehrlich, and Lars Goerigk. Effect of the damping function in dispersion corrected density functional theory. *Journal of Computational Chemistry*, 32(7):1456–1465, 2011. doi: 10.1002/jcc.21759.

[10] Florian Weigend and Reinhart Ahlrichs. Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Physical Chemistry Chemical Physics*, 7:3297–3305, 2005. doi: 10.1039/B508541A.

[11] Frank Neese. Software update: The ORCA program system—version 6.0. *WIREs Computational Molecular Science*, 15(2): e70019, 2025. doi: 10.1002/wcms.70019.

[12] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, et al. Gaussian 16. Gaussian, Inc., Wallingford, CT, 2016. URL `https://gaussian.com/citation/`.

[13] Jonathan E. Moussa and James J. P. Stewart. MOPAC: An open-source semiempirical molecular orbital program. *Journal of Open Source Software*, 11(119):8025, 2026. doi: 10.21105/joss.08025.

[14] FACCTs. ORCA 6.1 Manual, 2025. URL `https://www.faccts.de/docs/orca/6.1/manual/`. Online documentation; accessed 31 October 2026.

[15] K. Fukui. Formulation of the reaction coordinate. *The Journal of Physical Chemistry*, 74:4161–4163, 1970. doi: 10.1021/j100717a029.

[16] S. Maeda, Y. Harabuchi, Y. Ono, T. Taketsugu, and K. Morokuma. Intrinsic reaction coordinate: Calculation, bifurcation, and automated search. *International Journal of Quantum Chemistry*, 115:258–269, 2015. doi: 10.1002/qua.24757.

[17] RDKit. Getting Started in Python: Conformer Generation and Maximum Common Substructure, 2026. URL `https://www.rdkit.org/docs/GettingStartedInPython.html`. Online documentation; accessed 31 October 2026.

[18] Aric A. Hagberg, Daniel A. Schult, and Pieter J. Swart. Exploring network structure, dynamics, and function using NetworkX. In Gaël Varoquaux, Travis Vaught, and Jarrod Millman, editors, *Proceedings of the 7th Python in Science Conference*, pages 11–15, Pasadena, CA, 2008. URL `https://aric.hagberg.org/papers/hagberg-2008-exploring.pdf`.

[19] Karthik Valmeekam, Matthew Marquez, Sarath Sreedharan, and Subbarao Kambhampati. On the planning abilities of large language models: A critical investigation. In *Advances in Neural Information Processing Systems*, 2023. URL `https://arxiv.org/abs/2305.15771`.

[20] Zhibin Gou, Zhihong Shao, Yeyun Gong, Yelong Shen, Yujiu Yang, Nan Duan, and Weizhu Chen. CRITIC: Large language models can self-correct with tool-interactive critiquing. In *International Conference on Learning Representations*, 2024. URL `https://arxiv.org/abs/2305.11738`.

[21] Jie Huang, Xinyun Chen, Swaroop Mishra, Huaixiu Steven Zheng, Adams Wei Yu, Xinying Song, and Denny Zhou. Large language models cannot self-correct reasoning yet. In *International Conference on Learning Representations*, 2024. URL `https://arxiv.org/abs/2310.01798`.

[22] Luyu Gao, Aman Madaan, Shuyan Zhou, Uri Alon, Pengfei Liu, Yiming Yang, Jamie Callan, and Graham Neubig. PAL: Program-aided language models. In *Proceedings of the 40th International Conference on Machine Learning*, volume 202, pages 10764–10799, 2023. URL https://proceedings.mlr.press/v202/gao23f.html.

[23] Fred Zhang and Neel Nanda. Towards Best Practices of Activation Patching in Language Models: Metrics and Methods. In *International Conference on Learning Representations*, 2024. URL https://proceedings.iclr.cc/paper_files/paper/2024/hash/06a52a54c8ee03cd86771136bc91eb1f-Abstract-Conference.html.

[24] Bahar Fatemi, Bryan Perozzi, and Jonathan Halcrow. Talk like a Graph: Encoding Graphs for Large Language Models. In *International Conference on Learning Representations*, 2024. URL https://research.google/pubs/talk-like-a-graph-encoding-graphs-for-large-language-models/.

[25] S. Riniker and G. A. Landrum. Better Informed Distance Geometry: Using What We Know To Improve Conformation Generation. *Journal of Chemical Information and Modeling*, 55:2562–2574, 2015. doi: 10.1021/acs.jcim.5b00654.

[26] Shuzhe Wang, Jagna Witek, Gregory A. Landrum, and Sereina Riniker. Improving conformer generation for small rings and macrocycles based on distance geometry and experimental torsional-angle preferences. *Journal of Chemical Information and Modeling*, 60(4):2044–2058, 2020. doi: 10.1021/acs.jcim.0c00025.

[27] Thomas A. Halgren. Merck molecular force field. I. basis, form, scope, parameterization, and performance of MMFF94. *Journal of Computational Chemistry*, 17(5–6):490–519, 1996. doi: 10.1002/(SICI)1096-987X(199604)17:5/6<490::AID-JCC1>3.0.CO;2-P.

[28] Charles R. Harris, K. Jarrod Millman, Stéfan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J. Smith, Robert Kern, Matti Picus, Stephan Hoyer, Marten H. van Kerkwijk, Matthew Brett, Allan Haldane, Jaime Fernández del Río, Mark Wiebe, Pearu Peterson, Pierre Gérard-Marchant, Kevin Sheppard, Tyler Reddy, Warren Weckesser, Hameer Abbasi, Christoph Gohlke, and Travis E. Oliphant. Array programming with NumPy. *Nature*, 585(7825):357–362, 2020. doi: 10.1038/s41586-020-2649-2.

[29] Noel M. O'Boyle, Adam L. Tenderholt, and Karol M. Langner. cclib: A library for package-independent computational chemistry algorithms. *Journal of Computational Chemistry*, 29(5):839–845, 2008. doi: 10.1002/jcc.20823.

[30] Søren Smidstrup, Andreas Pedersen, Kurt Stokbro, and Hannes Jónsson. Improved initial guess for minimum energy path calculations. *The Journal of Chemical Physics*, 140(21):214106, 2014. doi: 10.1063/1.4878664.

[31] Graeme Henkelman and Hannes Jónsson. Improved tangent estimate in the nudged elastic band method for finding minimum energy paths and saddle points. *The Journal of Chemical Physics*, 113(22):9978–9985, 2000. doi: 10.1063/1.1323224.

[32] Ask Hjorth Larsen, Jens Jørgen Mortensen, Jakob Blomqvist, Ivano E. Castelli, Rune Christensen, Marcin Dułak, Jesper Friis, Michael N. Groves, Bjørk Hammer, Cory Hargus, et al. The atomic simulation environment—a Python library for working with atoms. *Journal of Physics: Condensed Matter*, 29(27):273002, 2017. doi: 10.1088/1361-648X/aa680e.

[33] Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization. In *International Conference on Learning Representations*, 2015. URL https://arxiv.org/abs/1412.6980.

[34] Jianlyu Chen, Shitao Xiao, Peitian Zhang, Kun Luo, Defu Lian, and Zheng Liu. M3-Embedding: Multi-linguality, multi-functionality, multi-granularity text embeddings through self-knowledge distillation. In *Findings of the Association for Computational Linguistics: ACL 2024*, pages 2318–2335, 2024. doi: 10.18653/v1/2024.findings-acl.137. URL https://aclanthology.org/2024.findings-acl.137/.

[35] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay. Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research*, 12:2825–2830, 2011. URL https://jmlr.org/papers/v12/pedregosa11a.html.

[36] Wes McKinney. Data structures for statistical computing in Python. In *Proceedings of the 9th Python in Science Conference*, 2010. doi: 10.25080/Majora-92bf1922-00a.

[37] John D. Hunter. Matplotlib: A 2d graphics environment. *Computing in Science & Engineering*, 9(3):90–95, 2007. doi: 10.1109/MCSE.2007.55.