

# COUNTING UNIPOTENT REPRESENTATIONS OF REAL REDUCTIVE GROUPS

DAN M. BARBASCH, JIA-JUN MA, BINYONG SUN, AND CHEN-BO ZHU

## CONTENTS

1.	Counting unipotent representations	1
2.	Parameterize of Unipotent representations	2
	References	3

## 1. COUNTING UNIPOTENT REPRESENTATIONS

In this section, let  $G_{\mathbb{C}}$  be a connected complex reductive group and  $\mathfrak{g}$  is its Lie algebra. Fix a antiholomorphic involution  $\sigma$  on  $G_{\mathbb{C}}$  and a corresponding Cartan involution  $\theta$  of  $G_{\mathbb{C}}$ . Let  $G$  be a finite central extension of a open subgroup of  $G_{\mathbb{C}}^{\sigma}$  and

$$\text{pr}: G \rightarrow G_{\mathbb{C}}^{\sigma}$$

be the canonical projection. Let  $K = \text{pr}^{-1}(G_{\mathbb{C}}^{\sigma})$ .

Let  ${}^a\mathfrak{h}$  be the abstract Cartan subalgebra of  $\mathfrak{g}$  and  ${}^aX$  be the lattice of abstract weight spaces. Let  ${}^aR \subseteq {}^aX$ ,  ${}^aR^+$  and  ${}^aQ$  be the abstract root system, the set of positive roots and the root lattice. Let

$$\mathbb{C} = \left\{ \mu \in {}^a\mathfrak{h}^* \mid \begin{array}{l} \text{either } \langle \text{Re}(\mu), \check{\alpha} \rangle > 0 \text{ or} \\ \langle \text{Re}(\mu), \check{\alpha} \rangle = 0 \text{ and } \sqrt{-1} \langle \text{Im}(\mu), \check{\alpha} \rangle > 0 \end{array} \right\}$$

and  $\overline{\mathbb{C}}$  be the closure of  $\mathbb{C}$  in  ${}^a\mathfrak{h}$ .

**1.1. Coherent family.** For each finite dimensional  $\mathfrak{g}$ -module or  $G_{\mathbb{C}}$ -module  $F$ , let  $F^*$  be its contragredient representation and let  $\Delta(F) \subseteq {}^aX$  denote the multi-set of weights in  $F$ .

Let  $\Pi_{\Lambda_0}(G_{\mathbb{C}})$  be the set of irreducible finite dimensional representations of  $G_{\mathbb{C}}$  with external weight in  $\Lambda_0$  and  $\mathcal{G}_{\Lambda}(G_{\mathbb{C}})$  be the subgroup generated by  $\Pi_{\Lambda_0}(G_{\mathbb{C}})$ . Let

$${}^aP := \{ \mu \in {}^aX \mid \mu \text{ is a } {}^a\mathfrak{h}\text{-weight of an } F \in \Pi_{\text{fin}}(G_{\mathbb{C}}) \}.$$

Via the highest weight theory, every  $W(G_{\mathbb{C}})$ -orbit  $W \cdot \mu$  in  ${}^aP$  corresponds with the irreducible finite dimensional representation  $F \in \Pi_{\text{fin}}(G_{\mathbb{C}})$  with external weight  $\mu$ .

Now the Grothendieck group  $\mathcal{G}(G_{\mathbb{C}})$  of finite dimensional representation of  $G_{\mathbb{C}}$  is identified with  $\mathbb{Z}[{}^aP/W]$ . In fact  $\mathcal{G}(G_{\mathbb{C}})$  is a  $\mathbb{Z}$ -algebra under the tensor product and equipped with the involution  $F \mapsto F^*$ .

Fix a  $W$ -invariant sub-lattice  $\Lambda_0 \subset {}^aX$  containing  ${}^aQ$ .

Take a lattice  $\Lambda = \lambda + \Lambda_0 \in \mathfrak{h}^*/\Lambda$  with  $\lambda \in \overline{\mathbb{C}}$ . Let

$$R(\lambda) := \{ \alpha \in {}^aR \mid \langle \lambda, \check{\alpha} \rangle = 0 \} \quad \text{and} \quad W(\lambda) := \langle s_{\alpha} \mid \alpha \in R(\lambda) \rangle.$$

---

2000 *Mathematics Subject Classification.* 22E45, 22E46.

*Key words and phrases.* orbit method, unitary dual, unipotent representation, classical group, theta lifting, moment map.

**Definition 1.1.** Suppose  $\mathcal{M}$  is an abelian group with  $\mathcal{G}_\Lambda(G_\mathbb{C})$ -action

$$\mathcal{G}_\Lambda(G_\mathbb{C}) \times \mathcal{M} \ni (F, m) \mapsto F \otimes m.$$

In addition, we fix a subgroup  $\mathcal{M}_\mu$  of  $\mathcal{M}$  for each  $\mu \in \Lambda$ .

A function  $f: \Lambda \rightarrow \mathcal{M}$  is called a coherent family based on  $\Lambda$  if it satisfies  $f(\mu) \in \mathcal{M}_\mu$  and

$$F \otimes f(\mu) = \sum_{\nu \in \Delta(F)} f(\mu + \nu) \quad \forall \mu \in \Lambda, F \in \Pi_{\Lambda_0}(G_\mathbb{C}).$$

Let  $\text{Coh}_\Lambda(\mathcal{M})$  be the abelian group of all coherent families based on  $\Lambda$  and value in  $\mathcal{M}$ .

In this paper, we will consider the following cases.

Suppose  $\mathcal{M} = \mathbb{Q}$  and  $F \otimes m = \dim(F) \cdot m$  for  $F \in \Pi_{\Lambda_0}(G_\mathbb{C})$  and  $m \in \mathcal{M}$ . We let  $\mathcal{M}_\mu = \mathcal{M}$  for every  $\mu \in \Lambda$ . When  $\Lambda = \Lambda_0$ , the set of  $W(G_\mathbb{C})$ -harmonic polynomials on  ${}^a\mathfrak{h}^*$  is naturally identified with  $\text{Coh}_\Lambda(\mathcal{M})$  via restriction (Vogan's result)

Let  $\mathcal{G}(\mathfrak{g}, K)$  be the Grothendieck group of finite length  $(\mathfrak{g}, K)$ -modules and  $\mathcal{G}_\mu(\mathfrak{g}, K)$  be the subgroup of  $\mathcal{G}(\mathfrak{g}, K)$  generated by the set of irreducible  $(\mathfrak{g}, K)$ -modules with infinitesimal character  $\mu$ .

Then  $\text{Coh}_\Lambda(\mathcal{G}(\mathfrak{g}, K))$  is the group of coherent families of Harish-Chandra modules.

## 2. PARAMETERIZE OF UNIPOTENT REPRESENTATIONS

We fix an abstract complex Cartan subgroup  $\mathbf{H}_a$  and  $\mathfrak{h}_a$  in  $\mathbf{G}$  and a set of simple roots  $\Pi_a$ . Let  $\mathcal{P}(\mathbf{G})$  be the set of all Langlands parameters of  $G$ -modules with character  $\rho$  (i.e. the infinitesimal character of the trivial representation). For  $\gamma \in \mathcal{P}(\mathbf{G})$ , let  $\mathcal{L}(\gamma)$ ,  $\mathcal{S}(\gamma)$  and  $\Phi_\gamma$  be the corresponding Langlands quotient, standard module and coherent family such that  $\Phi_\gamma(\rho) = \mathcal{L}(\gamma)$ . Let  $\mathcal{M}(\mathbf{G})$  be the span of  $\mathcal{L}(\gamma)$ . Let  $\{\mathcal{B}\}$  be the set of all blocks. Then  $\mathcal{P}(\mathbf{G}) = \bigsqcup_{\mathcal{B}} \mathcal{B}$ . The Weyl group  $W = W(G)$  acts on  $\mathcal{M}(\mathbf{G})$  by coherent continuation. Let  $\mathcal{M}_{\mathcal{B}}$  be the submodule of  $\mathcal{M}(\mathbf{G})$  spanned by  $\gamma \in \mathcal{B}$ , then

$$\mathcal{M}(\mathbf{G}) = \bigoplus_{\mathcal{B}} \mathcal{M}_{\mathcal{B}}$$

Let  $\tau(\gamma) \subset \Pi_a$  be the  $\tau$ -invariant of  $\gamma$ .

Let  $\mathcal{O}$  be even orbit.  $\lambda = \frac{1}{2}\check{h}$ . Define

$$S(\lambda) = \{ \alpha \in \Pi_a \mid \langle \alpha, \lambda \rangle = 0 \}.$$

Let  $\mathcal{P}_\lambda(\mathbf{G})$  be the set of all Langlands parameters with infinitesimal character  $\lambda$ . Let  $T_{\lambda, \rho}$  be the translation functor. Let

$$\mathcal{B}(S) = \{ \gamma \in \mathcal{B} \mid S \cap \tau(\gamma) = \emptyset \}$$

and

$$\mathcal{P}(\mathbf{G}, S) = \bigsqcup_{\mathcal{B}} \mathcal{B}(S)$$

Then

$$\begin{aligned} \mathcal{P}(\mathbf{G}, S) &\longrightarrow \mathcal{P}_\lambda(\mathbf{G}) \\ \gamma &\longmapsto T_{\lambda, \rho}(\gamma) \end{aligned}$$

Let  $\mathcal{O}$  be a complex nilpotent orbit in  $\mathfrak{g}$ . Let

$$\mathcal{B}(S, \mathcal{O}) = \{ \gamma \in \mathcal{B}(S) \mid \text{AV}_\mathbb{C}(\mathcal{L}(\gamma)) \subset \overline{\mathcal{O}} \}$$

Let

$$\begin{aligned} m_S(\sigma) &= [\sigma : \text{Ind}_{W(S)}^W \mathbf{1}] \\ m_{\mathcal{B}}(\sigma) &= [\sigma : \mathcal{M}_{\mathcal{B}}] \end{aligned}$$

Barbasch [10, Theorem 9.1] established the following theorem.

**Theorem 2.1.**

$$|\mathcal{B}(S, \mathcal{O})| = \sum_{\sigma} m_{\mathcal{B}}(\sigma) m_S(\sigma)$$

Here  $\sigma \times \sigma$  running over the  $W \times W$  appears in the double cell  $\mathcal{C}(\mathcal{O})$ .

*Proof.* We need to take the graded module of  $\mathcal{M}(\mathbf{G})$  with respect to the  $\overset{LR}{\leq}$ . By abuse of notation, we identify the basis  $\mathcal{P}(\mathbf{G})$  with its image in the graded module. Note that  $S \cap \tau(\lambda) = \emptyset$  if and only if  $W(S)$  acts on  $\gamma$  trivially by [70, Lemma 14.7]. On the other hand, by [70, Theorem 14.10, and page 58],  $\text{AV}_{\mathbf{C}}(\mathcal{L}(\gamma)) \subset \overline{\mathcal{O}}$  only if  $\gamma$  generate a  $W$ -module in the double cell of  $\mathcal{O}$ .  $\square$

Now assume  $S = S(\lambda)$ . By [12, Cor 5.30 b) and c)],  $[\sigma : \text{Ind}_{W(S)}^W \mathbf{1}] = [\mathbf{1}|_{W(S)} : \sigma] \leq 1$ .

## REFERENCES

- [1] J. Adams, *Discrete spectrum of the reductive dual pair  $(O(p, q), Sp(2m))$* , Invent. Math. **74** (1983), no. 3, 449–475.
- [2] J. Adams, D. Barbasch, and D. A. Vogan, *The Langlands classification and irreducible characters for real reductive groups*, Progress in Math., vol. 104, Birkhauser, 1991.
- [3] Jeffrey Adams and Fokko du Cloux, *Algorithms for representation theory of real reductive groups*, Journal of the Institute of Mathematics of Jussieu **8** (2009), no. 2, 209–259, DOI 10.1017/S1474748008000352.
- [4] J. Arthur, *On some problems suggested by the trace formula*, Lie group representations, II (College Park, Md.), Lecture Notes in Math. 1041 (1984), 1–49.
- [5] ———, *Unipotent automorphic representations: conjectures*, Orbites unipotentes et représentations, II, Astérisque **171-172** (1989), 13–71.
- [6] L. Auslander and B. Kostant, *Polarizations and unitary representations of solvable Lie groups*, Invent. Math. **14** (1971), 255–354.
- [7] D. Barbasch, *Unipotent representations for real reductive groups*, Proceedings of ICM (1990), Kyoto (2000), 769–777.
- [8] Dan Barbasch and David Vogan, *Weyl Group Representations and Nilpotent Orbits*, Representation Theory of Reductive Groups: Proceedings of the University of Utah Conference 1982, 1983, pp. 21–33.
- [9] D. Barbasch, *Orbital integrals of nilpotent orbits*, The mathematical legacy of Harish-Chandra, Proc. Sympos. Pure Math. **68** (2000), 97–110.
- [10] ———, *The unitary spherical spectrum for split classical groups*, J. Inst. Math. Jussieu **9** (2010), 265–356.
- [11] ———, *Unipotent representations and the dual pair correspondence*, J. Cogdell et al. (eds.), Representation Theory, Number Theory, and Invariant Theory, In Honor of Roger Howe. Progress in Math. **323** (2017), 47–85.
- [12] D. Barbasch and D. A. Vogan, *Unipotent representations of complex semisimple groups*, Annals of Math. **121** (1985), no. 1, 41–110.
- [13] R. Brylinski, *Dixmier algebras for classical complex nilpotent orbits via Kraft-Procesi models. I*, The orbit method in geometry and physics (Marseille, 2000). Progress in Math. **213** (2003), 49–67.
- [14] Roger W. Carter, *Finite groups of Lie type*, Wiley Classics Library, John Wiley & Sons, Ltd., Chichester, 1993.
- [15] W. Casselman, *Canonical extensions of Harish-Chandra modules to representations of  $G$* , Canad. J. Math. **41** (1989), 385–438.
- [16] F. Du Cloux, *Sur les représentations différentiables des groupes de Lie algébriques*, Ann. Sci. École Norm. Sup. **24** (1991), no. 3, 257–318.
- [17] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebra: an introduction*, Van Nostrand Reinhold Co., 1993.
- [18] A. Daszkiewicz, W. Kraśkiewicz, and T. Przebinda, *Nilpotent orbits and complex dual pairs*, J. Algebra **190** (1997), no. 2, 518 – 539.
- [19] ———, *Dual pairs and Kostant-Sekiguchi correspondence. II. Classification of nilpotent elements*, Central European J. Math. **3** (2005), 430–474.
- [20] J. Dixmier and P. Malliavin, *Factorisations de fonctions et de vecteurs indéfiniment différentiables*, Bull. Sci. Math. (2) **102** (1978), 307–330.

- [21] M. Duflo, *Théorie de Mackey pour les groupes de Lie algébriques*, Acta Math. **149** (1982), no. 3-4, 153–213.
- [22] R. Gomez and C.-B. Zhu, *Local theta lifting of generalized Whittaker models associated to nilpotent orbits*, Geom. Funct. Anal. **24** (2014), no. 3, 796–853.
- [23] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique IV: Étude locale des schémas et des morphismes de schémas. II*, Inst. Hautes Études Sci. Publ. Math. **24** (1965).
- [24] ———, *Éléments de géométrie algébrique IV: Étude locale des schémas et des morphismes de schémas. III*, Inst. Hautes Études Sci. Publ. Math. **28** (1966).
- [25] M. Harris, J.-S. Li, and B. Sun, *Theta correspondences for close unitary groups*, Arithmetic Geometry and Automorphic Forms, Adv. Lect. Math. (ALM) **19** (2011), 265–307.
- [26] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics, 52. New York-Heidelberg-Berlin: Springer-Verlag, 1983.
- [27] H. He, *Unipotent representations and quantum induction*, arXiv:math/0210372 (2002).
- [28] J.-S. Huang and J.-S. Li, *Unipotent representations attached to spherical nilpotent orbits*, Amer. J. Math. **121** (1999), no. 3, 497–517.
- [29] J.-S. Huang and C.-B. Zhu, *On certain small representations of indefinite orthogonal groups*, Represent. Theory **1** (1997), 190–206.
- [30] R. Howe,  *$\theta$ -series and invariant theory*, Automorphic Forms, Representations and  $L$ -functions, Proc. Sympos. Pure Math, vol. 33, 1979, pp. 275–285.
- [31] ———, *On a notion of rank for unitary representations of the classical groups*, Harmonic analysis and group representations, Liguori, Naples (1982), 223–331.
- [32] ———, *Transcending classical invariant theory*, J. Amer. Math. Soc. **2** (1989), 535–552.
- [33] ———, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, Piatetski-Shapiro, I. et al. (eds.), The Schur lectures (1992). Ramat-Gan: Bar-Ilan University, Isr. Math. Conf. Proc. 8, (1995), 1–182.
- [34] D. Jiang, B. Liu, and G. Savin, *Raising nilpotent orbits in wave-front sets*, Represent. Theory **20** (2016), 419–450.
- [35] A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Uspehi Mat. Nauk **17** (1962), 57–110.
- [36] B. Kostant, *Quantization and unitary representations*, Lectures in Modern Analysis and Applications III, Lecture Notes in Math. **170** (1970), 87–208.
- [37] H. Kraft and C. Procesi, *On the geometry of conjugacy classes in classical groups*, Comment. Math. Helv. **57** (1982), 539–602.
- [38] S. S. Kudla and S. Rallis, *Degenerate principal series and invariant distributions*, Israel J. Math. **69** (1990), 25–45.
- [39] S. S. Kudla, *Some extensions of the Siegel-Weil formula*, In: Gan W., Kudla S., Tschinkel Y. (eds) Eisenstein Series and Applications. Progress in Mathematics, vol 258. Birkhäuser Boston (2008), 205–237.
- [40] S. T. Lee and C.-B. Zhu, *Degenerate principal series and local theta correspondence II*, Israel J. Math. **100** (1997), 29–59.
- [41] ———, *Degenerate principal series of metaplectic groups and Howe correspondence*, D. Prasad et al. (eds.), Automorphic Representations and L-Functions, Tata Institute of Fundamental Research, India, (2013), 379–408.
- [42] J.-S. Li, *Singular unitary representations of classical groups*, Invent. Math. **97** (1989), no. 2, 237–255.
- [43] Q. Liu, *Algebraic Geometry and Arithmetic Curves*, Oxford University Press, 2006.
- [44] H. Y. Loke and J. Ma, *Invariants and  $K$ -spectrums of local theta lifts*, Compositio Math. **151** (2015), 179–206.
- [45] G. Lusztig and N. Spaltenstein, *Induced unipotent classes*, j. London Math. Soc. **19** (1979), 41–52.
- [46] G. Lusztig, *Intersection cohomology complexes on a reductive group*, Invent. Math. **75** (1984), no. 2, 205–272, DOI 10.1007/BF01388564. MR732546
- [47] G. W. Mackey, *Unitary representations of group extensions*, Acta Math. **99** (1958), 265–311.
- [48] W. M. McGovern, *Cells of Harish-Chandra modules for real classical groups*, Amer. J. of Math. **120** (1998), 211–228.
- [49] C. Mœglin, *Front d’onde des représentations des groupes classiques  $p$ -adiques*, Amer. J. Math. **118** (1996), 1313–1346.
- [50] ———, *Paquets d’Arthur Spéciaux Unipotents aux Places Archimédiennes et Correspondance de Howe*, J. Cogdell et al. (eds.), Representation Theory, Number Theory, and Invariant Theory, In Honor of Roger Howe. Progress in Math. **323** (2017), 469–502.

- [51] C. Mœglin, M.-F. Vignéras, and J.-L. Waldspurger, *Correspondances de Howe sur un corps  $p$ -adique*, Lecture Notes in Mathematics, vol. 1291, Springer, 1987.
- [52] K. Nishiyama, H. Ochiai, K. Taniguchi, H. Yamashita, and S. Kato, *Nilpotent orbits, associated cycles and Whittaker models for highest weight representations*, *Astérisque* **273** (2001), 1–163.
- [53] K. Nishiyama, H. Ochiai, and C.-B. Zhu, *Theta lifting of nilpotent orbits for symmetric pairs*, *Trans. Amer. Math. Soc.* **358** (2006), 2713–2734.
- [54] K. Nishiyama and C.-B. Zhu, *Theta lifting of unitary lowest weight modules and their associated cycles*, *Duke Math. J.* **125** (2004), no. 03, 415–465.
- [55] T. Ohta, *The closures of nilpotent orbits in the classical symmetric pairs and their singularities*, *Tohoku Math. J.* **43** (1991), no. 2, 161–211.
- [56] ———, *Induction of nilpotent orbits for real reductive groups and associated varieties of standard representations*, *Hiroshima Math. J.* **29** (1999), no. 2, 347–360.
- [57] ———, *Nilpotent orbits of  $\mathbb{Z}_4$ -graded Lie algebra and geometry of moment maps associated to the dual pair  $(U(p, q), U(r, s))$* , *Publ. RIMS* **41** (2005), no. 3, 723–756.
- [58] A. Paul and P. Trapa, *Some small unipotent representations of indefinite orthogonal groups and the theta correspondence*, *University of Aarhus Publ. Series* **48** (2007), 103–125.
- [59] V. L. Popov and E. B. Vinberg, *Invariant Theory*, *Algebraic Geometry IV: Linear Algebraic Groups, Invariant Theory*, *Encyclopedia of Mathematical Sciences*, vol. 55, Springer, 1994.
- [60] T. Przebinda, *The duality correspondence of infinitesimal characters*, *Colloq. Math.* **70** (1996), 93–102.
- [61] ———, *Characters, dual pairs, and unitary representations*, *Duke Math. J.* **69** (1993), no. 3, 547–592.
- [62] S. Rallis, *On the Howe duality conjecture*, *Compositio Math.* **51** (1984), 333–399.
- [63] S. Sahi, *Explicit Hilbert spaces for certain unipotent representations*, *Invent. Math.* **110** (1992), no. 2, 409–418.
- [64] J. Sekiguchi, *Remarks on real nilpotent orbits of a symmetric pair*, *J. Math. Soc. Japan* **39** (1987), no. 1, 127–138.
- [65] W. Schmid and K. Vilonen, *Characteristic cycles and wave front cycles of representations of reductive Lie groups*, *Annals of Math.* **151** (2000), no. 3, 1071–1118.
- [66] E. Sommers, *Lusztig’s canonical quotient and generalized duality*, *J. Algebra* **243** (2001), no. 2, 790–812.
- [67] T. A. Springer and R. Steinberg, *Seminar on algebraic groups and related finite groups; Conjugate classes*, *Lecture Notes in Math.*, vol. 131, Springer, 1970.
- [68] B. Sun and C.-B. Zhu, *A general form of Gelfand-Kazhdan criterion*, *Manuscripta Math.* **136** (2011), 185–197.
- [69] P. Trapa, *Special unipotent representations and the Howe correspondence*, *University of Aarhus Publication Series* **47** (2004), 210–230.
- [70] D. A. Vogan, *Irreducible characters of semisimple Lie groups. IV. Character-multiplicity duality*, *Duke Math. J.* **49** (1982), no. 4, 943–1073. MR683010
- [71] ———, *Unitary representations of reductive Lie groups*, *Ann. of Math. Stud.*, vol. 118, Princeton University Press, 1987.
- [72] ———, *Associated varieties and unipotent representations*, *Harmonic analysis on reductive groups*, *Proc. Conf., Brunswick/ME (USA) 1989*, *Prog. Math.* **101** (1991), 315–388.
- [73] ———, *The method of coadjoint orbits for real reductive groups*, *Representation theory of Lie groups (Park City, UT, 1998)*, *IAS/Park City Math. Ser.* **8** (2000), 179–238.
- [74] ———, *Unitary representations of reductive Lie groups*, *Mathematics towards the Third Millennium (Rome, 1999)*, *Accademia Nazionale dei Lincei*, (2000), 147–167.
- [75] N. R. Wallach, *Real reductive groups I*, *Academic Press Inc.*, 1988.
- [76] ———, *Real reductive groups II*, *Academic Press Inc.*, 1992.
- [77] H. Weyl, *The classical groups: their invariants and representations*, *Princeton University Press*, 1947.
- [78] S. Yamana, *Degenerate principal series representations for quaternionic unitary groups*, *Israel J. Math.* **185** (2011), 77–124.

THE DEPARTMENT OF MATHEMATICS, 310 MALOTT HALL, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853

*Email address:* `dmb14@cornell.edu`

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, 800 DONGCHUAN ROAD, SHANGHAI, 200240, CHINA

*Email address:* `hoxide@sjtu.edu.cn`

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, CHINA

*Email address:* `sun@math.ac.cn`

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076

*Email address:* `matzhucb@nus.edu.sg`