

COUNTING UNIPOTENT REPRESENTATIONS OF REAL REDUCTIVE GROUPS

DAN M. BARBASCH, JIA-JUN MA, BINYONG SUN, AND CHEN-BO ZHU

CONTENTS

1.	Counting unipotent representations	1
2.	Parameterize of Unipotent representations	3
3.	Counting in type A	4
4.	Counting in type BCD	8
	References	8

1. COUNTING UNIPOTENT REPRESENTATIONS

In this section, let $G_{\mathbb{C}}$ be a connected complex reductive group and \mathfrak{g} is its Lie algebra. Fix a antiholomorphic involution σ on $G_{\mathbb{C}}$ and a corresponding Cartan involution θ of $G_{\mathbb{C}}$. Let G be a finite central extension of a open subgroup of $G_{\mathbb{C}}^{\sigma}$ and

$$\text{pr}: G \rightarrow G_{\mathbb{C}}^{\sigma}$$

be the canonical projection. Let $K = \text{pr}^{-1}(G_{\mathbb{C}}^{\sigma})$ and $W = W(G_{\mathbb{C}})$.

Let ${}^a\mathfrak{h}$ be the abstract Cartan subalgebra of \mathfrak{g} and aX be the lattice of abstract weight spaces. Let ${}^aR \subseteq {}^aX$, ${}^aR^+$ and aQ be the abstract root system, the set of positive roots and the root lattice. Let

$$\mathbb{C} = \left\{ \mu \in {}^a\mathfrak{h}^* \mid \begin{array}{l} \text{either } \langle \text{Re}(\mu), \check{\alpha} \rangle > 0 \text{ or} \\ \langle \text{Re}(\mu), \check{\alpha} \rangle = 0 \text{ and } \sqrt{-1} \langle \text{Im}(\mu), \check{\alpha} \rangle > 0 \end{array} \right\}$$

and $\overline{\mathbb{C}}$ be the closure of \mathbb{C} in ${}^a\mathfrak{h}$.

We identify the set of infinitesimal characters with ${}^a\mathfrak{h}^*/W$.

1.1. Coherent family. For each finite dimensional \mathfrak{g} -module or $G_{\mathbb{C}}$ -module F , let F^* be its contragredient representation and let $\Delta(F) \subseteq {}^aX$ denote the multi-set of weights in F .

Let $\Pi_{\Lambda_0}(G_{\mathbb{C}})$ be the set of irreducible finite dimensional representations of $G_{\mathbb{C}}$ with external weight in Λ_0 and $\mathcal{G}_{\Lambda}(G_{\mathbb{C}})$ be the subgroup generated by $\Pi_{\Lambda_0}(G_{\mathbb{C}})$. Let

$${}^aP := \{ \mu \in {}^aX \mid \mu \text{ is a } {}^a\mathfrak{h}\text{-weight of an } F \in \Pi_{\text{fin}}(G_{\mathbb{C}}) \}.$$

Via the highest weight theory, every W -orbit $W \cdot \mu$ in aP corresponds with the irreducible finite dimensional representation $F \in \Pi_{\text{fin}}(G_{\mathbb{C}})$ with external weight μ .

Now the Grothendieck group $\mathcal{G}(G_{\mathbb{C}})$ of finite dimensional representation of $G_{\mathbb{C}}$ is identified with $\mathbb{Z}[{}^aP/W]$. In fact $\mathcal{G}(G_{\mathbb{C}})$ is a \mathbb{Z} -algebra under the tensor product and equipped with the involution $F \mapsto F^*$.

Fix a W -invariant sub-lattice $\Lambda_0 \subset {}^aX$ containing aQ .

2000 *Mathematics Subject Classification.* 22E45, 22E46.

Key words and phrases. orbit method, unitary dual, unipotent representation, classical group, theta lifting, moment map.

Take a lattice $\Lambda = \lambda + \Lambda_0 \in \mathfrak{h}^*/\Lambda$ with $\lambda \in \overline{\mathbb{C}}$. Let

$$R(\lambda) := \{ \alpha \in {}^a R \mid \langle \lambda, \check{\alpha} \rangle \in \mathbb{Z} \} \quad \text{and} \quad W(\lambda) := \langle s_\alpha \mid \alpha \in R(\lambda) \rangle.$$

Definition 1.1. Suppose \mathcal{M} is an abelian group with $\mathcal{G}_\Lambda(G_\mathbb{C})$ -action

$$\mathcal{G}_\Lambda(G_\mathbb{C}) \times \mathcal{M} \ni (F, m) \mapsto F \otimes m.$$

In addition, we fix a subgroup $\mathcal{M}_{[\mu]}$ of \mathcal{M} for each W -coset $[\mu] \in \Lambda/W$.

A function $f: \Lambda \rightarrow \mathcal{M}$ is called a coherent family based on Λ if it satisfies $f(\mu) \in \mathcal{M}_\mu$ and

$$F \otimes f(\mu) = \sum_{\nu \in \Delta(F)} f(\mu + \nu) \quad \forall \mu \in \Lambda, F \in \Pi_{\Lambda_0}(G_\mathbb{C}).$$

Let $\text{Coh}_\Lambda(\mathcal{M})$ be the abelian group of all coherent families based on Λ and value in \mathcal{M} .

In this paper, we will consider the following cases.

Example 1.2. Suppose $\mathcal{M} = \mathbb{Q}$ and $F \otimes m = \dim(F) \cdot m$ for $F \in \Pi_{\Lambda_0}(G_\mathbb{C})$ and $m \in \mathcal{M}$. We let $\mathcal{M}_\mu = \mathcal{M}$ for every $\mu \in \Lambda$. When $\Lambda = \Lambda_0$, the set of W -harmonic polynomials on ${}^a \mathfrak{h}^*$ is naturally identified with $\text{Coh}_\Lambda(\mathcal{M})$ via restriction (Vogan's result)

Example 1.3. Let $\mathcal{G}(\mathfrak{g}, K)$ be the Grothendieck group of finite length (\mathfrak{g}, K) -modules and $\mathcal{G}_\mu(\mathfrak{g}, K)$ be the subgroup of $\mathcal{G}(\mathfrak{g}, K)$ generated by the set of irreducible (\mathfrak{g}, K) -modules with infinitesimal character μ .

Then $\text{Coh}_\Lambda(\mathcal{G}(\mathfrak{g}, K))$ is the group of coherent families of Harish-Chandra modules.

Example 1.4. Fixing a Borel subalgebra $\mathfrak{b} \subset \mathfrak{g}$, let $\mathcal{G}(\mathfrak{g}, \mathfrak{b})$ be the Grothendieck group of the category $\mathcal{O}^{\mathfrak{b}}$. The space $\text{Coh}_\Lambda(\mathcal{G}(\mathfrak{g}, \mathfrak{b}))$ is defined similarly.

Note that the lattice Λ is stable under the $W(\lambda)$ action. We can define $W(\lambda)$ action on $\text{Coh}_\Lambda(\mathcal{M})$ by

$$w \cdot f(\mu) = f(w^{-1}\mu) \quad \forall \mu \in \Lambda, w \in W(\lambda).$$

Translation principal assumption. Recall that we have fixed a set of simple roots of W_Λ . We make the following assumption for $\text{Coh}_\Lambda(\mathcal{M})$.

- There is a basis $\{\Theta_\gamma \mid \gamma \in \text{Parm}\}$ of $\text{Coh}_\Lambda(\mathcal{M})$ where Parm is a parameter set;
- For every $\mu \in \Lambda$, the evaluation at μ is surjective;

$$\text{ev}_\mu: \text{Coh}_\Lambda(\mathcal{M}) \rightarrow \mathcal{M}_{[\mu]} \quad f \mapsto f(\mu)$$

is surjective;

- a subset $\tau(\gamma)$ of the simple roots of W_Λ is attached to each $\gamma \in \text{Parm}$ such that $s \cdot \Theta_\gamma = -\Theta_\gamma$;
- $\Theta_\gamma(\mu) = 0$ if and only if $\tau(\gamma) \cap R_\mu \neq \emptyset$.
- $\{\Theta_\gamma(\mu) \mid \tau(\gamma) \cap R_\mu = \emptyset\}$ form a basis of \mathcal{M}_γ .

The translation principle assumption implies

$$(1.1) \quad \text{Ker ev}_\mu = \text{span} \{ \Theta_\gamma \mid \tau(\gamma) \cap R_\mu \neq \emptyset \}$$

Now we have the following counting lemma.

Lemma 1.5. For each μ , we have

$$\dim \mathcal{M}_{[\mu]} = \dim(\text{Coh}_\Lambda(\mathcal{M}))_{W_\mu} = [\text{Coh}_\Lambda(\mathcal{M}), 1_{W_\mu}].$$

Proof. Clearly

$$\text{span} \{ \Theta_\gamma - w\Theta_\gamma \mid \gamma \in \text{Parm}, w \in W_\Lambda \} \subseteq \text{ker ev}_\mu$$

since $w \cdot \Theta_\gamma(\mu) = \Theta_\gamma(w^{-1} \cdot \mu) = \Theta_\gamma(\mu)$ for $w \in W_\mu$. Combine this with (1.1), we conclude that

$$\text{span} \{ \Theta_\gamma - w\Theta_\gamma \mid \gamma \in \text{Parm}, w \in W_\Lambda \} = \text{ker ev}_\mu.$$

Therefore, ev_μ induces an isomorphism $(\text{Coh}_\Lambda(\mathcal{M}))_{W_\mu} \rightarrow \mathcal{M}_{[\mu]}$. Now the dimension equality follows. \square

Example 1.6. For the case of category \mathcal{O} . We can take $\text{Parm} = W_\Lambda$. Let $\Theta_w(\mu) = L(w\mu)$ for each $w \in W_\Lambda$ with $\tau(w) = \{s_\alpha \mid \alpha \in \Delta^+, w\alpha \notin R^+\}$.

Example 1.7. In the Harish-Chandra module case, Parm consists of certain irreducible K -equivariant local systems on a K -orbit of the flag variety of G . $\tau(\gamma)$ is the τ -invariants of the parameter γ .

2. PARAMETERIZE OF UNIPOTENT REPRESENTATIONS

We fix an abstract complex Cartan subgroup \mathbf{H}_a and \mathfrak{h}_a in \mathbf{G} and a set of simple roots Π_a . Let $\mathcal{P}(\mathbf{G})$ be the set of all Langlands parameters of G -modules with character ρ (i.e. the infinitesimal character of the trivial representation). For $\gamma \in \mathcal{P}(\mathbf{G})$, let $\mathcal{L}(\gamma)$, $\mathcal{S}(\gamma)$ and Φ_γ be the corresponding Langlands quotient, standard module and coherent family such that $\Phi_\gamma(\rho) = \mathcal{L}(\gamma)$. Let $\mathcal{M}(\mathbf{G})$ be the span of $\mathcal{L}(\gamma)$. Let $\{\mathcal{B}\}$ be the set of all blocks. Then $\mathcal{P}(\mathbf{G}) = \bigsqcup_{\mathcal{B}} \mathcal{B}$. The Weyl group $W = W(G)$ acts on $\mathcal{M}(\mathbf{G})$ by coherent continuation. Let $\mathcal{M}_{\mathcal{B}}$ be the submodule of $\mathcal{M}(\mathbf{G})$ spanned by $\gamma \in \mathcal{B}$, then

$$\mathcal{M}(\mathbf{G}) = \bigoplus_{\mathcal{B}} \mathcal{M}_{\mathcal{B}}$$

Let $\tau(\gamma) \subset \Pi_a$ be the τ -invariant of γ .

Let \mathcal{O} be even orbit. $\lambda = \frac{1}{2}\check{h}$. Define

$$S(\lambda) = \{ \alpha \in \Pi_a \mid \langle \alpha, \lambda \rangle = 0 \}.$$

Let $\mathcal{P}_\lambda(\mathbf{G})$ be the set of all Langlands parameters with infinitesimal character λ . Let $T_{\lambda,\rho}$ be the translation functor. Let

$$\mathcal{B}(S) = \{ \gamma \in \mathcal{B} \mid S \cap \tau(\gamma) = \emptyset \}$$

and

$$\mathcal{P}(\mathbf{G}, S) = \bigsqcup_{\mathcal{B}} \mathcal{B}(S)$$

Then

$$\begin{aligned} \mathcal{P}(\mathbf{G}, S) &\longrightarrow \mathcal{P}_\lambda(\mathbf{G}) \\ \gamma &\longmapsto T_{\lambda,\rho}(\gamma) \end{aligned}$$

Let \mathcal{O} be a complex nilpotent orbit in \mathfrak{g} . Let

$$\mathcal{B}(S, \mathcal{O}) = \{ \gamma \in \mathcal{B}(S) \mid \text{AV}_{\mathbb{C}}(\mathcal{L}(\gamma)) \subset \overline{\mathcal{O}} \}$$

Let

$$\begin{aligned} m_S(\sigma) &= [\sigma : \text{Ind}_{W(S)}^W \mathbf{1}] \\ m_{\mathcal{B}}(\sigma) &= [\sigma : \mathcal{M}_{\mathcal{B}}] \end{aligned}$$

Barbasch [10, Theorem 9.1] established the following theorem.

Theorem 2.1.

$$|\mathcal{B}(S, \mathcal{O})| = \sum_{\sigma} m_{\mathcal{B}}(\sigma) m_S(\sigma)$$

Here $\sigma \times \sigma$ running over the $W \times W$ appears in the double cell $\mathcal{C}(\mathcal{O})$.

Proof. We need to take the graded module of $\mathcal{M}(\mathbf{G})$ with respect to the $\overset{LR}{\leq}$. By abuse of notation, we identify the basis $\mathcal{P}(\mathbf{G})$ with its image in the graded module. Note that $S \cap \tau(\lambda) = \emptyset$ if and only if $W(S)$ acts on γ trivially by [70, Lemma 14.7]. On the other hand, by [70, Theorem 14.10, and page 58], $\text{AV}_{\mathbb{C}}(\mathcal{L}(\gamma)) \subset \overline{\mathcal{O}}$ only if γ generate a W -module in the double cell of \mathcal{O} . \square

Now assume $S = S(\lambda)$. By [12, Cor 5.30 b) and c)], $[\sigma : \text{Ind}_{W(S)}^W \mathbf{1}] = [\mathbf{1}|_{W(S)} : \sigma] \leq 1$.

3. COUNTING IN TYPE A

The results in this section are well known to the experts.

Let \mathbf{YD} be the set of Young diagrams viewed as a finite multiset of positive integers. The set of nilpotent orbits in $\text{GL}_n(\mathbb{C})$ is identified with Young diagram of n boxes.

Let $\check{G}_{\mathbb{C}} = \text{GL}_n(\mathbb{C})$. Fix an orbit $\check{\mathcal{O}} \in \text{Nil}(ckG)$, let $\check{\mathcal{O}}_e$ (resp. $\check{\mathcal{O}}_o$) be the partition consists of all even (resp. odd) rows in $\check{\mathcal{O}}$.

Let S_n denote the Weyl group of $\text{GL}_n(\mathbb{C})$. Let $W_n := S_n \ltimes \{\pm 1\}^n$ denote the Weyl group of type B_n or C_n . Let sgn denote the sign representation of the Weyl group. The group W_n is naturally embeded in S_{2n} . For W_n , let ϵ denote the unique non-trivial character which is trivial on S_n . Note that ϵ is also the restriction of the sgn of S_{2n} on W_n .

3.1. Special unipotent representations of $G = \text{GL}_n(\mathbb{C})$. By [12], the set of unipotent representations of $G = \text{GL}_n(\mathbb{C})$ one-one corresponds to nilpotent orbits in $\text{Nil}(\check{G}_{\mathbb{C}})$. Suppose $\check{\mathcal{O}}$ has rows

$$\mathbf{r}_1(\check{\mathcal{O}}) \geq \mathbf{r}_2(\check{\mathcal{O}}) \geq \cdots \geq \mathbf{r}_k(\check{\mathcal{O}}) > 0.$$

Then $\mathcal{O} := d_{\text{BV}}(\check{\mathcal{O}})$ has columns $\mathbf{c}_i(\mathcal{O}) = \mathbf{r}_i(\check{\mathcal{O}})$ for all $i \in \mathbb{N}^+$. The map $\check{\mathcal{O}} \mapsto \mathcal{O}$ is a bijection.

We set $\mathbf{CP} = \mathbf{YD}$ be the set of Young diagrams. For $\tau \in \mathbf{CP}$ which has k columns, let $1_{\mathbf{c}_k}$ be the trivial representations of $\text{GL}_{\mathbf{c}_k}(\mathbb{C})$.

$$\pi_{\tau} = 1_{\mathbf{c}_1(\tau)} \times 1_{\mathbf{c}_2(\tau)} \times \cdots \times 1_{\mathbf{c}_k(\tau)}.$$

The Vogan duality gives a duality between Harish-Chandra cells. In this case, Harish-Chandra cells is the double cell of Lusztig. Now we have a duality

$$\pi_{\tau} \leftrightarrow \pi_{\tau^t}.$$

Let $\tau' := \nabla(\tau)$ be the partition obtained by deleting the first column of τ . Let $\theta_{a,b}$ (resp. $\Theta_{a,b}$) be the theta lift (resp. big theta lift) from $\text{GL}_a(\mathbb{C})$ to $\text{GL}_b(\mathbb{C})$. Then we have

$$\pi_{\tau} = \theta_{|\tau'|, |\tau|}(\pi_{\tau}).$$

3.2. Counting unipotent representations of $\text{GL}_n(\mathbb{R})$. Now let $\check{\mathcal{O}} \in \text{Nil}(\check{G}_{\mathbb{C}})$. Recall the decomposition $\check{\mathcal{O}} = \check{\mathcal{O}}_e \cup \check{\mathcal{O}}_o$. Let $n_e = |\check{\mathcal{O}}_e|$, $n_o = |\check{\mathcal{O}}_o|$ and $\lambda_{\check{\mathcal{O}}} = \frac{1}{2}\check{h}$.

Then

$$\begin{aligned} W(\lambda_{\check{\mathcal{O}}}) &\cong S_{|\check{\mathcal{O}}_e|} \times S_{|\check{\mathcal{O}}_o|}. \\ W_{\lambda_{\check{\mathcal{O}}}} &= \prod_j S_{\mathbf{c}_j(\check{\mathcal{O}}_e)} \times \prod_j S_{\mathbf{c}_j(\check{\mathcal{O}}_o)} \end{aligned}$$

By the formula of a -function, one can easily see that The cell in $W(\lambda_{\check{\mathcal{O}}})$ consists of the unique representatoin $J_{W_{\lambda_{\check{\mathcal{O}}}}}^{W[\lambda_{\check{\mathcal{O}}}]}(1)$. Now the W -cell is $J_{W_{\lambda_{\check{\mathcal{O}}}}}^W 1$ corresponds to the orbit $\mathcal{O} = \check{\mathcal{O}}^t$ under the Springer correspondence. [**WLOG, we assume $\check{\mathcal{O}} = \check{\mathcal{O}}_o$.**

Let $\sigma \in \widehat{S}_n$. We identify σ with a Young diagram. Let $c_i = \mathbf{c}_i(\sigma)$. Then $\sigma = J_{W'}^{S_n} \epsilon_{W'}$ where $W' = \prod S_{c_i}$ (see Carter's book). This implies Lusztig's a -function takes value

$$a(\sigma) = \sum_i c_i(c_i - 1)/2$$

Comparing the above with the dimension formula of nilpotent Orbits [17, Collary 6.1.4], we get (for the formula, see Bai ZQ-Xie Xun's paper on GK dimension of $SU(p, q)$)

$$\frac{1}{2} \dim(\sigma) = \dim(L(\lambda)) = n(n-1)/2 - a(\sigma).$$

Here $\dim(\sigma)$ is the dimension of nilpotent orbit attached to the Young diagram of σ (it is the Springer correspondence, regular orbit maps to trivial representation, note that $a(\text{triv}) = 0$), $L(\lambda)$ is any highest weight module in the cell of σ .

Return to our question, let $S' = \prod_i S_{\mathbf{c}_i(\check{\sigma})}$. We want to find the component σ_0 in $\text{Ind}_{S'}^{S_n} 1$ whose $a(\sigma_0)$ is maximal, i.e. the Young diagram of σ_0 is minimal.

By the branching rule, $\sigma \subset \text{Ind}_{S'}^{S_n} 1$ is given by adding rows of length $\mathbf{c}_i(\check{\sigma})$ repeatedly (Each time add at most one box in each column). Now it is clear that $\sigma_0 = \check{\sigma}^t$ is desired.

This agrees with the Barbasch-Vogan duality d_{BV} given by

$$\check{\sigma} \xrightarrow{\text{Springer}} \check{\sigma} \xrightarrow{\otimes \text{sgn}} \check{\sigma}^t \xrightarrow{\text{Springer}} \check{\sigma}^t.$$

]

The $W_{[\lambda_{\check{\sigma}}]}$ -module $\text{Coh}_{[\lambda_{\check{\sigma}}]}$ is given by the following formula:

$$\begin{aligned} \text{Coh}_{[\lambda_{\check{\sigma}}]} &\cong \mathcal{C}_{n_e} \otimes \mathcal{C}_{n_o} \quad \text{with} \\ \mathcal{C}_n &:= \bigoplus_{\substack{s,a,b \\ 2s+a+b=n}} \text{Ind}_{W_s \times S_a \times S_b}^{S_n} \epsilon \otimes 1 \otimes 1. \end{aligned}$$

According to Vogan duality, we can obtain the above formula by tensoring sgn on the formula of the unitary groups in [8, Section 4].

By branching rules of the symmetric groups, $\text{Unip}_{\check{\sigma}}(G)$ can be parameterized by painted partition.

$$\text{PP}_{\check{\sigma}} = \{ \tau: \text{Box}(\check{\sigma}^t) \rightarrow \{ \bullet, c, d \} \mid \begin{array}{l} \text{"}\bullet\text{" occurs with even} \\ \text{multiplicity in each column} \end{array} \}.$$

[The typical diagram of all columns with even length $2c$ are

•	...	•	•	...	•
⋮	⋮	⋮	⋮	⋮	⋮
•	...	•	c	...	c
•	...	•	d	...	d

The typical diagram of all columns with odd length $2c + 1$ are

•	...	•	•	...	•
⋮	⋮	⋮	⋮	⋮	⋮
•	...	•	•	...	•
c	...	c	d	...	d

]

Let $\text{sgn}_n: \text{GL}_n(\mathbb{R}) \rightarrow \{ \pm 1 \}$ be the sign of determinant. Let 1_n be the trivial representation of $\text{GL}_n(\mathbb{R})$. For $\tau \in \text{PP}_{\check{\sigma}}$, we attach the representation

$$(3.1) \quad \pi_\tau := \bigtimes_j \underbrace{1_j \times \cdots \times 1_j}_{c_j\text{-terms}} \times \underbrace{\text{sgn}_j \times \cdots \times \text{sgn}_j}_{d_j\text{-terms}}.$$

Here

- j running over all column lengths in $\check{\mathcal{O}}^t$,
- d_j is the number of columns of length j ending with the symbol “d”,
- c_j is the number of columns of length j ending with the symbol “•” or “c”, and
- “ \times ” denote the parabolic induction.

3.3. Special Unipotent representations of $G = \mathrm{GL}_m(\mathbb{H})$. Suppose that \mathcal{O} is the complexification of a rational nilpotent $\mathrm{GL}_m(\mathbb{H})$ -orbit. Then \mathcal{O} has only even length columns. Therefore, $\mathrm{Unip}_{\check{\mathcal{O}}}(G) \neq \emptyset$ only if $\check{\mathcal{O}} = \check{\mathcal{O}}_e$.

In this case the coherent continuation representation is given by

$$\mathrm{Coh}_{[\lambda_{\check{\mathcal{O}}}]}(G) = \mathrm{Ind}_{W_m}^{S_{2m}} \epsilon$$

and $\mathrm{Unip}_{\check{\mathcal{O}}}(G)$ is a singleton. For each partition τ only having even columns, we define

$$\pi_{\tau} := \bigotimes_i 1_{\mathbf{c}_i(\tau)/2}.$$

3.4. Counting special unipotent representations of $\mathrm{U}(p, q)$. We call the parity of $|\check{\mathcal{O}}|$ the “good parity”. The other parity is called the “bad parity”. We write $\check{\mathcal{O}} = \check{\mathcal{O}}_g \cup \check{\mathcal{O}}_b$ where $\check{\mathcal{O}}_g$ and $\check{\mathcal{O}}_b$ consist of good parity length rows and bad parity rows respectively.

Let $(n_g, n_b) = (|\check{\mathcal{O}}_g|, |\check{\mathcal{O}}_b|)$. Now as the $S_{n_g} \times S_{n_b}$

$$\bigoplus_{\substack{p, q \in \mathbb{N} \\ p+q=n}} \mathrm{Coh}_{[\lambda_{\check{\mathcal{O}}}]}(\mathrm{U}(p, q)) = \mathcal{C}_{gd} \otimes \mathcal{C}_{bd}$$

where

$$\begin{aligned} \mathcal{C}_{gd} &= \bigoplus_{\substack{s, a, b \in \mathbb{N} \\ 2s+a+b=n_g}} \mathrm{Ind}_{W_s \times S_a \times S_b}^{S_{n_g}} 1 \otimes \mathrm{sgn} \otimes \mathrm{sgn} \\ \mathcal{C}_{bd} &= \begin{cases} \mathrm{Ind}_{W_{\frac{n_b}{2}}}^{S_{n_b}} 1 & \text{if } n_b \text{ is even} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

By the above formula, we have

Lemma 3.1. (i) The set $\mathrm{Unip}_{\check{\mathcal{O}}_b}(\mathrm{U}(p, q)) \neq \emptyset$ if and only if $p = q$ and each row length in $\check{\mathcal{O}}$ has even multiplicity.

(ii) Suppose $\mathrm{Unip}_{\check{\mathcal{O}}_b}(\mathrm{U}(p, p)) \neq \emptyset$, let $\check{\mathcal{O}}'$ be the Young diagram such that $\mathbf{r}_i(\check{\mathcal{O}}') = \mathbf{r}_{2i}(\check{\mathcal{O}}_b)$ and π' be the unique special unipotent representation in $\mathrm{Unip}_{\check{\mathcal{O}}'}(\mathrm{GL}_p(\mathbb{C}))$. Then the unique element in $\mathrm{Unip}_{\check{\mathcal{O}}_b}(\mathrm{U}(p, p))$ is given by

$$\pi := \mathrm{Ind}_P^{\mathrm{U}(p, p)} \pi'$$

where P is a parabolic subgroup in $\mathrm{U}(p, p)$ with Levi factor equals to $\mathrm{GL}_p(\mathbb{C})$.

(iii) In general, when $\mathrm{Unip}_{\check{\mathcal{O}}_b}(\mathrm{U}(p, p)) \neq \emptyset$, we have a natural bijection

$$\begin{aligned} \mathrm{Unip}_{\check{\mathcal{O}}_g}(\mathrm{U}(n_1, n_2)) &\longrightarrow \mathrm{Unip}_{\check{\mathcal{O}}}(\mathrm{U}(n_1 + p, n_2 + p)) \\ \pi_0 &\longmapsto \mathrm{Ind}_P^{\mathrm{U}(n_1+p, n_2+p)} \pi' \otimes \pi_0 \end{aligned}$$

where P is a parabolic subgroup with Levi factor $\mathrm{GL}_p(\mathbb{C}) \times \mathrm{U}(n_1, n_2)$.

The above lemma ensure us to reduce the problem to the case when $\check{\mathcal{O}} = \check{\mathcal{O}}_g$. Now assume $\check{\mathcal{O}} = \check{\mathcal{O}}_g$ and so $\mathrm{Coh}_{[\check{\mathcal{O}}]}$ corresponds to the blocks of the infinitesimal character of the trivial representation.

By [8, Theorem 4.2], Harish-Chandra cells in $\mathrm{Coh}_{[\check{\mathcal{O}}]}$ are in one-one correspondence to real nilpotent orbits in $\mathcal{O} := d_{\mathrm{BV}}(\check{\mathcal{O}}) = \check{\mathcal{O}}^t$.


[From the branching rule, the cell is parametered by painted partition

$$\text{PP}(\mathbf{U}) := \{ \tau \in \text{PP} \mid \begin{array}{l} \text{Im}(\tau) \subseteq \{ \bullet, s, r \} \\ \text{"}\bullet\text{" occurs even times in each row} \end{array} \}.$$

The bijection $\text{PP}(\mathbf{U}) \rightarrow \text{SYD}, \tau \mapsto \mathcal{O}$ is given by the following recipe: The shape of \mathcal{O} is the same as that of τ . \mathcal{O} is the unique (upto row switching) signed Young diagram such that

$$\mathcal{O}(i, \mathbf{r}_i(\tau)) := \begin{cases} +, & \text{when } \tau(i, \mathbf{r}_i(\tau)) = r; \\ -, & \text{otherwise, i.e. } \tau(i, \mathbf{r}_i(\tau)) \in \{ \bullet, s \}. \end{cases}$$

Example 3.2.



]

Now the following lemma is clear.

Lemma 3.3. *When $\check{\mathcal{O}} = \check{\mathcal{O}}_g$, the associated variety of every special unipotent representations in $\text{Unip}_{\check{\mathcal{O}}}(\mathbf{U})$ is irreducible. Moreover, the following map is a bijection.*

$$\begin{array}{ccc} \text{Unip}_{\check{\mathcal{O}}_g}(\mathbf{U}(n_1, n_2)) & \longrightarrow & \{ \text{rational forms of } \check{\mathcal{O}}^t \} \\ \pi_0 & \mapsto & \text{AV}^{\text{weak}}(\pi_0). \end{array}$$

□

Remark. When $\check{\mathcal{O}}_b \neq \emptyset$, the special unipotent representations can have reducible associated variety. Meanwhile, it is easy to see that the map $\text{Unip}_{\check{\mathcal{O}}}(\mathbf{U}) \ni \pi \mapsto \text{AV}^{\text{weak}}(\pi)$ is still injective.

We will show that every elements in $\text{Unip}_{\check{\mathcal{O}}_g}$ can be constructed by iterated theta lifting. For each τ , let \mathcal{O} be the corresponding real nilpotent orbit. Let $\text{Sign}(\mathcal{O})$ be the signature of \mathcal{O} , $\nabla(\mathcal{O})$ be the signed Young diagram obtained by deleting the first column of \mathcal{O} . Suppose \mathcal{O} has k -columns. Inductively we have a sequence of unitary groups $\mathbf{U}(p_i, q_i)$ with $(p_i, q_i) = \text{Sign}(\nabla^i(\mathcal{O}))$ for $i = 0, \dots, k$. Then

$$(3.2) \quad \pi_\tau = \theta_{\mathbf{U}(p_1, q_1)}^{\mathbf{U}(p_0, q_0)} \theta_{\mathbf{U}(p_2, q_2)}^{\mathbf{U}(p_1, q_1)} \cdots \theta_{\mathbf{U}(p_k, q_k)}^{\mathbf{U}(p_{k-1}, q_{k-1})}(1)$$

where 1 is the trivial representation of $\mathbf{U}(p_k, q_k)$.

Suppose $\check{\mathcal{O}} = \check{\mathcal{O}}_g$. Form the duality between cells of $\mathbf{U}(p, q)$ and $\text{GL}(n, \mathbb{R})$. We have an ad-hoc (bijective) duality between unipotent representations:

$$\begin{array}{ccc} d_{\text{BV}}: \text{Unip}_{\check{\mathcal{O}}}(\mathbf{U}) & \rightarrow & \text{Unip}_{\check{\mathcal{O}}^t}(\text{GL}(\mathbb{R})) \\ \pi_\tau & \mapsto & \pi_{d_{\text{BV}}(\tau)} \end{array}$$

Here $\check{\mathcal{O}}^t = d_{\text{BV}}(\check{\mathcal{O}})$ and $d_{\text{BV}}(\tau)$ is the painted bipartition obtained by transposing τ and replace s and r by c and d respectively. See (3.2) and (3.1) for the definition of special unipotent representations on the two sides.

4. COUNTING IN TYPE BCD

In this section, we count special unipotent representations of real orthogonal groups (type B, D), symplectic groups (type C) and metaplectic groups (type \tilde{C}).

Recall that

$$\text{good parity} = \begin{cases} \text{odd} & \text{in type } C \text{ and } D \\ \text{even} & \text{in type } C \text{ and } D \end{cases}$$

We decompose $\check{\mathcal{O}} = \check{\mathcal{O}}_g \cup \check{\mathcal{O}}_b$ as before. We also set $(2m+1, 2m') = (|\check{\mathcal{O}}_g|, |\check{\mathcal{O}}_b|)$

4.1. Counting special unipotent representation of $G = \text{Sp}(2n, \mathbb{R})$. The dual group of $G_{\mathbb{C}}$ is $\check{G}_{\mathbb{C}} = \text{SO}(2n+1, \mathbb{C})$. By Vogan duality, the dual group for class $[\lambda_{\check{\mathcal{O}}}]$ is

$$\text{SO}(n_g, \mathbb{C}) \times \text{SO}(n_b, \mathbb{C}).$$

Now we have

$$\text{Coh}_{[\lambda_{\check{\mathcal{O}}}]}(G) \otimes \text{sgn} = \mathcal{C}_g \otimes \mathcal{C}_b$$

with

$$\begin{aligned} \mathcal{C}_g &= \bigoplus_{p+q=2m+1} \text{Coh}_{[\rho]}(\text{SO}(p, q)) \\ &= \bigoplus_{2b+s+r+c+d=m} \text{Ind}_{W_b \ltimes \{-1\}^b \times W_s \times W_r \times W_c \times W_d}^{W_m} 1 \otimes \det \otimes \det \otimes 1 \otimes 1 \\ \mathcal{C}_b &= \text{Coh}_{[\rho]}(\text{SO}^*(2m)) = \bigoplus_{2s+b=m} \text{Ind}_{W_s \ltimes \{-1\}^s \times S_b}^{W_m} 1 \otimes \text{sgn} \otimes \text{sgn} \end{aligned}$$

REFERENCES

- [1] J. Adams, *Discrete spectrum of the reductive dual pair $(O(p, q), \text{Sp}(2m))$* , Invent. Math. **74** (1983), no. 3, 449–475.
- [2] J. Adams, D. Barbasch, and D. A. Vogan, *The Langlands classification and irreducible characters for real reductive groups*, Progress in Math., vol. 104, Birkhauser, 1991.
- [3] Jeffrey Adams and Fokko du Cloux, *Algorithms for representation theory of real reductive groups*, Journal of the Institute of Mathematics of Jussieu **8** (2009), no. 2, 209–259, DOI 10.1017/S1474748008000352.
- [4] J. Arthur, *On some problems suggested by the trace formula*, Lie group representations, II (College Park, Md.), Lecture Notes in Math. 1041 (1984), 1–49.
- [5] ———, *Unipotent automorphic representations: conjectures*, Orbites unipotentes et représentations, II, Astérisque **171-172** (1989), 13–71.
- [6] L. Auslander and B. Kostant, *Polarizations and unitary representations of solvable Lie groups*, Invent. Math. **14** (1971), 255–354.
- [7] D. Barbasch, *Unipotent representations for real reductive groups*, Proceedings of ICM (1990), Kyoto (2000), 769–777.
- [8] Dan Barbasch and David Vogan, *Weyl Group Representations and Nilpotent Orbits*, Representation Theory of Reductive Groups: Proceedings of the University of Utah Conference 1982, 1983, pp. 21–33.
- [9] D. Barbasch, *Orbital integrals of nilpotent orbits*, The mathematical legacy of Harish-Chandra, Proc. Sympos. Pure Math. **68** (2000), 97–110.
- [10] ———, *The unitary spherical spectrum for split classical groups*, J. Inst. Math. Jussieu **9** (2010), 265–356.
- [11] ———, *Unipotent representations and the dual pair correspondence*, J. Cogdell et al. (eds.), Representation Theory, Number Theory, and Invariant Theory, In Honor of Roger Howe. Progress in Math. **323** (2017), 47–85.
- [12] D. Barbasch and D. A. Vogan, *Unipotent representations of complex semisimple groups*, Annals of Math. **121** (1985), no. 1, 41–110.
- [13] R. Brylinski, *Dixmier algebras for classical complex nilpotent orbits via Kraft-Procesi models. I*, The orbit method in geometry and physics (Marseille, 2000). Progress in Math. **213** (2003), 49–67.
- [14] Roger W. Carter, *Finite groups of Lie type*, Wiley Classics Library, John Wiley & Sons, Ltd., Chichester, 1993.

- [15] W. Casselman, *Canonical extensions of Harish-Chandra modules to representations of G* , Canad. J. Math. **41** (1989), 385–438.
- [16] F. Du Cloux, *Sur les représentations différentiables des groupes de Lie algébriques*, Ann. Sci. École Norm. Sup. **24** (1991), no. 3, 257–318.
- [17] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebra: an introduction*, Van Nostrand Reinhold Co., 1993.
- [18] A. Daszkiewicz, W. Kraśkiewicz, and T. Przebinda, *Nilpotent orbits and complex dual pairs*, J. Algebra **190** (1997), no. 2, 518 – 539.
- [19] ———, *Dual pairs and Kostant-Sekiguchi correspondence. II. Classification of nilpotent elements*, Central European J. Math. **3** (2005), 430–474.
- [20] J. Dixmier and P. Malliavin, *Factorisations de fonctions et de vecteurs indéfiniment différentiables*, Bull. Sci. Math. (2) **102** (1978), 307–330.
- [21] M. Duflo, *Théorie de Mackey pour les groupes de Lie algébriques*, Acta Math. **149** (1982), no. 3-4, 153–213.
- [22] R. Gomez and C.-B. Zhu, *Local theta lifting of generalized Whittaker models associated to nilpotent orbits*, Geom. Funct. Anal. **24** (2014), no. 3, 796–853.
- [23] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique IV: Étude locale des schémas et des morphismes de schémas. II*, Inst. Hautes Études Sci. Publ. Math. **24** (1965).
- [24] ———, *Éléments de géométrie algébrique IV: Étude locale des schémas et des morphismes de schémas. III*, Inst. Hautes Études Sci. Publ. Math. **28** (1966).
- [25] M. Harris, J.-S. Li, and B. Sun, *Theta correspondences for close unitary groups*, Arithmetic Geometry and Automorphic Forms, Adv. Lect. Math. (ALM) **19** (2011), 265–307.
- [26] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics, 52. New York-Heidelberg-Berlin: Springer-Verlag, 1983.
- [27] H. He, *Unipotent representations and quantum induction*, arXiv:math/0210372 (2002).
- [28] J.-S. Huang and J.-S. Li, *Unipotent representations attached to spherical nilpotent orbits*, Amer. J. Math. **121** (1999), no. 3, 497–517.
- [29] J.-S. Huang and C.-B. Zhu, *On certain small representations of indefinite orthogonal groups*, Represent. Theory **1** (1997), 190–206.
- [30] R. Howe, *θ -series and invariant theory*, Automorphic Forms, Representations and L -functions, Proc. Sympos. Pure Math, vol. 33, 1979, pp. 275–285.
- [31] ———, *On a notion of rank for unitary representations of the classical groups*, Harmonic analysis and group representations, Liguori, Naples (1982), 223–331.
- [32] ———, *Transcending classical invariant theory*, J. Amer. Math. Soc. **2** (1989), 535–552.
- [33] ———, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, Piatetski-Shapiro, I. et al. (eds.), The Schur lectures (1992). Ramat-Gan: Bar-Ilan University, Isr. Math. Conf. Proc. 8, (1995), 1–182.
- [34] D. Jiang, B. Liu, and G. Savin, *Raising nilpotent orbits in wave-front sets*, Represent. Theory **20** (2016), 419–450.
- [35] A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Uspehi Mat. Nauk **17** (1962), 57–110.
- [36] B. Kostant, *Quantization and unitary representations*, Lectures in Modern Analysis and Applications III, Lecture Notes in Math. **170** (1970), 87–208.
- [37] H. Kraft and C. Procesi, *On the geometry of conjugacy classes in classical groups*, Comment. Math. Helv. **57** (1982), 539–602.
- [38] S. S. Kudla and S. Rallis, *Degenerate principal series and invariant distributions*, Israel J. Math. **69** (1990), 25–45.
- [39] S. S. Kudla, *Some extensions of the Siegel-Weil formula*, In: Gan W., Kudla S., Tschinkel Y. (eds) Eisenstein Series and Applications. Progress in Mathematics, vol 258. Birkhäuser Boston (2008), 205–237.
- [40] S. T. Lee and C.-B. Zhu, *Degenerate principal series and local theta correspondence II*, Israel J. Math. **100** (1997), 29–59.
- [41] ———, *Degenerate principal series of metaplectic groups and Howe correspondence*, D. Prasad et al. (eds.), Automorphic Representations and L-Functions, Tata Institute of Fundamental Research, India, (2013), 379–408.
- [42] J.-S. Li, *Singular unitary representations of classical groups*, Invent. Math. **97** (1989), no. 2, 237–255.
- [43] Q. Liu, *Algebraic Geometry and Arithmetic Curves*, Oxford University Press, 2006.
- [44] H. Y. Loke and J. Ma, *Invariants and K -spectrums of local theta lifts*, Compositio Math. **151** (2015), 179–206.

- [45] G. Lusztig and N. Spaltenstein, *Induced unipotent classes*, j. London Math. Soc. **19** (1979), 41–52.
- [46] G. Lusztig, *Intersection cohomology complexes on a reductive group*, Invent. Math. **75** (1984), no. 2, 205–272, DOI 10.1007/BF01388564. MR732546
- [47] G. W. Mackey, *Unitary representations of group extensions*, Acta Math. **99** (1958), 265–311.
- [48] W. M. McGovern, *Cells of Harish-Chandra modules for real classical groups*, Amer. J. of Math. **120** (1998), 211–228.
- [49] C. Mœglin, *Front d’onde des représentations des groupes classiques p -adiques*, Amer. J. Math. **118** (1996), 1313–1346.
- [50] ———, *Paquets d’Arthur Spéciaux Unipotents aux Places Archimédiennes et Correspondance de Howe*, J. Cogdell et al. (eds.), Representation Theory, Number Theory, and Invariant Theory, In Honor of Roger Howe. Progress in Math. **323** (2017), 469–502.
- [51] C. Mœglin, M.-F. Vignéras, and J.-L. Waldspurger, *Correspondances de Howe sur un corps p -adique*, Lecture Notes in Mathematics, vol. 1291, Springer, 1987.
- [52] K. Nishiyama, H. Ochiai, K. Taniguchi, H. Yamashita, and S. Kato, *Nilpotent orbits, associated cycles and Whittaker models for highest weight representations*, Astérisque **273** (2001), 1–163.
- [53] K. Nishiyama, H. Ochiai, and C.-B. Zhu, *Theta lifting of nilpotent orbits for symmetric pairs*, Trans. Amer. Math. Soc. **358** (2006), 2713–2734.
- [54] K. Nishiyama and C.-B. Zhu, *Theta lifting of unitary lowest weight modules and their associated cycles*, Duke Math. J. **125** (2004), no. 03, 415–465.
- [55] T. Ohta, *The closures of nilpotent orbits in the classical symmetric pairs and their singularities*, Tohoku Math. J. **43** (1991), no. 2, 161–211.
- [56] ———, *Induction of nilpotent orbits for real reductive groups and associated varieties of standard representations*, Hiroshima Math. J. **29** (1999), no. 2, 347–360.
- [57] ———, *Nilpotent orbits of \mathbb{Z}_4 -graded Lie algebra and geometry of moment maps associated to the dual pair $(U(p, q), U(r, s))$* , Publ. RIMS **41** (2005), no. 3, 723–756.
- [58] A. Paul and P. Trapa, *Some small unipotent representations of indefinite orthogonal groups and the theta correspondence*, University of Aarhus Publ. Series **48** (2007), 103–125.
- [59] V. L. Popov and E. B. Vinberg, *Invariant Theory*, Algebraic Geometry IV: Linear Algebraic Groups, Invariant Theory, Encyclopedia of Mathematical Sciences, vol. 55, Springer, 1994.
- [60] T. Przebinda, *The duality correspondence of infinitesimal characters*, Colloq. Math. **70** (1996), 93–102.
- [61] ———, *Characters, dual pairs, and unitary representations*, Duke Math. J. **69** (1993), no. 3, 547–592.
- [62] S. Rallis, *On the Howe duality conjecture*, Compositio Math. **51** (1984), 333–399.
- [63] S. Sahi, *Explicit Hilbert spaces for certain unipotent representations*, Invent. Math. **110** (1992), no. 2, 409–418.
- [64] J. Sekiguchi, *Remarks on real nilpotent orbits of a symmetric pair*, J. Math. Soc. Japan **39** (1987), no. 1, 127–138.
- [65] W. Schmid and K. Vilonen, *Characteristic cycles and wave front cycles of representations of reductive Lie groups*, Annals of Math. **151** (2000), no. 3, 1071–1118.
- [66] E. Sommers, *Lusztig’s canonical quotient and generalized duality*, J. Algebra **243** (2001), no. 2, 790–812.
- [67] T. A. Springer and R. Steinberg, *Seminar on algebraic groups and related finite groups; Conjugate classes*, Lecture Notes in Math., vol. 131, Springer, 1970.
- [68] B. Sun and C.-B. Zhu, *A general form of Gelfand-Kazhdan criterion*, Manuscripta Math. **136** (2011), 185–197.
- [69] P. Trapa, *Special unipotent representations and the Howe correspondence*, University of Aarhus Publication Series **47** (2004), 210–230.
- [70] D. A. Vogan, *Irreducible characters of semisimple Lie groups. IV. Character-multiplicity duality*, Duke Math. J. **49** (1982), no. 4, 943–1073. MR683010
- [71] ———, *Unitary representations of reductive Lie groups*, Ann. of Math. Stud., vol. 118, Princeton University Press, 1987.
- [72] ———, *Associated varieties and unipotent representations*, Harmonic analysis on reductive groups, Proc. Conf., Brunswick/ME (USA) 1989, Prog. Math. **101** (1991), 315–388.
- [73] ———, *The method of coadjoint orbits for real reductive groups*, Representation theory of Lie groups (Park City, UT, 1998). IAS/Park City Math. Ser. **8** (2000), 179–238.
- [74] ———, *Unitary representations of reductive Lie groups*, Mathematics towards the Third Millennium (Rome, 1999). Accademia Nazionale dei Lincei, (2000), 147–167.

- [75] N. R. Wallach, *Real reductive groups I*, Academic Press Inc., 1988.
- [76] ———, *Real reductive groups II*, Academic Press Inc., 1992.
- [77] H. Weyl, *The classical groups: their invariants and representations*, Princeton University Press, 1947.
- [78] S. Yamana, *Degenerate principal series representations for quaternionic unitary groups*, Israel J. Math. **185** (2011), 77–124.

THE DEPARTMENT OF MATHEMATICS, 310 MALOTT HALL, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853

Email address: dmb14@cornell.edu

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, 800 DONGCHUAN ROAD, SHANGHAI, 200240, CHINA

Email address: hoxide@sjtu.edu.cn

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, CHINA

Email address: sun@math.ac.cn

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076

Email address: matzhucb@nus.edu.sg