

SPECIAL UNIPOTENT REPRESENTATIONS: COUNTING

DAN M. BARBASCH, JIA-JUN MA, BINYONG SUN, AND CHEN-BO ZHU

CONTENTS

1.		1
2.	Special non-special switch	1
3.	Counting the multiplicity of non-special representations	4
	References	6

1.

2. SPECIAL NON-SPECIAL SWITCH

2.0.1. *The case of $\star = \tilde{C}$.* Suppose that $\star = \tilde{C}$. We define an involution \natural on the set $\{\bullet, s, r, c, d\}$ by sending \bullet, s, r, c, d to \bullet, r, s, d, c respectively.

Let $\wp \subset \text{PP}_\star(\check{\mathcal{O}})$ such that $\wp \neq \emptyset$. Pick a pair $(2k-1, 2k) \in \wp$, and let $\wp' = \wp - \{(2k-1, 2k)\}$.

Lemma 2.1. *For each $\tau' \in \text{PBP}_\star(\check{\mathcal{O}}, \wp')$, there is a unique $\tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp)$ such that*

$$(2.1) \quad \begin{aligned} \mathcal{P}_\tau(i, j) &:= \begin{cases} \natural(\mathcal{Q}_{\tau'}(i, j)) & \text{if } i = k \\ \mathcal{P}_{\tau'}(i, j) & \text{otherwise} \end{cases} \\ \mathcal{Q}_\tau(i, j) &:= \begin{cases} \natural(\mathcal{P}_{\tau'}(i, j)) & \text{if } i = k \\ \mathcal{Q}_{\tau'}(i, j) & \text{otherwise} \end{cases} \end{aligned}$$

Moreover, the map $\tau' \mapsto \tau$ yields a bijection

$$(2.2) \quad \text{PBP}_\star(\check{\mathcal{O}}, \wp') \longrightarrow \text{PBP}_\star(\check{\mathcal{O}}, \wp).$$

Proof. One first verify that $(\iota(\check{\mathcal{O}}, \wp), \mathcal{P}_\tau) \times (j(\check{\mathcal{O}}, \wp), \mathcal{Q}_\tau) \times \tilde{C}$ is a valid painted bi-partition. We can define the inverse of (2.2) by (2.1) (switching the role of τ and τ'). This proves the bijectivity of (2.2). \square

[We now verify τ is a valid painted bipartition.

The relevant portion of \mathcal{P} and \mathcal{Q} are boxes with indexes (i, j) such that $k-1 \leq i \leq k$ and $\frac{\mathbf{r}_{2k}(\check{\mathcal{O}})}{2} \leq j \leq \frac{\mathbf{r}_{2k-1}(\check{\mathcal{O}})}{2}$

$$(2.3) \quad \begin{array}{ccccc} & w_1 & x_2 & & w_3 & x_0 & & & w_5 & y_0 & & w_7 & y_2 \\ & * & & & * & r & & & * & s & & * & \\ & * & & & * & \vdots & & & * & \vdots & & * & \\ & * & & \times & * & \vdots & & & * & \vdots & \times & * & \\ & * & & & * & \vdots & & & * & \vdots & & * & \\ & * & & & * & r & & & * & s & & * & \\ & w_2 & & & w_4 & x_1 & & & w_6 & y_1 & & w_8 & \end{array}$$

2000 *Mathematics Subject Classification.* 22E45, 22E46.

Key words and phrases. orbit method, unitary dual, unipotent representation, classical group, theta lifting, moment map.

Either all w_i are \emptyset (\star are all \emptyset) or all none-empty (\star are all \bullet). When w_i are all \emptyset or \bullet , the validity of τ is easy.

For the rest cases, w_1, w_3, w_5, w_7 must be painted by \bullet . Therefore, it suffice to consider the switch

$$w_2 \times w_4 x_1 \longleftrightarrow w_6 y_1 \times w_8 \quad .$$

	τ	τ^s	τ^{ns}	
	$\begin{array}{ c c c } \hline x'_0 & \star & \star \\ \hline x'_1 & x'_2 & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline x_1 & x_2 & x_3 & \\ \hline & & s & \\ & & \vdots & \\ & & s & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline y_0 & \star & \star & \star \\ \hline r & y_2 & & \\ \hline & & \vdots & \\ & & r & \\ & & y_1 & \\ & & y_3 & \\ \hline \end{array}$ \times	
$\begin{array}{ c c c } \hline x'_1 = & s & \\ \hline \end{array}$ then $\begin{array}{ c c c } \hline z_0 = & \emptyset / & s \\ \hline x_0 = & \emptyset / & \bullet \\ \hline x_1 = & \bullet & \\ \hline x_2 = & x'_2 & \\ \hline \end{array}$	$\begin{array}{ c c c } \hline x'_0 & \star & \star \\ \hline s & x_2 & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline \bullet & x_2 & \bullet & \\ \hline & & s & \\ & & \vdots & \\ & & s & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline r & x_2 & & \\ \hline & & \vdots & \\ & & r & \\ & & c & \\ & & d & \\ \hline \end{array}$ \times	$x_2 \neq r$
		$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline \bullet & r & \bullet & \\ \hline & & s & \\ & & \vdots & \\ & & s & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline r & c & & \\ \hline & & \vdots & \\ & & r & \\ & & r & \\ & & d & \\ \hline \end{array}$ \times	$x_2 = r$
$\begin{array}{ c c c } \hline x'_1 \neq & s & \\ \hline \end{array}$ then $\begin{array}{ c c c } \hline x_1 = & x'_1 & \\ \hline x_2 = & x'_2 & \\ \hline \end{array}$	$\begin{array}{ c c c } \hline x'_0 & \star & \star \\ \hline x'_1 & x_2 & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline x_1 & x_2 & s & \\ \hline & & s & \\ & & \vdots & \\ & & s & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline x_0 & \star & \star & \star \\ \hline r & x_2 & & \\ \hline & & \vdots & \\ & & r & \\ & & r & \\ & & x_1 & \\ \hline \end{array}$ \times	$\begin{array}{ c c c } \hline x'_0 \neq & c & \\ \hline \end{array}$ then $\begin{array}{ c c c c } \hline x'_0 = & \emptyset / s / & r & \\ \hline x_0 = & \emptyset / \bullet / & r & \\ \hline \end{array}$
		$\begin{array}{ c c c c } \hline c & \star & \star & \star \\ \hline d & x_2 & s & \\ \hline & & s & \\ & & \vdots & \\ & & s & \\ \hline \end{array}$ \times	$\begin{array}{ c c c c } \hline r & \star & \star & \star \\ \hline r & x_2 & & \\ \hline & & \vdots & \\ & & r & \\ & & c & \\ & & d & \\ \hline \end{array}$ \times	$\begin{array}{ c c c } \hline x'_0 = & c & \\ \hline \end{array}$ then $\begin{array}{ c c c c } \hline x_0 = & x'_0 = & c & \\ \hline x_1 = & x'_1 = & d & \\ \hline x_2 = & x'_2 = & \emptyset / d & \\ \hline \end{array}$

TABLE 1. “special-non-special” switch

We can list all the cases:

τ'				τ			
s	\times	r	d	s	c	\times	r
s	\times	d	d	s	c	\times	d
c	\times	r	d	c	c	\times	r
c	\times	d	d	c	c	\times	d

]

Lemma 2.2. Suppose $\star \in \{B, D\}$ and $\wp \in \text{PP}_\star(\check{\mathcal{O}})$. Let $\check{\mathcal{O}}_1$ be the \star' -good orbit having the rows

$$\mathbf{r}_1(\check{\mathcal{O}}_1) := \mathbf{r}_1(\check{\mathcal{O}}) + 2, \quad \text{and} \quad \mathbf{r}_{i+1}(\check{\mathcal{O}}_1) := \mathbf{r}_i(\check{\mathcal{O}}) \forall i = 1, 2, \dots$$

Let

$$\wp_1 = \{ (i+1, i+2) \mid (i, i+1) \in \wp \}.$$

Then the naive descent map yields a bijection

$$\text{PBP}_\star(\check{\mathcal{O}}_1, \wp_1) \xrightarrow{\nabla_{\text{naive}}} \text{PBP}_\star(\check{\mathcal{O}}, \wp)$$

Proof. Clear by the definition of ∇_{naive} . \square

Proposition 2.3. Suppose $\star \in \{B, C, \tilde{C}, D\}$. For $\wp \subset \text{PP}_\star(\check{\mathcal{O}})$, $|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| = |\text{PBP}_\star(\check{\mathcal{O}}, \emptyset)|$

Proof. Suppose $\star \in \{C, \tilde{C}\}$. The proposition following by applying Lemma 2.1 finitely many times to reduce \wp to \emptyset .

Suppose $\star \in \{B, D\}$. By Lemma 2.2, the problem translate to $\star' \in \{C, \tilde{C}\}$ which we already proved. \square

Then $\mathbf{r}_1(\check{\mathcal{O}}) > \mathbf{r}_2(\check{\mathcal{O}}) \geq 0$.

For every $\tau = (\imath, \mathcal{P}) \times (j, \mathcal{Q}) \times \star \in \text{PBP}_\star(\check{\mathcal{O}})$, its leg is defined to be the pair

$$\text{Leg}(\tau) := \Lambda_{\max(\imath'-1, 0), 1}(\imath, \mathcal{P}) \times \Lambda_{\max(j'-1, 0), 1}(j, \mathcal{Q}),$$

and its body is defined to be the pair

$$\text{Body}(\tau) := \bar{\Lambda}_{\max(\imath'-1, 0), 1}(\imath, \mathcal{P}) \times \bar{\Lambda}_{\max(j'-1, 0), 1}(j, \mathcal{Q}).$$

Note that if $\tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp)$, then $\text{Leg}(\tau)$ is represented by the first pair in (2.4) where

$$x_1 \neq \emptyset, \quad x_0 = \emptyset \Leftrightarrow x_2 = \emptyset \Leftrightarrow \mathbf{r}_2(\check{\mathcal{O}}) = 0,$$

and the grey part consisting of $l - l' - 1$ boxes with label r .

$$(2.4) \quad \text{Leg}(\tau) : \begin{array}{c} \boxed{x_2} \\ \times \\ \begin{array}{c} \boxed{x_0} \\ r \\ \vdots \\ r \\ \boxed{x_1} \end{array} \end{array} \quad \text{Leg}(\bar{\tau}) : \begin{array}{c} \begin{array}{c} \boxed{y_0} \\ s \\ \vdots \\ s \\ \boxed{y_1} \end{array} \\ \times \\ \boxed{y_2} \end{array},$$

Likewise, for every $\bar{\tau} \in \text{PBP}_\star(\check{\mathcal{O}}, \bar{\wp})$, $\text{Leg}(\bar{\tau})$ is represented by the second pair of (2.4) where

$$y_1 \neq \emptyset, \quad y_0 = \emptyset \Leftrightarrow y_2 = \emptyset \Leftrightarrow \mathbf{r}_2(\check{\mathcal{O}}) = 0,$$

and the grey part consisting of $l - l' - 1$ boxes with label s .

The following proposition is much easier to check than Proposition ???. We omit the details.

Proposition 2.4. *Suppose that $\star = \tilde{C}$ and $(1, 2) \in \wp$. For every $\tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp)$ such that $\text{Leg}(\tau)$ is represented by the first pair in (2.4), there is a unique element $\bar{\tau} \in \text{PBP}_\star(\check{\mathcal{O}}, \bar{\wp})$ such that*

$$\text{Body}(\bar{\tau}) = \text{Body}(\tau)$$

and $\text{Leg}(\bar{\tau})$ is represented by the second pair in (2.4) with

$$y_i = \bullet, r, s, d, c, \text{ or } \emptyset \quad (i = 0, 1, 2),$$

respectively if

$$x_i = \bullet, s, r, c, d, \text{ or } \emptyset.$$

Moreover, the map

$$\text{PBP}_\star(\check{\mathcal{O}}, \wp) \rightarrow \text{PBP}_\star(\check{\mathcal{O}}, \bar{\wp}), \quad \tau \mapsto \bar{\tau}.$$

is bijective.

3. COUNTING THE MULTIPLICITY OF NON-SPECIAL REPRESENTATIONS

In this section, we assume $\star \in \{B, C, \tilde{C}, D\}$. Let $\check{\mathcal{O}}$ be a \star -good parity orbit.

For $\star \in \{B, D\}$, we write

$$\text{PBP}_\star^d(\check{\mathcal{O}}, \wp) := \{ \tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp) \mid x_\tau = d \}.$$

$$\text{PBP}_\star^{rc}(\check{\mathcal{O}}, \wp) := \{ \tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp) \mid x_\tau \in \{r, c\} \}.$$

$$\text{PBP}_\star^s(\check{\mathcal{O}}, \wp) := \{ \tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp) \mid x_\tau = s \}.$$

$$\text{PBP}_\star^{-s}(\check{\mathcal{O}}, \wp) := \{ \tau \in \text{PBP}_\star(\check{\mathcal{O}}, \wp) \mid x_\tau \neq s \}.$$

Note that $\text{PBP}_\star^{-s}(\check{\mathcal{O}}, \wp) = \text{PBP}_\star^d(\check{\mathcal{O}}, \wp) \sqcup \text{PBP}_\star^{rc}(\check{\mathcal{O}}, \wp)$. We write

$$\text{PBP}_{D,sc}(\check{\mathcal{O}}_0) := \{ \tau_0 \in \text{PBP}_D(\check{\mathcal{O}}_0) \mid \mathcal{P}_{\tau_0}^{-1}(\{s, c\}) \neq \emptyset \}.$$

We define $\text{PBP}_{D,sc}^\sharp$ similarly.

We will prove the following counting lemma:

Proposition 3.1. *Let $\wp \in \text{PP}(\check{\mathcal{O}})$, we have $|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| = |\text{PBP}_\star(\check{\mathcal{O}}, \emptyset)|$. When $\star \in \{B, D\}$, for $\sharp \in \{-s, s, d, rc\}$, we have*

$$(3.1) \quad |\text{PBP}_\star^\sharp(\check{\mathcal{O}}, \wp)| = |\text{PBP}_\star^\sharp(\check{\mathcal{O}}, \emptyset)|$$

Proof. We prove the proposition by induction. Assume for each good parity orbit $\check{\mathcal{O}}'$ such that $\mathbf{r}_k(\check{\mathcal{O}}') = 0$ the proposition holds.

Suppose $\star \in \{C, \tilde{C}\}$.

When $(1, 2) \in \text{PP}_\star(\check{\mathcal{O}})$, and $(1, 2) \in \wp$. Let $\bar{\wp} = \wp - \{(1, 2)\}$. By the switching algorithm, we have

$$|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| = |\text{PBP}_\star(\check{\mathcal{O}}, \bar{\wp})|.$$

Therefore, we can assume $(1, 2) \notin \wp$ without loss of generality.

Suppose $(1, 2) \in \text{PP}_\star(\check{\mathcal{O}})$, we have a bijection

$$\text{PBP}_\star(\check{\mathcal{O}}, \wp) \xrightarrow{\nabla} \text{PBP}_{\star'}(\check{\mathcal{O}}', \wp')$$

for each $(1, 2) \notin \wp \subset \text{PP}_\star(\check{\mathcal{O}})$. By the induction hypothesis,

$$|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| = |\text{PBP}_{\star'}(\check{\mathcal{O}}', \wp')| = |\text{PBP}_{\star'}(\check{\mathcal{O}}', \emptyset)| = |\text{PBP}_\star(\check{\mathcal{O}}, \emptyset)|.$$

Suppose $(1, 2) \notin \text{PP}_\star(\check{\mathcal{O}})$, we have a bijection

$$\text{PBP}_\star(\check{\mathcal{O}}, \wp) \xrightarrow{\nabla} \text{PBP}_{\star'}^{-s}(\check{\mathcal{O}}', \wp')$$

for each $\wp \subset \text{PP}_\star(\check{\mathcal{O}})$. Therefore, by induction hypothesis, we have

$$|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| = |\text{PBP}_{\star'}^{-s}(\check{\mathcal{O}}', \wp')| = |\text{PBP}_{\star'}^{-s}(\check{\mathcal{O}}', \emptyset)| = |\text{PBP}_\star(\check{\mathcal{O}}, \emptyset)|.$$

Note that under the

$$\text{PBP}_{\star'}(\check{\mathcal{O}}', \wp') \longrightarrow \text{PBP}_\star(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_D(\check{\mathcal{O}}')$$

Suppose $\star \in \{B, D\}$. Assume $(2, 3) \in \text{PBP}_\star(\check{\mathcal{O}})$. Then

$$\delta: \text{PBP}_\star(\check{\mathcal{O}}, \wp) \longrightarrow \text{PBP}_\star(\check{\mathcal{O}}', \wp') \times \text{PBP}_D(\check{\mathcal{O}}_0)$$

is a bijection for each $\wp \in \text{PP}_\star(\check{\mathcal{O}})$. Therefore

$$|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| = |\text{PBP}_{\star'}(\check{\mathcal{O}}', \wp')| |\text{PBP}_D(\check{\mathcal{O}}_0)| = |\text{PBP}_{\star'}(\check{\mathcal{O}}', \emptyset)| |\text{PBP}_D(\check{\mathcal{O}}_0)| = |\text{PBP}_\star(\check{\mathcal{O}}, \emptyset)|.$$

Since $x_\tau = d \Leftrightarrow x_{\tau_0} = d$ and $x_\tau = s \Leftrightarrow x_{\tau_0} = s$, δ induces bijections

$$\text{PBP}_\star^\sharp(\check{\mathcal{O}}, \wp) \longrightarrow \text{PBP}_\star(\check{\mathcal{O}}', \wp') \times \text{PBP}_D^\sharp(\check{\mathcal{O}}_0)$$

for $\sharp \in \{s, -s, d, rc\}$. Now (3.1) follows.

Assume $(2, 3) \notin \text{PBP}_\star(\check{\mathcal{O}})$. Note that the image of δ as the following

$$\begin{aligned} \text{Im}(\delta) &= \{(\tau'', \tau_0) \in \text{PBP}_\star(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_D(\check{\mathcal{O}}_{\mathbf{t}}, \emptyset) \mid x_{\tau''} = d, \text{ or } \mathcal{P}_{\tau_0}^{-1}(\{s, c\}) \neq \emptyset\} \\ &= \text{PBP}_\star^d(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_D(\check{\mathcal{O}}_0) \sqcup \text{PBP}_\star^{rc}(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_{D,sc}(\check{\mathcal{O}}_0). \end{aligned}$$

Now

$$\begin{aligned} &|\text{PBP}_\star(\check{\mathcal{O}}, \wp)| \\ &= |\text{PBP}_\star^d(\check{\mathcal{O}}, \wp'')| |\text{PBP}_d(\check{\mathcal{O}}_0)| + |\text{PBP}_\star^{rc}(\check{\mathcal{O}}, \wp'')| |\text{PBP}_{\star}^{sc}(\check{\mathcal{O}}_0)| \\ &= |\text{PBP}_\star(\check{\mathcal{O}}, \wp)| \end{aligned}$$

The equation (3.1) follows from the bijection:

$$\text{PBP}_\star^\sharp(\check{\mathcal{O}}, \wp) \xrightarrow{\delta} \text{PBP}_\star^d(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_D^\sharp(\check{\mathcal{O}}_0) \sqcup \text{PBP}_\star^{rc}(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_{D,sc}^\sharp(\check{\mathcal{O}}_0).$$

This finished the proof of the proposition. \square

In the rest of this section we assume that $\check{\mathcal{O}} \neq \emptyset$. Let $\check{\mathcal{O}}'$ be the dual descent of $\check{\mathcal{O}}$ as defined in the Introduction. Then $\check{\mathcal{O}}'$ has \star' -good parity, where \star' is the Howe dual of \star . Put

$$l' := l_{\star', \check{\mathcal{O}}'}.$$

The following equation of signatures will be crucial in our computation of the local system in the next section.

Proposition 3.2. *Suppose $\mathbf{r}_2(\check{\mathcal{O}}) > 0$. Let $\check{\mathcal{O}}'' := \check{\nabla}^2(\check{\mathcal{O}})$ and $\wp'' = \check{\nabla}^2(\wp)$. Consider the map*

$$(3.2) \quad \delta: \text{PBP}_\star(\check{\mathcal{O}}) \longrightarrow \text{PBP}_\star(\check{\mathcal{O}}'', \wp'') \times \text{PBP}_{\star_{\mathbf{t}}}(\check{\mathcal{O}}_{\mathbf{t}}, \emptyset), \quad \tau \mapsto (\nabla^2(\tau), \tau_{\mathbf{t}})$$

Then δ is an injection.

- When $\star = C^*$ or $\mathbf{r}_2(\check{\mathcal{O}}) > \mathbf{r}_3(\check{\mathcal{O}})$, the map δ is a bijection. Moreover,

$$\text{Sign}(\tau) = (\mathbf{c}_2(\mathcal{O}), \mathbf{c}_2(\mathcal{O})) + \text{Sign}(\nabla^2(\tau)) + \text{Sign}(\tau_{\mathbf{t}}).$$

- When $\star \in \{B, D\}$ and $\mathbf{r}_2(\check{\mathcal{O}}) = \mathbf{r}_3(\check{\mathcal{O}})$, the map δ is an injection, whose image equals

$$\{(\tau'', \tau_0) \in \text{PBP}_\star(\check{\mathcal{O}}, \wp'') \times \text{PBP}_D(\check{\mathcal{O}}_{\mathbf{t}}, \emptyset) \mid x_{\tau''} = d, \text{ or } \mathcal{P}_{\tau_0}^{-1}(\{s, c\}) \neq \emptyset\}.$$

Moreover,

$$\text{Sign}(\tau) = (\mathbf{c}_2(\mathcal{O}) - 1, \mathbf{c}_2(\mathcal{O}) - 1) + \text{Sign}(\nabla^2(\tau)) + \text{Sign}(\tau_{\mathbf{t}}).$$

REFERENCES

- [1] J. Adams, *Discrete spectrum of the reductive dual pair $(O(p, q), Sp(2m))$* , Invent. Math. **74** (1983), no. 3, 449–475.
- [2] J. Adams, B. Barbasch, and D. A. Vogan, *The Langlands classification and irreducible characters for real reductive groups*, Progress in Math., vol. 104, Birkhauser, 1991.
- [3] Jeffrey Adams and Fokko du Cloux, *Algorithms for representation theory of real reductive groups*, Journal of the Institute of Mathematics of Jussieu **8** (2009), no. 2, 209–259, DOI 10.1017/S1474748008000352.
- [4] J. Arthur, *On some problems suggested by the trace formula*, Lie group representations, II (College Park, Md.), Lecture Notes in Math. 1041 (1984), 1–49.
- [5] ———, *Unipotent automorphic representations: conjectures*, Orbites unipotentes et représentations, II, Astérisque **171-172** (1989), 13–71.
- [6] L. Auslander and B. Kostant, *Polarizations and unitary representations of solvable Lie groups*, Invent. Math. **14** (1971), 255–354.
- [7] D. Barbasch, *The unitary dual for complex classical Lie groups*, Invent. Math. **96** (1989), no. 1, 103–176.
- [8] ———, *Unipotent representations for real reductive groups*, Proceedings of ICM (1990), Kyoto (2000), 769–777.
- [9] Dan Barbasch and David Vogan, *Weyl Group Representations and Nilpotent Orbits*, Representation Theory of Reductive Groups: Proceedings of the University of Utah Conference 1982, 1983, pp. 21–33.
- [10] D. Barbasch, *Orbital integrals of nilpotent orbits*, The mathematical legacy of Harish-Chandra, Proc. Sympos. Pure Math. **68** (2000), 97–110.
- [11] ———, *The unitary spherical spectrum for split classical groups*, J. Inst. Math. Jussieu **9** (2010), 265–356.
- [12] ———, *Unipotent representations and the dual pair correspondence*, J. Cogdell et al. (eds.), Representation Theory, Number Theory, and Invariant Theory, In Honor of Roger Howe. Progress in Math. **323** (2017), 47–85.
- [13] D. Barbasch, J.-J. Ma, B.-Y. Sun, and C.-B. Zhu, *The counting the special unipotent representations*, in preparation.
- [14] D. Barbasch and D. A. Vogan, *Weyl group representations and nilpotent orbits*, in Representation theory of reductive groups (Park City, Utah, 1982), Progress in Math. **40** (1983), 21–33.
- [15] ———, *Unipotent representations of complex semisimple groups*, Annals of Math. **121** (1985), no. 1, 41–110.
- [16] R. Brylinski, *Dixmier algebras for classical complex nilpotent orbits via Kraft-Procesi models. I*, The orbit method in geometry and physics (Marseille, 2000). Progress in Math. **213** (2003), 49–67.
- [17] W. Borho, *Recent advances in enveloping algebras of semisimple Lie-algebras*, Séminaire Bourbaki, Exp. No. 489 (1976/77), 1–18.
- [18] Roger W. Carter, *Finite groups of Lie type*, Wiley Classics Library, John Wiley & Sons, Ltd., Chichester, 1993.
- [19] W. Casselman, *Canonical extensions of Harish-Chandra modules to representations of G* , Canad. J. Math. **41** (1989), 385–438.
- [20] F. Du Cloux, *Sur les représentations différentiables des groupes de Lie algébriques*, Ann. Sci. École Norm. Sup. **24** (1991), no. 3, 257–318.
- [21] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebra: an introduction*, Van Nostrand Reinhold Co., 1993.
- [22] A. Daszkiewicz, W. Kraśkiewicz, and T. Przebinda, *Nilpotent orbits and complex dual pairs*, J. Algebra **190** (1997), no. 2, 518 – 539.
- [23] ———, *Dual pairs and Kostant-Sekiguchi correspondence. II. Classification of nilpotent elements*, Central European J. Math. **3** (2005), 430–474.
- [24] J. Dixmier and P. Malliavin, *Factorisations de fonctions et de vecteurs indéfiniment différentiables*, Bull. Sci. Math. (2) **102** (1978), 307–330.
- [25] M. Duflo, *Théorie de Mackey pour les groupes de Lie algébriques*, Acta Math. **149** (1982), no. 3-4, 153–213.
- [26] R. Gomez and C.-B. Zhu, *Local theta lifting of generalized Whittaker models associated to nilpotent orbits*, Geom. Funct. Anal. **24** (2014), no. 3, 796–853.
- [27] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique IV: Étude locale des schémas et des morphismes de schémas. II*, Inst. Hautes Études Sci. Publ. Math. **24** (1965).

- [28] ———, *Éléments de géométrie algébrique IV: Étude locale des schémas et des morphismes de schémas. III*, Inst. Hautes Études Sci. Publ. Math. **28** (1966).
- [29] M. Harris, J.-S. Li, and B. Sun, *Theta correspondences for close unitary groups*, Arithmetic Geometry and Automorphic Forms, Adv. Lect. Math. (ALM) **19** (2011), 265–307.
- [30] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics, 52. New York-Heidelberg-Berlin: Springer-Verlag, 1983.
- [31] H. He, *Unipotent representations and quantum induction*, arXiv:math/0210372 (2002).
- [32] ———, *Unitary representations and theta correspondence for type I classical groups*, J. Funct. Anal. **199** (2003), no. 1, 92–121.
- [33] J.-S. Huang and J.-S. Li, *Unipotent representations attached to spherical nilpotent orbits*, Amer. J. Math. **121** (1999), no. 3, 497–517.
- [34] J.-S. Huang and C.-B. Zhu, *On certain small representations of indefinite orthogonal groups*, Represent. Theory **1** (1997), 190–206.
- [35] R. Howe, *θ -series and invariant theory*, Automorphic Forms, Representations and L -functions, Proc. Sympos. Pure Math, vol. 33, 1979, pp. 275–285.
- [36] ———, *On a notion of rank for unitary representations of the classical groups*, Harmonic analysis and group representations, Liguori, Naples (1982), 223–331.
- [37] ———, *Transcending classical invariant theory*, J. Amer. Math. Soc. **2** (1989), 535–552.
- [38] ———, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, Piatetski-Shapiro, I. et al. (eds.), The Schur lectures (1992). Ramat-Gan: Bar-Ilan University, Isr. Math. Conf. Proc. 8, (1995), 1–182.
- [39] D. Jiang, B. Liu, and G. Savin, *Raising nilpotent orbits in wave-front sets*, Represent. Theory **20** (2016), 419–450.
- [40] A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Uspehi Mat. Nauk **17** (1962), 57–110.
- [41] B. Kostant, *Quantization and unitary representations*, Lectures in Modern Analysis and Applications III, Lecture Notes in Math. **170** (1970), 87–208.
- [42] H. Kraft and C. Procesi, *On the geometry of conjugacy classes in classical groups*, Comment. Math. Helv. **57** (1982), 539–602.
- [43] S. S. Kudla and S. Rallis, *Degenerate principal series and invariant distributions*, Israel J. Math. **69** (1990), 25–45.
- [44] S. S. Kudla, *Some extensions of the Siegel-Weil formula*, In: Gan W., Kudla S., Tschinkel Y. (eds) Eisenstein Series and Applications. Progress in Mathematics, vol 258. Birkhäuser Boston (2008), 205–237.
- [45] S. T. Lee and C.-B. Zhu, *Degenerate principal series and local theta correspondence II*, Israel J. Math. **100** (1997), 29–59.
- [46] ———, *Degenerate principal series of metaplectic groups and Howe correspondence*, D. Prasad et al. (eds.), Automorphic Representations and L-Functions, Tata Institute of Fundamental Research, India, (2013), 379–408.
- [47] J.-S. Li, *Singular unitary representations of classical groups*, Invent. Math. **97** (1989), no. 2, 237–255.
- [48] Q. Liu, *Algebraic Geometry and Arithmetic Curves*, Oxford University Press, 2006.
- [49] H. Y. Loke and J. Ma, *Invariants and K -spectrums of local theta lifts*, Compositio Math. **151** (2015), 179–206.
- [50] G. Lusztig and N. Spaltenstein, *Induced unipotent classes*, j. London Math. Soc. **19** (1979), 41–52.
- [51] G. Lusztig, *Intersection cohomology complexes on a reductive group*, Invent. Math. **75** (1984), no. 2, 205–272, DOI 10.1007/BF01388564. MR732546
- [52] G. W. Mackey, *Unitary representations of group extensions*, Acta Math. **99** (1958), 265–311.
- [53] W. M. McGovern, *Cells of Harish-Chandra modules for real classical groups*, Amer. J. of Math. **120** (1998), 211–228.
- [54] C. Mœglin, *Front d’onde des représentations des groupes classiques p -adiques*, Amer. J. Math. **118** (1996), 1313–1346.
- [55] ———, *Paquets d’Arthur Spéciaux Unipotents aux Places Archimédiennes et Correspondance de Howe*, J. Cogdell et al. (eds.), Representation Theory, Number Theory, and Invariant Theory, In Honor of Roger Howe. Progress in Math. **323** (2017), 469–502.
- [56] C. Mœglin and D. Renard, *Sur les paquets d’Arthur des groupes classiques réels*, J. Eur. Math. Soc. **22** (2020), 1827–1892.
- [57] C. Mœglin, M.-F. Vignéras, and J.-L. Waldspurger, *Correspondances de Howe sur un corps p -adique*, Lecture Notes in Mathematics, vol. 1291, Springer, 1987.

- [58] K. Nishiyama, H. Ochiai, K. Taniguchi, H. Yamashita, and S. Kato, *Nilpotent orbits, associated cycles and Whittaker models for highest weight representations*, *Astérisque* **273** (2001), 1–163.
- [59] K. Nishiyama, H. Ochiai, and C.-B. Zhu, *Theta lifting of nilpotent orbits for symmetric pairs*, *Trans. Amer. Math. Soc.* **358** (2006), 2713–2734.
- [60] K. Nishiyama and C.-B. Zhu, *Theta lifting of unitary lowest weight modules and their associated cycles*, *Duke Math. J.* **125** (2004), 415–465.
- [61] T. Ohta, *The closures of nilpotent orbits in the classical symmetric pairs and their singularities*, *Tohoku Math. J.* **43** (1991), no. 2, 161–211.
- [62] ———, *Induction of nilpotent orbits for real reductive groups and associated varieties of standard representations*, *Hiroshima Math. J.* **29** (1999), no. 2, 347–360.
- [63] ———, *Nilpotent orbits of \mathbb{Z}_4 -graded Lie algebra and geometry of moment maps associated to the dual pair $(U(p, q), U(r, s))$* , *Publ. RIMS* **41** (2005), no. 3, 723–756.
- [64] A. Paul and P. Trapa, *Some small unipotent representations of indefinite orthogonal groups and the theta correspondence*, *University of Aarhus Publ. Series* **48** (2007), 103–125.
- [65] V. L. Popov and E. B. Vinberg, *Invariant Theory*, *Algebraic Geometry IV: Linear Algebraic Groups*, *Invariant Theory*, *Encyclopedia of Mathematical Sciences*, vol. 55, Springer, 1994.
- [66] T. Przebinda, *The duality correspondence of infinitesimal characters*, *Colloq. Math.* **70** (1996), 93–102.
- [67] ———, *Characters, dual pairs, and unitary representations*, *Duke Math. J.* **69** (1993), no. 3, 547–592.
- [68] S. Rallis, *On the Howe duality conjecture*, *Compositio Math.* **51** (1984), 333–399.
- [69] David A. Renard and Peter E. Trapa, *Irreducible genuine characters of the metaplectic group: Kazhdan-Lusztig algorithm and Vogan duality*, *Represent. Theory* **4** (2000), 245–295, DOI 10.1090/S1088-4165-00-00105-9.
- [70] S. Sahi, *Explicit Hilbert spaces for certain unipotent representations*, *Invent. Math.* **110** (1992), no. 2, 409–418.
- [71] J. Sekiguchi, *Remarks on real nilpotent orbits of a symmetric pair*, *J. Math. Soc. Japan* **39** (1987), no. 1, 127–138.
- [72] W. Schmid and K. Vilonen, *Characteristic cycles and wave front cycles of representations of reductive Lie groups*, *Annals of Math.* **151** (2000), no. 3, 1071–1118.
- [73] E. Sommers, *Lusztig’s canonical quotient and generalized duality*, *J. Algebra* **243** (2001), no. 2, 790–812.
- [74] T. A. Springer and R. Steinberg, *Seminar on algebraic groups and related finite groups; Conjugate classes*, *Lecture Notes in Math.*, vol. 131, Springer, 1970.
- [75] B. Sun and C.-B. Zhu, *A general form of Gelfand-Kazhdan criterion*, *Manuscripta Math.* **136** (2011), 185–197.
- [76] P. Trapa, *Special unipotent representations and the Howe correspondence*, *University of Aarhus Publication Series* **47** (2004), 210–230.
- [77] D. A. Vogan, *Irreducible characters of semisimple Lie groups. IV. Character-multiplicity duality*, *Duke Math. J.* **49** (1982), no. 4, 943–1073. MR683010
- [78] ———, *Unitary representations of reductive Lie groups*, *Ann. of Math. Stud.*, vol. 118, Princeton University Press, 1987.
- [79] ———, *Associated varieties and unipotent representations*, *Harmonic analysis on reductive groups*, *Proc. Conf., Brunswick/ME (USA) 1989*, *Prog. Math.* **101** (1991), 315–388.
- [80] ———, *The method of coadjoint orbits for real reductive groups*, *Representation theory of Lie groups (Park City, UT, 1998)*, *IAS/Park City Math. Ser.* **8** (2000), 179–238.
- [81] ———, *Unitary representations of reductive Lie groups*, *Mathematics towards the Third Millennium (Rome, 1999)*, *Accademia Nazionale dei Lincei*, (2000), 147–167.
- [82] N. R. Wallach, *Real reductive groups I*, *Academic Press Inc.*, 1988.
- [83] ———, *Real reductive groups II*, *Academic Press Inc.*, 1992.
- [84] H. Weyl, *The classical groups: their invariants and representations*, *Princeton University Press*, 1947.
- [85] S. Yamana, *Degenerate principal series representations for quaternionic unitary groups*, *Israel J. Math.* **185** (2011), 77–124.

THE DEPARTMENT OF MATHEMATICS, 310 MALOTT HALL, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853

Email address: dmb14@cornell.edu

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, 800 DONGCHUAN ROAD, SHANGHAI, 200240, CHINA

Email address: hoxide@sjtu.edu.cn

INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU, 310058, CHINA

Email address: sunbinyong@zju.edu.cn

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076

Email address: matzhucb@nus.edu.sg