

The weak Arthur packets of real classical groups

Ma, Jia-Jun

(joint with Dan Barbasch, Binyong Sun and Chengbo Zhu)

School of Mathematical Sciences, Xiamen University
Department of Mathematics, Xiamen University Malaysia Campus

November 10, 2023

(BIRS-IASM, Arthur packets)

Jordan-Chevalley decomposition

- For each $g \in G := \mathrm{GL}_n(\mathbb{C})$, $\exists!$ pair (s, u) such that

$$g = su = us,$$

Jordan-Chevalley decomposition

- For each $g \in G := \mathrm{GL}_n(\mathbb{C})$, $\exists!$ pair (s, u) such that

$$g = su = us,$$

- s is semisimple, and u is unipotent ($u - 1$ is nilpotent.)

Jordan-Chevalley decomposition

- For each $g \in G := \mathrm{GL}_n(\mathbb{C})$, $\exists!$ pair (s, u) such that

$$g = su = us,$$

- s is semisimple, and u is unipotent ($u - 1$ is nilpotent.)

Camille Jordan



Claude Chevalley



Jordan-Chevalley decomposition

- For each $g \in G := \mathrm{GL}_n(\mathbb{C})$, $\exists!$ pair (s, u) such that


$$g = su = us,$$

- s is semisimple, and u is unipotent ($u - 1$ is nilpotent.)

Camille Jordan



Claude Chevalley



- Classification of conjugation classes in G :

$$G / \sim = \bigsqcup_{s \in G_{\text{s.s.}} / \sim} \{su \mid u \in G_s\} / \sim \xleftrightarrow{\text{bij.}} \bigsqcup_{s \in G_{\text{s.s.}} / \sim} \text{unip}(G_s)$$

Finite group of Lie type: $G := \mathbf{G}(\mathbb{F}_q)$

- Deligne-Lusztig and Lusztig (1970s–1980s)

Finite group of Lie type: $G := \mathbf{G}(\mathbb{F}_q)$

- Deligne-Lusztig and Lusztig (1970s–1980s)
- *Jordan decomposition*:

Finite group of Lie type: $G := \mathbf{G}(\mathbb{F}_q)$

- Deligne-Lusztig and Lusztig (1970s–1980s)

- *Jordan decomposition*:

Define dual group \check{G} . E.g. $\mathrm{Sp}_{2n} = G \longleftrightarrow \check{G} = \mathrm{SO}_{2n+1}$

Deligne



$$\mathrm{Irr}(G) = \bigsqcup_{s \in \check{G}_{\mathrm{s.s.}} / \sim} \mathcal{E}(G, s).$$

Lusztig



Finite group of Lie type: $G := \mathbf{G}(\mathbb{F}_q)$

- Deligne-Lusztig and Lusztig (1970s–1980s)
- *Jordan decomposition*:

Define dual group \check{G} . E.g. $\mathrm{Sp}_{2n} = G \longleftrightarrow \check{G} = \mathrm{SO}_{2n+1}$

Deligne



$$\mathrm{Irr}(G) = \bigsqcup_{s \in \check{G}_{\mathrm{s.s.}} / \sim} \mathcal{E}(G, s).$$

Lusztig



Lusztig's map to the unipotent packet.

$$\mathcal{E}(G, s) \xrightarrow[\mathcal{L}_s]{bij.} \mathcal{E}(\check{G}_s, 1)$$

Representations of Real Lie groups


- $G \in \{ \mathrm{GL}_n(\mathbb{R}), \mathrm{U}(p, q), \mathrm{O}(p, q), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \dots \}$

Representations of Real Lie groups

- $G \in \{ \mathrm{GL}_n(\mathbb{R}), \mathrm{U}(p, q), \mathrm{O}(p, q), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \dots \}$
 $\{ \text{discrete series} \} \subset \{ \text{tempered} \} \subset \{ \text{unitary} \} \subset \mathrm{Irr}(G)$


Representations of Real Lie groups

- $G \in \{ \mathrm{GL}_n(\mathbb{R}), \mathrm{U}(p, q), \mathrm{O}(p, q), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \dots \}$
 $\{ \text{discrete series} \} \subset \{ \text{tempered} \} \subset \{ \text{unitary} \} \subset \mathrm{Irr}(G)$
- Unitary dual of $\mathrm{SL}_2(\mathbb{R})$:




Representations of Real Lie groups

- $G \in \{ \mathrm{GL}_n(\mathbb{R}), \mathrm{U}(p, q), \mathrm{O}(p, q), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \dots \}$
 $\{ \text{discrete series} \} \subset \{ \text{tempered} \} \subset \{ \text{unitary} \} \subset \mathrm{Irr}(G)$
- Unitary dual of $\mathrm{SL}_2(\mathbb{R})$:



Representations of Real Lie groups

- $G \in \{ \mathrm{GL}_n(\mathbb{R}), \mathrm{U}(p, q), \mathrm{O}(p, q), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \dots \}$
 $\{ \text{discrete series} \} \subset \{ \text{tempered} \} \subset \{ \text{unitary} \} \subset \mathrm{Irr}(G)$
- Unitary dual of $\mathrm{SL}_2(\mathbb{R})$:



Elias M. Stein



- *Open problem:* Structure of the unitary dual!

- $W_{\mathbb{R}} = \mathbb{C}^{\times} \cup j\mathbb{C}^{\times}$ such that $jzj^{-1} = \bar{z}, j^2 = -1 \in \mathbb{C}^{\times}$.

Langlands and Arthur

- $W_{\mathbb{R}} = \mathbb{C}^{\times} \cup j\mathbb{C}^{\times}$ such that $jzj^{-1} = \bar{z}, j^2 = -1 \in \mathbb{C}^{\times}$.
- Langlands dual group ${}^L G = \check{G} \rtimes \langle j \rangle$.
Eg. $\mathrm{Sp}_{2n}(\mathbb{R}) = G \longleftrightarrow {}^L G = \mathrm{SO}_{2n+1}(\mathbb{C}) \times \langle j \rangle$

Langlands and Arthur

- $W_{\mathbb{R}} = \mathbb{C}^{\times} \cup j\mathbb{C}^{\times}$ such that $jzj^{-1} = \bar{z}, j^2 = -1 \in \mathbb{C}^{\times}$.
- Langlands dual group ${}^L G = \check{G} \rtimes \langle j \rangle$.
Eg. $\mathrm{Sp}_{2n}(\mathbb{R}) = G \longleftrightarrow {}^L G = \mathrm{SO}_{2n+1}(\mathbb{C}) \rtimes \langle j \rangle$
- *Langlands*



$$\mathrm{Irr}(G) \xrightarrow{\text{finite to one}} \{ \phi: W_{\mathbb{R}} \rightarrow {}^L G \} / \check{G}.$$

Langlands and Arthur

- $W_{\mathbb{R}} = \mathbb{C}^{\times} \cup j\mathbb{C}^{\times}$ such that $jzj^{-1} = \bar{z}, j^2 = -1 \in \mathbb{C}^{\times}$.
- Langlands dual group ${}^L G = \check{G} \rtimes \langle j \rangle$.
Eg. $\mathrm{Sp}_{2n}(\mathbb{R}) = G \longleftrightarrow {}^L G = \mathrm{SO}_{2n+1}(\mathbb{C}) \times \langle j \rangle$
- *Langlands*



$$\mathrm{Irr}(G) \xrightarrow{\text{finite to one}} \{ \phi: W_{\mathbb{R}} \rightarrow {}^L G \} / \check{G}.$$

- *Arthur* $\mathrm{Irr}_{\mathrm{temp}}(G) \subset \mathrm{Irr}_A(G) \subset \mathrm{Irr}_{\mathrm{unit}}(G)$



$$\mathrm{Irr}_A(G) \xrightarrow{\text{finite to finite}} \{ \psi: W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G \} / \check{G}.$$

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$
- Unipotent parameter: $\psi|_{\mathbb{C}^\times} = \text{trivial}$

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$
- Unipotent parameter: $\psi|_{\mathbb{C}^\times} = \text{trivial}$
 \Leftrightarrow nilpotent orbit of a real form of \check{G} .

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$
- Unipotent parameter: $\psi|_{\mathbb{C}^\times} = \text{trivial}$
 \Leftrightarrow nilpotent orbit of a real form of \check{G} .



- *Conjecture:* \exists unipotent Arthur packets

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$
- Unipotent parameter: $\psi|_{\mathbb{C}^\times} = \text{trivial}$
 \Leftrightarrow nilpotent orbit of a real form of \check{G} .



- *Conjecture:* \exists unipotent Arthur packets

“On some problems suggested by the trace formula” 1980's

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$
- Unipotent parameter: $\psi|_{\mathbb{C}^\times} = \text{trivial}$
 \Leftrightarrow nilpotent orbit of a real form of \check{G} .



■ *Conjecture:* \exists unipotent Arthur packets

“On some problems suggested by the trace formula” 1980's

Mœglin Renard



■ *Reduction to unipotent Arthur packet*

Arthur's unipotent representations

- $\psi : W_{\mathbb{R}} \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G$
- Unipotent parameter: $\psi|_{\mathbb{C}^\times} = \text{trivial}$
 \Leftrightarrow nilpotent orbit of a real form of \check{G} .



■ *Conjecture:* \exists unipotent Arthur packets

“On some problems suggested by the trace formula” 1980's

Mœglin Renard



■ *Reduction to unipotent Arthur packet*

“Sur Les paquets d'Arthur des groupes classiques réels”
(2020 JEMS)

Barbasch-Vogan's definition of special unip. repn.

Barbasch



Vogan



■ $\text{Unip}(G) := \text{repn. with smallest "size"}$

Barbasch-Vogan's definition of special unip. repn.

Barbasch



Vogan



$\text{Unip}(G) := \text{repn. with smallest "size"}$

- Wavefront cycle/Associated cycle:

$$AC : \text{Irr}(G) \longrightarrow \mathcal{K}(\text{equiv. sheaves on nil-cone})$$

Barbasch-Vogan's definition of special unip. repn.

Barbasch



Vogan




$\text{Unip}(G) := \text{repn. with smallest "size"}$

- Wavefront cycle/Associated cycle:

$AC : \text{Irr}(G) \longrightarrow \mathcal{K}(\text{equiv. sheaves on nil-cone})$

- E.g.: $\text{Lie}(\text{SL}_2(\mathbb{R})) \cong \mathbb{R}^3$.



Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- \check{O} : a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{O}}$.

Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- $\check{\mathcal{O}}$: a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{\mathcal{O}}}$.
 \rightsquigarrow the maximal primitive ideal $\mathcal{I}_{\check{\mathcal{O}}} \subset \mathcal{U}(\mathfrak{g})$.

Duflo



Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- \check{O} : a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{O}}$.

\rightsquigarrow the maximal primitive ideal $\mathcal{I}_{\check{O}} \subset \mathcal{U}(\mathfrak{g})$.

- *Definition* (Barbasch-Vogan):

Duflo



Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- \check{O} : a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{O}}$.

Duflo

\rightsquigarrow the maximal primitive ideal $\mathcal{I}_{\check{O}} \subset \mathcal{U}(\mathfrak{g})$.

- *Definition* (Barbasch-Vogan):

An irr. adm. G -module is called *special unipotent* if

$$\mathrm{Ann}_{\mathcal{U}(\mathfrak{g})}(\pi) = \mathcal{I}_{\check{O}}.$$



Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- $\check{\mathcal{O}}$: a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{\mathcal{O}}}$.

Duflo

\rightsquigarrow the maximal primitive ideal $\mathcal{I}_{\check{\mathcal{O}}} \subset \mathcal{U}(\mathfrak{g})$.

- *Definition* (Barbasch-Vogan):

An irr. adm. G -module is called *special unipotent* if



$$\mathrm{Ann}_{\mathcal{U}(\mathfrak{g})}(\pi) = \mathcal{I}_{\check{\mathcal{O}}}.$$

$$\iff \pi \text{ has inf. char. } \chi_{\check{\mathcal{O}}} \text{ and } \mathrm{AV}_{\mathbb{C}}(\pi) = \overline{\mathcal{O}}$$

Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- $\check{\mathcal{O}}$: a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{\mathcal{O}}}$.

Duflo



\rightsquigarrow the maximal primitive ideal $\mathcal{I}_{\check{\mathcal{O}}} \subset \mathcal{U}(\mathfrak{g})$.

- *Definition* (Barbasch-Vogan):

An irr. adm. G -module is called *special unipotent* if

$$\mathrm{Ann}_{\mathcal{U}(\mathfrak{g})}(\pi) = \mathcal{I}_{\check{\mathcal{O}}}.$$

$$\iff \pi \text{ has inf. char. } \chi_{\check{\mathcal{O}}} \text{ and } \mathrm{AV}_{\mathbb{C}}(\pi) = \overline{\mathcal{O}}$$

Spaltenstein

- $\mathrm{Nil}^{\mathrm{special}}(\mathbf{G}) \ni \mathcal{O} :=$ the



dual of $\check{\mathcal{O}}$.

Barbasch-Vogan's definition of special unip. repn. II

G : a real reductive group.

- $\check{\mathcal{O}}$: a nilpotent orbit in $\check{G} \rightsquigarrow$ inf. char. $\chi_{\check{\mathcal{O}}}$.

Duflo



\rightsquigarrow the maximal primitive ideal $\mathcal{I}_{\check{\mathcal{O}}} \subset \mathcal{U}(\mathfrak{g})$.

- *Definition* (Barbasch-Vogan):

An irr. adm. G -module is called *special unipotent* if

$$\mathrm{Ann}_{\mathcal{U}(\mathfrak{g})}(\pi) = \mathcal{I}_{\check{\mathcal{O}}}.$$

$$\iff \pi \text{ has inf. char. } \chi_{\check{\mathcal{O}}} \text{ and } \mathrm{AV}_{\mathbb{C}}(\pi) = \overline{\mathcal{O}}$$

Spaltenstein

- $\mathrm{Nil}^{\mathrm{special}}(\mathbf{G}) \ni \mathcal{O} :=$ the



dual of $\check{\mathcal{O}}$.

- $\mathrm{Unip}_{\check{\mathcal{O}}}(G) := \{ \text{special unipotent repn. attached to } \check{\mathcal{O}} \}.$

Conjecture/Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

Conjecture/Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

- *Philosophy:* $\mathrm{Unip}_{\emptyset}(G)$ = the **building blocks** of the unitary dual.

Conjecture/Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

- *Philosophy:* $\mathrm{Unip}_{\emptyset}(G)$ = the **building blocks** of the unitary dual.

Special unip. repn. are enough in general

E.g., Complex Spin group, by Wong-Zhang.

c.f. Mason-Brown's talk.

Conjecture/Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

- *Philosophy:* $\mathrm{Unip}_{\emptyset}(G)$ = the **building blocks** of the unitary dual.

Special unip. repn. are enough in general

E.g., Complex Spin group, by Wong-Zhang.

c.f. Mason-Brown's talk.

- *Conjecture (1980s):* $\mathrm{Unip}_{\emptyset}(G)$ consists of **unitary** repn.

Conjecture/Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

- *Philosophy:* $\mathrm{Unip}_{\emptyset}(G)$ = the **building blocks** of the unitary dual.

Special unip. repn. are enough in general

E.g., Complex Spin group, by Wong-Zhang.

c.f. Mason-Brown's talk.

- *Conjecture (1980s):* $\mathrm{Unip}_{\emptyset}(G)$ consists of **unitary** repn.

Question 1: Size of $\mathrm{Unip}_{\emptyset}(G)$?

Question 2: Construction of unipotent repn.

Question 3: Character formula?

Conjecture / Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

- *Philosophy:* $\mathrm{Unip}_{\emptyset}(G)$ = the **building blocks** of the unitary dual.

Special unip. repn. are enough in general

E.g., Complex Spin group, by Wong-Zhang.

c.f. Mason-Brown's talk.

- *Conjecture (1980s):* $\mathrm{Unip}_{\emptyset}(G)$ consists of **unitary** repn.

Question 1: Size of $\mathrm{Unip}_{\emptyset}(G)$?

Question 2: Construction of unipotent repn.

Question 3: Character formula?

- *Barbasch-Vogan* 1985: Classification for **complex groups**.

Conjecture / Open problems

- *Major open problem:* Classify the **unitary dual**:

$$\mathrm{Irr}_{\mathrm{unit}}(G) = \{ \text{irr. unitary repn. of } G \}.$$

- *Philosophy:* $\mathrm{Unip}_{\emptyset}(G)$ = the **building blocks** of the unitary dual.

Special unip. repn. are enough in general

E.g., Complex Spin group, by Wong-Zhang.

c.f. Mason-Brown's talk.

- *Conjecture (1980s):* $\mathrm{Unip}_{\emptyset}(G)$ consists of **unitary** repn.

Question 1: Size of $\mathrm{Unip}_{\emptyset}(G)$?

Question 2: Construction of unipotent repn.

Question 3: Character formula?

- *Barbasch-Vogan* 1985: Classification for **complex groups**.

- *Barbasch* 1989: Proved the conj. for **complex classical groups**.

Special unip. repn. of simply conn. classical groups

Theorem (Barbasch-M.-Sun-Zhu)

Suppose G is a simply connected real classical group, i.e. one of the following groups

$$\mathrm{SU}(p, q), \mathrm{Spin}(p, q), \mathrm{Spin}(2n, \mathbb{H}), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \mathrm{Sp}(p, q)$$

Arthur-Barbasch-Vogan's unitarity conj. for special unipotent repn. holds:

Special unip. repn. of simply conn. classical groups

Theorem (Barbasch-M.-Sun-Zhu)

Suppose G is a simply connected real classical group, i.e. one of the following groups


$$\mathrm{SU}(p, q), \mathrm{Spin}(p, q), \mathrm{Spin}(2n, \mathbb{H}), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{Mp}(2n, \mathbb{R}), \mathrm{Sp}(p, q)$$

Arthur-Barbasch-Vogan's unitarity conj. for special unipotent repn. holds:

All *special unipotent repn.* of G are *unitarizable*.

Answer for *exceptional groups*

J. Adams, S. Miller, M. van Leeuwen, and D. A. Vogan



Counting G -module with a specific ann. variety

- Fix inf. char. $\mu \in \mathfrak{h}^*/W$

Counting G -module with a specific ann. variety

- Fix inf. char. $\mu \in \mathfrak{h}^*/W$
- integral Weyl group

$$W(\mu) := \{ w \in W \mid \langle \mu - w\mu, \check{\alpha} \rangle \in \mathbb{Z}, \forall \alpha \in \Delta(\mathfrak{g}, \mathfrak{h}) \}$$

double cell $\mathcal{D} \subset \text{Irr}(W(\mu)) \rightsquigarrow$ the special repn. τ_0

\rightsquigarrow truncated ind. $J_{W(\mu)}^W \tau_0 \xrightarrow{\text{Springer corr.}} \mathcal{O}.$

Counting G -module with a specific ann. variety

- Fix inf. char. $\mu \in \mathfrak{h}^* / W$
- integral Weyl group

$$W(\mu) := \{ w \in W \mid \langle \mu - w\mu, \check{\alpha} \rangle \in \mathbb{Z}, \forall \alpha \in \Delta(\mathfrak{g}, \mathfrak{h}) \}$$

double cell $\mathcal{D} \subset \text{Irr}(W(\mu)) \rightsquigarrow$ the special repn. τ_0

\rightsquigarrow truncated ind. $J_{W(\mu)}^W \tau_0 \xrightarrow{\text{Springer corr.}} \mathcal{O}.$

$$W_\mu = \{ w \in W \mid w\mu = \mu \}.$$

Counting G -module with a specific ann. variety

- Fix inf. char. $\mu \in \mathfrak{h}^* / W$
- integral Weyl group

$$W(\mu) := \{ w \in W \mid \langle \mu - w\mu, \check{\alpha} \rangle \in \mathbb{Z}, \forall \alpha \in \Delta(\mathfrak{g}, \mathfrak{h}) \}$$

double cell $\mathcal{D} \subset \text{Irr}(W(\mu)) \rightsquigarrow$ the special repn. τ_0

\rightsquigarrow truncated ind. $J_{W(\mu)}^W \tau_0 \xrightarrow{\text{Springer corr.}} \mathcal{O}.$

$$W_\mu = \{ w \in W \mid w\mu = \mu \}.$$

- $\mathcal{G}_\mu(G)$: the coherent continuation repn. of $W(\mu)$.

Counting G -module with a specific ann. variety

- Fix inf. char. $\mu \in \mathfrak{h}^* / W$
- integral Weyl group

$$W(\mu) := \{ w \in W \mid \langle \mu - w\mu, \alpha \rangle \in \mathbb{Z}, \forall \alpha \in \Delta(\mathfrak{g}, \mathfrak{h}) \}$$

double cell $\mathcal{D} \subset \text{Irr}(W(\mu)) \rightsquigarrow$ the special repn. τ_0

\rightsquigarrow truncated ind. $J_{W(\mu)}^W \tau_0 \xrightarrow{\text{Springer corr.}} \mathcal{O}$.

$$W_\mu = \{ w \in W \mid w\mu = \mu \}.$$

- $\mathcal{G}_\mu(G)$: the coherent continuation repn. of $W(\mu)$.

Lemma:

$$\begin{aligned} & \# \{ \pi \in \text{Irr}_\mu(\mathfrak{g}, K)(G) \mid \text{AV}_C(\pi) = \overline{\mathcal{O}} \} \\ &= \sum_{\substack{\mathcal{D} \rightsquigarrow \mathcal{O} \\ \tau \in \mathcal{D}}} [\tau : 1_{W_\mu}] \cdot [\tau : \mathcal{G}_\mu(G)] \end{aligned}$$

Counting G -module with a specific ann. variety

- Fix inf. char. $\mu \in \mathfrak{h}^* / W$
- integral Weyl group

$$W(\mu) := \{ w \in W \mid \langle \mu - w\mu, \alpha \rangle \in \mathbb{Z}, \forall \alpha \in \Delta(\mathfrak{g}, \mathfrak{h}) \}$$

double cell $\mathcal{D} \subset \text{Irr}(W(\mu)) \rightsquigarrow$ the special repr. τ_0

\rightsquigarrow truncated ind. $J_{W(\mu)}^W \tau_0 \xrightarrow{\text{Springer corr.}} \mathcal{O}$.

$$W_\mu = \{ w \in W \mid w\mu = \mu \}.$$

- $\mathcal{G}_\mu(G)$: the coherent continuation repr. of $W(\mu)$.

Lemma: If \mathfrak{g} has no E_8 factor, then

$$\begin{aligned} & \# \{ \pi \in \text{Irr}_\mu(\mathfrak{g}, K)(G) \mid \text{AV}_C(\pi) = \overline{\mathcal{O}} \} \\ &= \sum_{\substack{\mathcal{D} \rightsquigarrow \mathcal{O} \\ \tau \in \mathcal{D}}} [\tau : 1_{W_\mu}] \cdot [\tau : \mathcal{G}_\mu(G)] \end{aligned}$$

Applications

- $\text{Unip}_{\check{O}}(\text{SL}_n(\mathbb{H})) = \text{restriction of } \text{Unip}_{\check{O}}(\text{GL}_n(\mathbb{H})).$

Applications

- $\text{Unip}_{\check{O}}(\text{SL}_n(\mathbb{H})) = \text{restriction of } \text{Unip}_{\check{O}}(\text{GL}_n(\mathbb{H})).$
- $\text{Unip}_{\check{O}}(\text{SL}_n(\mathbb{R})) = \text{components in the rest. of } \text{Unip}_{\check{O}}(\text{GL}_n(\mathbb{R})).$

Applications

- $\text{Unip}_{\mathcal{O}}(\text{SL}_n(\mathbb{H})) = \text{restriction of } \text{Unip}_{\mathcal{O}}(\text{GL}_n(\mathbb{H})).$
- $\text{Unip}_{\mathcal{O}}(\text{SL}_n(\mathbb{R})) = \text{components in the rest. of } \text{Unip}_{\mathcal{O}}(\text{GL}_n(\mathbb{R})).$
- $\text{Unip}_{\mathcal{O}}(\text{SU}(p, q)) = \text{rest. of } \text{Unip}_{\mathcal{O}}(\text{det double cover of } \text{U}(p, q)).$

Applications

- $\text{Unip}_{\mathcal{O}}(\text{SL}_n(\mathbb{H})) = \text{restriction of } \text{Unip}_{\mathcal{O}}(\text{GL}_n(\mathbb{H})).$
- $\text{Unip}_{\mathcal{O}}(\text{SL}_n(\mathbb{R})) = \text{components in the rest. of } \text{Unip}_{\mathcal{O}}(\text{GL}_n(\mathbb{R})).$
- $\text{Unip}_{\mathcal{O}}(\text{SU}(p, q)) = \text{rest. of } \text{Unip}_{\mathcal{O}}(\text{det double cover of } \text{U}(p, q)).$
- genuine special unipotent repn. of $\text{Spin}(p, q)$ are some obvious irreducibly parabolically induced module.

Nilpotent orbits with “good/bad parity”

- Bad parity (must occur with even multiplicity in $\check{\mathcal{O}}$):


$$\begin{cases} \text{even number,} & \text{when } \check{G} \text{ is type } B \text{ or } D \\ \text{odd number,} & \text{when } \check{G} \text{ is type } C \end{cases}$$

- $\check{\mathcal{O}}$ has “good parity” if $\check{\mathcal{O}}$ only contains

$$\begin{cases} \text{odd rows,} & \text{when } \check{G} \text{ is type } B \text{ or } D \\ \text{even rows,} & \text{when } \check{G} \text{ is type } C \end{cases}$$


- $\lambda_{\check{\mathcal{O}}}$ is integral.

\mathcal{O}



$\mathrm{Sp}(14, \mathbb{C})$

$\check{\mathcal{O}}$



$\mathrm{SO}(15, \mathbb{C})$

Reduction to the “good parity”

- Consider $G = \mathrm{Sp}(2n, \mathbb{R})$.
- $\check{\mathcal{O}}$ decompose into two parts $\check{\mathcal{O}}_g$ (good parity) and $\check{\mathcal{O}}_b$ (bad parity).
- Assume $\check{\mathcal{O}}_b = \{r_1, r_1, \dots, r_k, r_k\}$.

Reduction to the “good parity”

- Consider $G = \mathrm{Sp}(2n, \mathbb{R})$.
- $\check{\mathcal{O}}$ decompose into two parts $\check{\mathcal{O}}_g$ (good parity) and $\check{\mathcal{O}}_b$ (bad parity).
- Assume $\check{\mathcal{O}}_b = \{r_1, r_1, \dots, r_k, r_k\}$.

Theorem Let $\check{\mathcal{O}}'_b = \{r_1, \dots, r_k\} \in \mathrm{Nil}_{\mathrm{GL}}$.

$$\begin{aligned} \mathrm{Unip}_{\check{\mathcal{O}}'_b}(\mathrm{GL}) \times \mathrm{Unip}_{\check{\mathcal{O}}_g}(\mathrm{Sp}) &\xrightarrow{\text{bij.}} \mathrm{Unip}_{\check{\mathcal{O}}}(\mathrm{Sp}) \\ (\pi', \pi_0) &\mapsto \mathrm{Ind}_{\mathrm{GL}_{|\check{\mathcal{O}}'_b|} \times \mathrm{Sp}(2n_0, \mathbb{R}) \times U}^{\mathrm{Sp}(2n, \mathbb{R})} \pi' \otimes \pi_0 \end{aligned}$$

$$\mathrm{Unip}_{\check{\mathcal{O}}'_b}(\mathrm{GL}) = \left\{ \mathrm{Ind}_{j=1}^k \otimes \mathrm{sgn}_{\mathrm{GL}(r_j, \mathbb{R})}^{\epsilon_j} \mid \epsilon_j \in \mathbb{Z}/2\mathbb{Z} \right\}$$

Reduction to the “good parity”

- Consider $G = \mathrm{Sp}(2n, \mathbb{R})$.
- $\check{\mathcal{O}}$ decompose into two parts $\check{\mathcal{O}}_g$ (good parity) and $\check{\mathcal{O}}_b$ (bad parity).
- Assume $\check{\mathcal{O}}_b = \{r_1, r_1, \dots, r_k, r_k\}$.

Theorem Let $\check{\mathcal{O}}'_b = \{r_1, \dots, r_k\} \in \mathrm{Nil}_{\mathrm{GL}}$.

$$\begin{aligned} \mathrm{Unip}_{\check{\mathcal{O}}'_b}(\mathrm{GL}) \times \mathrm{Unip}_{\check{\mathcal{O}}_g}(\mathrm{Sp}) &\xrightarrow{\text{bij.}} \mathrm{Unip}_{\check{\mathcal{O}}}(\mathrm{Sp}) \\ (\pi', \pi_0) &\mapsto \mathrm{Ind}_{\mathrm{GL}_{|\check{\mathcal{O}}'_b|} \times \mathrm{Sp}(2n_0, \mathbb{R}) \ltimes U}^{\mathrm{Sp}(2n, \mathbb{R})} \pi' \otimes \pi_0 \end{aligned}$$

$$\mathrm{Unip}_{\check{\mathcal{O}}'_b}(\mathrm{GL}) = \left\{ \mathrm{Ind}_{j=1}^k \otimes \mathrm{sgn}_{\mathrm{GL}(r_j, \mathbb{R})}^{\epsilon_j} \mid \epsilon_j \in \mathbb{Z}/2\mathbb{Z} \right\}$$

- Use *theta correspondence* to study $\mathrm{Unip}_{\check{\mathcal{O}}_g}(G)$.
- We assume $\check{\mathcal{O}}$ has **good parity** from now on.

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

$$W(\chi_{\check{\mathcal{O}}}) = S_n \ltimes \{\pm 1\}^n,$$

$$\mathcal{G}_{\rho_G}(G) = \sum_{\substack{b,c,d,r, \\ \sigma \in \hat{S}_r}} \mathrm{Ind}_{S_r \times W_{2b} \times W_c \times W_d}^{W_n} \mathrm{sgn} \otimes (\sigma \times \sigma) \otimes \mathbf{1} \otimes \mathbf{1}.$$



$$\text{max. torus } \mathbb{C}^b \times \mathrm{U}(1)^r \times \mathbb{R}^{c+d}$$

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

$$W(\chi_{\check{\mathcal{O}}}) = S_n \ltimes \{\pm 1\}^n,$$

$$\mathcal{G}_{\rho_G}(G) = \sum_{\substack{b,c,d,r, \\ \sigma \in \hat{S}_r}} \mathrm{Ind}_{S_r \times W_{2b} \times W_c \times W_d}^{W_n} \mathrm{sgn} \otimes (\sigma \times \sigma) \otimes \mathbf{1} \otimes \mathbf{1}.$$



$$\text{max. torus } \mathbb{C}^b \times \mathrm{U}(1)^r \times \mathbb{R}^{c+d}$$

$$\check{\mathcal{O}} \rightsquigarrow \text{special repn. } \tau = \text{Springer}(d_{BV}(\check{\mathcal{O}}))$$

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

$$W(\chi_{\check{\mathcal{O}}}) = S_n \ltimes \{ \pm 1 \}^n,$$

$$\mathcal{G}_{\rho_G}(G) = \sum_{\substack{b,c,d,r, \\ \sigma \in \hat{S}_r}} \mathrm{Ind}_{S_r \times W_{2b} \times W_c \times W_d}^{W_n} \mathrm{sgn} \otimes (\sigma \times \sigma) \otimes \mathbf{1} \otimes \mathbf{1}.$$



$$\text{max. torus } \mathbb{C}^b \times \mathrm{U}(1)^r \times \mathbb{R}^{c+d}$$

$$\check{\mathcal{O}} \rightsquigarrow \text{special repn. } \tau = \text{Springer}(d_{BV}(\check{\mathcal{O}}))$$

$$\text{Lusztig left cell } \mathcal{C} = \{ \tau_1, \dots, \tau_{2l} \} \text{ containing } \tau$$

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

$$W(\chi_{\check{\mathcal{O}}}) = S_n \ltimes \{ \pm 1 \}^n,$$

$$\mathcal{G}_{\rho_G}(G) = \sum_{\substack{b,c,d,r, \\ \sigma \in \hat{S}_r}} \mathrm{Ind}_{S_r \times W_{2b} \times W_c \times W_d}^{W_n} \mathrm{sgn} \otimes (\sigma \times \sigma) \otimes \mathbf{1} \otimes \mathbf{1}.$$



$$\text{max. torus } \mathbb{C}^b \times \mathrm{U}(1)^r \times \mathbb{R}^{c+d}$$

$$\check{\mathcal{O}} \rightsquigarrow \text{special repr. } \tau = \text{Springer}(d_{BV}(\check{\mathcal{O}}))$$

$$\text{Lusztig left cell } \mathcal{C} = \{ \tau_1, \dots, \tau_{2l} \} \text{ containing } \tau$$

- $[\tau_i : 1_{W_{\lambda_{\check{\mathcal{O}}}}}] = 1$ and $[\tau_i : \mathcal{G}_{\rho_G}(G)] = [\tau : \mathcal{G}_{\rho_G}(G)]$

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

$$W(\chi_{\check{\mathcal{O}}}) = S_n \ltimes \{\pm 1\}^n,$$

$$\mathcal{G}_{\rho_G}(G) = \sum_{\substack{b,c,d,r, \\ \sigma \in \hat{S}_r}} \mathrm{Ind}_{S_r \times W_{2b} \times W_c \times W_d}^{W_n} \mathrm{sgn} \otimes (\sigma \times \sigma) \otimes \mathbf{1} \otimes \mathbf{1}.$$



$$\text{max. torus } \mathbb{C}^b \times \mathrm{U}(1)^r \times \mathbb{R}^{c+d}$$

$$\check{\mathcal{O}} \rightsquigarrow \text{special repr. } \tau = \text{Springer}(d_{BV}(\check{\mathcal{O}}))$$

$$\text{Lusztig left cell } \mathcal{C} = \{\tau_1, \dots, \tau_{2^l}\} \text{ containing } \tau$$

- $[\tau_i : 1_{W_{\lambda_{\check{\mathcal{O}}}}}] = 1$ and $[\tau_i : \mathcal{G}_{\rho_G}(G)] = [\tau : \mathcal{G}_{\rho_G}(G)]$
 $\# \mathrm{Unip}_{\check{\mathcal{O}}}(G) = 2^l \cdot [\tau : \mathcal{G}_{\rho_G}(G)]$

Counting unipotent representations I

- Example: $G = \mathrm{Sp}(2n, \mathbb{R})$ and $\chi_{\check{\mathcal{O}}} \in \rho_G + \text{weight lattice}$.

$$W(\chi_{\check{\mathcal{O}}}) = S_n \ltimes \{\pm 1\}^n,$$

$$\mathcal{G}_{\rho_G}(G) = \sum_{\substack{b,c,d,r, \\ \sigma \in \hat{S}_r}} \mathrm{Ind}_{S_r \times W_{2b} \times W_c \times W_d}^{W_n} \mathrm{sgn} \otimes (\sigma \times \sigma) \otimes \mathbf{1} \otimes \mathbf{1}.$$



$$\text{max. torus } \mathbb{C}^b \times \mathrm{U}(1)^r \times \mathbb{R}^{c+d}$$

$$\check{\mathcal{O}} \rightsquigarrow \text{special repr. } \tau = \text{Springer}(d_{BV}(\check{\mathcal{O}}))$$


$$\text{Lusztig left cell } \mathcal{C} = \{\tau_1, \dots, \tau_{2^l}\} \text{ containing } \tau$$

- $[\tau_i : 1_{W_{\lambda_{\check{\mathcal{O}}}}}] = 1$ and $[\tau_i : \mathcal{G}_{\rho_G}(G)] = [\tau : \mathcal{G}_{\rho_G}(G)]$
 $\# \text{Unip}_{\check{\mathcal{O}}}(G) = 2^l \cdot [\tau : \mathcal{G}_{\rho_G}(G)]$


- $[\tau : \mathcal{G}_{\rho_G}(G)]$ is counted by painted bi-partitions $\text{PBP}(\check{\mathcal{O}})$.

Example of PBP

$$\check{\mathcal{O}} = [7, 5, 5, 5, 3, 1, 1] =$$




$\text{PBP}_{\check{\mathcal{O}}}(\text{Sp}(2n, \mathbb{R}))$



\times









\mapsto

Associated character











\cup

Inductive structure of nilpotent orbits

$\widehat{\mathbf{G}}_i$	$\mathrm{SO}(15, \mathbb{C})$	$\mathrm{O}(10, \mathbb{C})$	$\mathrm{SO}(7, \mathbb{C})$	$\mathrm{O}(4, \mathbb{C})$
$\check{\mathcal{O}}_i$				
\mathcal{O}_i				
\mathbf{G}_i	$\mathrm{Sp}(14, \mathbb{C})$	$\mathrm{O}(10, \mathbb{C})$	$\mathrm{Sp}(6, \mathbb{C})$	$\mathrm{O}(4, \mathbb{C})$

Inductive structure of nilpotent orbits

$\widehat{\mathbf{G}}_i$	$\mathrm{SO}(15, \mathbb{C})$	$\mathrm{O}(10, \mathbb{C})$	$\mathrm{SO}(7, \mathbb{C})$	$\mathrm{O}(4, \mathbb{C})$
$\check{\mathcal{O}}_i$				
\mathcal{O}_i				
\mathbf{G}_i	$\mathrm{Sp}(14, \mathbb{C})$	$\mathrm{O}(10, \mathbb{C})$	$\mathrm{Sp}(6, \mathbb{C})$	$\mathrm{O}(4, \mathbb{C})$

Kraft



Procesi





resolution of singularities of nilpotent orbit closures.




Example of descent sequences

\mathbf{G}_i^\vee $\mathrm{SO}(15, \mathbb{C})$ $\mathrm{O}(10, \mathbb{C})$ $\mathrm{SO}(7, \mathbb{C})$ $\mathrm{O}(4, \mathbb{C})$

$\check{\mathcal{O}}_i$



\mathcal{O}_i



G_i









$\mathrm{Sp}(14, \mathbb{R})$

$\mathrm{O}(4, 6)$

$\mathrm{Sp}(6, \mathbb{R})$

$\mathrm{O}(2, 2)$

Example of descent sequences

G_i^\vee	$\mathrm{SO}(15, \mathbb{C})$	$\mathrm{O}(10, \mathbb{C})$	$\mathrm{SO}(7, \mathbb{C})$	$\mathrm{O}(4, \mathbb{C})$
$\check{\mathcal{O}}_i$				
\mathcal{O}_i				
G_i	$\mathrm{Sp}(14, \mathbb{R})$	$\mathrm{O}(4, 6)$	$\mathrm{Sp}(6, \mathbb{R})$	$\mathrm{O}(2, 2)$

Ohta's resolution of singularities of a nilpotent orbit closure in symmetric pairs.

Construction of elements in $\text{Unip}_{\mathfrak{O}}(G)$

- $\chi = \bigotimes_{j=0}^{2a} \chi_j$, a 1-dim repn. of $\prod_{j=0}^{2a} G_j$.
- $\chi_j \in \{\mathbf{1}, \text{sgn}^{+,-}, \text{sgn}^{-,+}, \det\}$
- Define a smooth repn. of $G = G_{2a}$ (the symplectic group).

$$\pi_{\chi} := (\omega_{G_{2a}, G_{2a-1}} \hat{\otimes} \omega_{G_{2a-1}, G_{2a-2}} \hat{\otimes} \cdots \hat{\otimes} \omega_{G_1, G_0} \otimes \chi)_{G_{2a-1} \times G_{2a-2} \times \cdots \times G_0}$$

Construction of elements in $\text{Unip}_{\check{\mathcal{O}}}(G)$

- $\chi = \bigotimes_{j=0}^{2a} \chi_j$, a 1-dim repn. of $\prod_{j=0}^{2a} G_j$.
- $\chi_j \in \{\mathbf{1}, \text{sgn}^{+,-}, \text{sgn}^{-,+}, \det\}$
- Define a smooth repn. of $G = G_{2a}$ (the symplectic group).

$$\pi_{\chi} := (\omega_{G_{2a}, G_{2a-1}} \hat{\otimes} \omega_{G_{2a-1}, G_{2a-2}} \hat{\otimes} \cdots \hat{\otimes} \omega_{G_1, G_0} \otimes \chi)_{G_{2a-1} \times G_{2a-2} \times \cdots \times G_0}$$

Theorem (Barbasch-M.-Sun-Zhu)

Suppose $\check{\mathcal{O}}$ is an orbit with good parity. Then

- *either $\pi_{\chi} = 0$ or*

Construction of elements in $\text{Unip}_{\check{\mathcal{O}}}(G)$

- $\chi = \bigotimes_{j=0}^{2a} \chi_j$, a 1-dim repn. of $\prod_{j=0}^{2a} G_j$.
- $\chi_j \in \{\mathbf{1}, \text{sgn}^{+, -}, \text{sgn}^{-, +}, \det\}$
- Define a smooth repn. of $G = G_{2a}$ (the symplectic group).

$$\pi_{\chi} := (\omega_{G_{2a}, G_{2a-1}} \hat{\otimes} \omega_{G_{2a-1}, G_{2a-2}} \hat{\otimes} \cdots \hat{\otimes} \omega_{G_1, G_0} \otimes \chi)_{G_{2a-1} \times G_{2a-2} \times \cdots \times G_0}$$

Theorem (Barbasch-M.-Sun-Zhu)

Suppose $\check{\mathcal{O}}$ is an orbit with good parity. Then

- either $\pi_{\chi} = 0$ or
- $\pi_{\chi} \in \text{Unip}_{\check{\mathcal{O}}}(G)$ and unitarizable.

Construction of elements in $\text{Unip}_{\check{\mathcal{O}}}(G)$

- $\chi = \bigotimes_{j=0}^{2a} \chi_j$, a 1-dim repn. of $\prod_{j=0}^{2a} G_j$.
- $\chi_j \in \{\mathbf{1}, \text{sgn}^{+, -}, \text{sgn}^{-, +}, \det\}$
- Define a smooth repn. of $G = G_{2a}$ (the symplectic group).

$$\pi_{\chi} := (\omega_{G_{2a}, G_{2a-1}} \hat{\otimes} \omega_{G_{2a-1}, G_{2a-2}} \hat{\otimes} \cdots \hat{\otimes} \omega_{G_1, G_0} \otimes \chi)_{G_{2a-1} \times G_{2a-2} \times \cdots \times G_0}$$

Theorem (Barbasch-M.-Sun-Zhu)

Suppose $\check{\mathcal{O}}$ is an orbit with good parity. Then

- either $\pi_{\chi} = 0$ or
- $\pi_{\chi} \in \text{Unip}_{\check{\mathcal{O}}}(G)$ and unitarizable.
- Moreover,

$$\text{Unip}_{\check{\mathcal{O}}}(G) = \{ \pi_{\chi} \mid \pi_{\chi} \neq 0 \}.$$

Example: Coincidences of theta lifting

Lift to $G = \mathrm{Sp}(6, \mathbb{R})$ from real forms of $\mathbf{G} = \mathrm{O}(4, \mathbb{C})$.

$\check{\mathcal{O}} = 3^2 1^1$ and $\mathcal{O} = 2^3$.

		$\mathrm{Sp}(6, \mathbb{R})$	
$\mathrm{O}(4, 0)$		$\theta(\mathrm{sgn}^{+, -})$	
$\mathrm{O}(3, 1)$	$\theta(\mathbf{1})$	$\theta(\mathrm{sgn}^{+, -})$	$\theta(\mathrm{sgn}^{-, +})$
$\mathrm{O}(2, 2)$	$\theta(\mathbf{1})$	$\theta(\mathrm{sgn}^{+, -})$	$\theta(\mathrm{sgn}^{-, +})$
$\mathrm{O}(1, 3)$	$\theta(\mathbf{1})$	$\theta(\mathrm{sgn}^{+, -})$	$\theta(\mathrm{sgn}^{-, +})$
$\mathrm{O}(0, 4)$			$\theta(\mathrm{sgn}^{-, +})$

Matching unipotent representations with PBP

- $\text{PBP}(\check{\mathcal{O}})$ is complicate.

Matching unipotent representations with PBP

- $\text{PBP}(\check{\mathcal{O}})$ is complicate.
- $\text{LS}(\check{\mathcal{O}}) = \{ \text{AC}(\pi_\chi) \}$ is also complicate.

Matching unipotent representations with PBP

- $\text{PBP}(\check{\mathcal{O}})$ is complicate.
- $\text{LS}(\check{\mathcal{O}}) = \{ \text{AC}(\pi_\chi) \}$ is also complicate.
- Proof of Exhaustion

Matching unipotent representations with PBP

- $\text{PBP}(\check{\mathcal{O}})$ is complicate.
- $\text{LS}(\check{\mathcal{O}}) = \{ \text{AC}(\pi_\chi) \}$ is also complicate.
- **Proof of Exhaustion**

Define descent of painted bi-part., **compatible with the theta!**

$$\begin{array}{ccccc}
 \text{LS}(\check{\mathcal{O}}) & \xleftarrow{\text{AC}} & \text{PBP}^{\text{ext}}(\check{\mathcal{O}}) & \longleftrightarrow & \bigcup \text{Unip}_{\check{\mathcal{O}}}(G) \\
 \textcolor{red}{\vartheta}^{\text{geo}} \uparrow & & \nabla \downarrow & & \uparrow \theta \\
 \text{LS}(\check{\mathcal{O}}') & \xleftarrow{\text{AC}} & \text{PBP}^{\text{ext}}(\check{\mathcal{O}}') & \longleftrightarrow & \bigcup \text{Unip}_{\check{\mathcal{O}}'}(G')
 \end{array}$$

Matching unipotent representations with PBP

- $\text{PBP}(\check{\mathcal{O}})$ is complicate.
- $\text{LS}(\check{\mathcal{O}}) = \{ \text{AC}(\pi_\chi) \}$ is also complicate.
- **Proof of Exhaustion**

Define descent of painted bi-part., **compatible with the theta!**

$$\begin{array}{ccccc}
 \text{LS}(\check{\mathcal{O}}) & \xleftarrow{\text{AC}} & \text{PBP}^{\text{ext}}(\check{\mathcal{O}}) & \longleftrightarrow & \bigcup \text{Unip}_{\check{\mathcal{O}}}(G) \\
 \textcolor{red}{\vartheta}^{\text{geo}} \uparrow & & \nabla \downarrow & & \uparrow \theta \\
 \text{LS}(\check{\mathcal{O}}') & \xleftarrow{\text{AC}} & \text{PBP}^{\text{ext}}(\check{\mathcal{O}}') & \longleftrightarrow & \bigcup \text{Unip}_{\check{\mathcal{O}}'}(G')
 \end{array}$$

$$\pi_\tau := \Theta(\pi_{\nabla(\tau)} \otimes \chi'_\tau) \otimes \chi_\tau$$

Matching unipotent representations with PBP

- $\text{PBP}(\check{\mathcal{O}})$ is complicate.
- $\text{LS}(\check{\mathcal{O}}) = \{ \text{AC}(\pi_\chi) \}$ is also complicate.
- **Proof of Exhaustion**

Define descent of painted bi-part., **compatible with the theta!**

$$\begin{array}{ccccc}
 \text{LS}(\check{\mathcal{O}}) & \xleftarrow{\text{AC}} & \text{PBP}^{\text{ext}}(\check{\mathcal{O}}) & \longleftrightarrow & \bigcup \text{Unip}_{\check{\mathcal{O}}}(G) \\
 \uparrow \scriptstyle \textcolor{red}{\vartheta}^{\text{geo}} & & \downarrow \nabla & & \uparrow \theta \\
 \text{LS}(\check{\mathcal{O}}') & \xleftarrow{\text{AC}} & \text{PBP}^{\text{ext}}(\check{\mathcal{O}}') & \longleftrightarrow & \bigcup \text{Unip}_{\check{\mathcal{O}}'}(G')
 \end{array}$$

$$\pi_\tau := \Theta(\pi_{\nabla(\tau)} \otimes \chi'_\tau) \otimes \chi_\tau$$

- The **injectivity** of theta lifting is crucial!

About the proof

- θ -lifting $\approx \sqrt{\text{parabolic induction}}$
- Structure of degenerate principle series (Algebra).
- Estimation of matrix coeff. integrals (Analysis) \rightsquigarrow unitarity

About the proof

- θ -lifting $\approx \sqrt{\text{parabolic induction}}$
- Structure of degenerate principle series (Algebra).
- Estimation of matrix coeff. integrals (Analysis) \rightsquigarrow unitarity
- Sharp formula of Asso. Cycles: **lower bound=upper bound**
 - *Algebra+Analysis*: \rightsquigarrow lower bound.
 - *Algebraic Geometry*: double fibration of moment maps
 \rightsquigarrow upper bound
- Exhaustion: **lower bound=upper bound** (Combinatorics)
 - Character theory (Kazhdan-Lusztig-Vogan theory)
 \rightsquigarrow upper bound by counting “tableaux”.
 - Asso. Cycle+Injectivity of θ \rightsquigarrow lower bound.

- *Definition for metaplectic groups*

On the notion of metaplectic Barbasch-Vogan duality

<https://arxiv.org/abs/2010.16089>

- *Construction and unitarity using θ -lifting*

Special unipotent representations: orthogonal and symplectic groups

<https://arxiv.org/abs/1712.05552>

- *Counting unipotent representations*

<https://arxiv.org/abs/2205.05266>

Thank you for your attention!

